


INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C12N 15/86, 15/12, A61K 48/00		A2	(11) International Publication Number: WO 94/12649
			(43) International Publication Date: 9 June 1994 (09.06.94)
(21) International Application Number: PCT/US93/11667			(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(22) International Filing Date: 2 December 1993 (02.12.93)			
(30) Priority Data:			
07/985,478	3 December 1992 (03.12.92)	US	
08/130,682	1 October 1993 (01.10.93)	US	
08/136,742	13 October 1993 (13.10.93)	US	
(71) Applicant: GENZYME CORPORATION [US/US]; One Kendall Square, Cambridge, MA 02139 (US).			
(72) Inventors: GREGORY, Richard, J.; 4789 Gateshead Road, Carlsbad, CA 92008 (US). ARMENTANO, Donna; 33 Carver Road, Watertown, MA 02172 (US). COUTURE, Larry, A.; 67 Circle Drive, Framingham, MA 01701 (US). SMITH, Alan, E.; 88 Cleveland Road, Wellesley, MA 02181 (US).			
(74) Agents: HANLEY, Elizabeth, A. et al.; Lahive & Cockfield, 60 State Street, Boston, MA 02109 (US).			

(54) Title: GENE THERAPY FOR CYSTIC FIBROSIS

(57) Abstract

Gene Therapy vectors, which are especially useful for cystic fibrosis, and methods for using the vectors are disclosed. In preferred embodiments, the vectors are adenovirus-based. Advantages of adenovirus-based vectors for gene therapy are that they appear to be relatively safe and can be manipulated to encode the desired gene product and at the same time are inactivated in terms of their ability to replicate in a normal lytic viral life cycle. Additionally, adenovirus has a natural tropism for airway epithelia. Therefore, adenovirus-based vectors are particularly preferred for respiratory gene therapy applications such as gene therapy for cystic fibrosis. In one embodiment, the adenovirus-based gene therapy vector comprises an adenovirus 2 serotype genome in which the E1a and E1b regions of the genome, which are involved in early stages of viral replication have been deleted and replaced by genetic material of interest (e.g., DNA encoding the cystic fibrosis transmembrane regulator protein). In another embodiment, the adenovirus-based therapy vector is a pseudo-adenovirus (PAV). PAVs contain no potentially harmful viral genes, have a theoretical capacity for foreign material of nearly 36 kb, may be produced in reasonably high titers and maintain the tropism of the parent adenovirus for dividing and non-dividing human target cell types.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

GENE THERAPY FOR CYSTIC FIBROSIS

Related Applications

This application is a continuation-in-part application of United States Serial Number 08/130,682, filed on October 1, 1993 which is a continuation-in-part application of United States Serial Number 07/985,478, filed on December 2, 1992, which is a continuation-in-part application of United States Serial Number 07/613,592, filed on November 15, 1990, which is in turn a continuation-in-part application of United States Serial Number 07/589,295, filed on September 27, 1990, which is itself a continuation-in-part application of United States Serial Number 07/488,307, filed on March 5, 1990. The contents of all of the above copending patent applications are incorporated herein by reference. Definitions of language or terms not provided in the present application are the same as those set forth in the copending applications. Any reagents or materials used in the examples of the present application whose source is not expressly identified also is the same as those described in the copending application, e.g., $\Delta F508$ CFTR gene and CFTR antibodies.

Background of the Invention

Cystic Fibrosis (CF) is the most common fatal genetic disease in humans (Boat, T.F. et al. in *The Metabolic Basis of Inherited Diseases* (Scriver, C.R. et al. eds., McGraw-Hill, New York (1989)). Approximately one in every 2,500 infants in the United States is born with the disease. At the present time, there are approximately 30,000 CF patients in the United States. Despite current standard therapy, the median age of survival is only 26 years. Disease of the pulmonary airways is the major cause of morbidity and is responsible for 95% of the mortality. The first manifestation of lung disease is often a cough, followed by progressive dyspnea. Tenacious sputum becomes purulent because of colonization of *Staphylococcus* and then with *Pseudomonas*. Chronic bronchitis and bronchiectasis can be partially treated with current therapy, but the course is punctuated by increasingly frequent exacerbations of the pulmonary disease. As the disease progresses, the patient's activity is progressively limited. End-stage lung disease is heralded by increasing hypoxemia, pulmonary hypertension, and cor pulmonale.

The upper airways of the nose and sinuses are also involved in CF. Most patients with CF develop chronic sinusitis. Nasal polyps occur in 15-20% of patients and are common by the second decade of life. Gastrointestinal problems are also frequent in CF; infants may suffer meconium ileus. Exocrine pancreatic insufficiency, which produces symptoms of malabsorption, is present in the large majority of patients with CF. Males are almost uniformly infertile and fertility is decreased in females.

Based on both genetic and molecular analyses, a gene associated with CF was isolated as part of 21 individual cDNA clones and its protein product predicted (Kerem, B.S. et al. (1989) *Science* 245:1073-1080; Riordan, J.R. et al. (1989) *Science* 245:1066-1073;

Rommens, J.M. et al. (1989) *Science* 245:1059-1065)). United States Serial Number 07/488,307 describes the construction of the gene into a continuous strand, expression of the gene as a functional protein and confirmation that mutations of the gene are responsible for CF. (See also Gregory, R.J. et al. (1990) *Nature* 347:382-386; Rich, D.P. et al. (1990) *Nature* 347:358-362). The co-pending patent application also discloses experiments which show that proteins expressed from wild type but not a mutant version of the cDNA complemented the defect in the cAMP regulated chloride channel shown previously to be characteristic of CF.

The protein product of the CF associated gene is called the cystic fibrosis transmembrane conductance regulator (CFTR) (Riordan, J.R. et al. (1989) *Science* 245:1066-1073). CFTR is a protein of approximately 1480 amino acids made up of two repeated elements, each comprising six transmembrane segments and a nucleotide binding domain. The two repeats are separated by a large, polar, so-called R-domain containing multiple potential phosphorylation sites. Based on its predicted domain structure, CFTR is a member of a class of related proteins which includes the multi-drug resistance (MDR) or P-glycoprotein, bovine adenylyl cyclase, the yeast STE6 protein as well as several bacterial amino acid transport proteins (Riordan, J.R. et al. (1989) *Science* 245:1066-1073; Hyde, S.C. et al. (1990) *Nature* 346:362-365). Proteins in this group, characteristically, are involved in pumping molecules into or out of cells.

CFTR has been postulated to regulate the outward flow of anions from epithelial cells in response to phosphorylation by cyclic AMP-dependent protein kinase or protein kinase C (Riordan, J.R. et al. (1989) *Science* 245:1066-1073; Welsh, 1986; Frizzell, R.A. et al. (1986) *Science* 233:558-560; Welsh, M.J. and Liedtke, C.M. (1986) *Nature* 322:467; Li, M. et al. (1988) *Nature* 331:358-360; Huang, T-C. et al. (1989) *Science* 244:1351-1353).

Sequence analysis of the CFTR gene of CF chromosomes has revealed a variety of mutations (Cutting, G.R. et al. (1990) *Nature* 346:366-369; Dean, M. et al. (1990) *Cell* 61:863-870; and Kerem, B-S. et al. (1989) *Science* 245:1073-1080; Kerem, B-S. et al. (1990) *Proc. Natl. Acad. Sci. USA* 87:8447-8451). Population studies have indicated that the most common CF mutation, a deletion of the 3 nucleotides that encode phenylalanine at position 508 of the CFTR amino acid sequence ($\Delta F508$), is associated with approximately 70% of the cases of cystic fibrosis. This mutation results in the failure of an epithelial cell chloride channel to respond to cAMP (Frizzell R.A. et al. (1986) *Science* 233:558-560; Welsh, M.J. (1986) *Science* 232:1648-1650.; Li, M. et al. (1988) *Nature* 331:358-360; Quinton, P.M. (1989) *Clin. Chem.* 35:726-730). In airway cells, this leads to an imbalance in ion and fluid transport. It is widely believed that this causes abnormal mucus secretion, and ultimately results in pulmonary infection and epithelial cell damage.

Studies on the biosynthesis (Cheng, S.H. et al. (1990) *Cell* 63:827-834; Gregory, R.J. et al. (1991) *Mol. Cell Biol.* 11:3886-3893) and localization (Denning, G.M. et al. (1992) *J. Cell Biol.* 118:551-559) of CFTR $\Delta F508$, as well as other CFTR mutants, indicate that many CFTR mutant proteins are not processed correctly and, as a result, are not delivered to the

plasma membrane (Gregory, R.J. et al. (1991) *Mol. Cell Biol.* 11:3886-3893). These conclusions are consistent with earlier functional studies which failed to detect cAMP-stimulated Cl^- channels in cells expressing CFTR ΔF508 (Rich, D.P. et al. (1990) *Nature* 347:358-363; Anderson, M.P. et al. (1991) *Science* 251:679-682).

- 5 To date, the primary objectives of treatment for CF have been to control infection, promote mucus clearance, and improve nutrition (Boat, T.F. et al. in *The Metabolic Basis of Inherited Diseases* (Scriver, C.R. et al. eds., McGraw-Hill, New York (1989)). Intensive antibiotic use and a program of postural drainage with chest percussion are the mainstays of therapy. However, as the disease progresses, frequent hospitalizations are required.
- 10 Nutritional regimens include pancreatic enzymes and fat-soluble vitamins. Bronchodilators are used at times. Corticosteroids have been used to reduce inflammation, but they may produce significant adverse effects and their benefits are not certain. In extreme cases, lung transplantation is sometimes attempted (Marshall, S. et al. (1990) *Chest* 98:1488).

- Most efforts to develop new therapies for CF have focused on the pulmonary
- 15 complications. Because CF mucus consists of a high concentration of DNA, derived from lysed neutrophils, one approach has been to develop recombinant human DNase (Shak, S. et al. (1990) *Proc. Natl. Sci. Acad USA* 87:9188). Preliminary reports suggest that aerosolized enzyme may be effective in reducing the viscosity of mucus. This could be helpful in clearing the airways of obstruction and perhaps in reducing infections. In an attempt to limit
- 20 damage caused by an excess of neutrophil derived elastase, protease inhibitors have been tested. For example, alpha-1-antitrypsin purified from human plasma has been aerosolized to deliver enzyme activity to lungs of CF patients (McElvaney, N. et al. (1991) *The Lancet* 337:392). Another approach would be the use of agents to inhibit the action of oxidants derived from neutrophils. Although biochemical parameters have been successfully
- 25 measured, the long term beneficial effects of these treatments have not been established.

- Using a different rationale, other investigators have attempted to use pharmacological agents to reverse the abnormally decreased chloride secretion and increased sodium absorption in CF airways. Defective electrolyte transport by airway epithelia is thought to alter the composition of the respiratory secretions and mucus (Boat, T.F. et al. in *The*
- 30 *Metabolic Basis of Inherited Diseases* (Scriver, C.R. et al. eds., McGraw-Hill, New York (1989); Quinton, P.M. (1990) *FASEB J.* 4:2709-2717). Hence, pharmacological treatments aimed at correcting the abnormalities in electrolyte transport could be beneficial. Trials are in progress with aerosolized versions of the drug amiloride; amiloride is a diuretic that inhibits sodium channels, thereby inhibiting sodium absorption. Initial results indicate that the drug
- 35 is safe and suggest a slight change in the rate of disease progression, as measured by lung function tests (Knowles, M. et al. (1990) *N. Eng. J. Med.* 322: 1189-1194; App, E. (1990) *Am. Rev. Respir. Dis.* 141:605). Nucleotides, such as ATP or UTP, stimulate purinergic receptors in the airway epithelium. As a result, they open a class of chloride channel that is different from CFTR chloride channels. *In vitro* studies indicate that ATP and UTP can stimulate

chloride secretion (Knowles, M. et al. (1991) *N. Eng. J. Med.* 325:533). Preliminary trials to test the ability of nucleotides to stimulate secretion *in vivo*, and thereby correct the electrolyte transport abnormalities are underway.

5 Despite progress in therapy, cystic fibrosis remains a lethal disease, and no current therapy treats the basic defect. However, two general approaches may prove feasible. These are: 1) protein replacement therapy to deliver the wild type protein to patients to augment their defective protein, and; 2) gene replacement therapy to deliver wild type copies of the CF associated gene. Since the most life threatening manifestations of CF involve pulmonary complications, epithelial cells of the upper airways are appropriate target cells for therapy.

10 The feasibility of gene therapy has been established by introducing a wild type cDNA into epithelial cells from a CF patient and demonstrating complementation of the hallmark defect in chloride ion transport (Rich, D.P. et al. (1990) *Nature* 347:358-363). This initial work involved cells in tissue culture, however, subsequent work has shown that to deliver the gene to the airways of whole animals, defective adenoviruses may be useful (Rosenfeld, 15 (1992) *Cell* 68:143-155). However, the safety and effectiveness of using defective adenoviruses remain to be demonstrated.

Summary of the Invention

20 In general, the instant invention relates to vectors for transferring selected genetic material of interest (e.g., DNA or RNA) to cells *in vivo*. In preferred embodiments, the vectors are adenovirus-based. Advantages of adenovirus-based vectors for gene therapy are that they appear to be relatively safe and can be manipulated to encode the desired gene product and at the same time are inactivated in terms of their ability to replicate in a normal lytic viral life cycle. Additionally, adenovirus has a natural tropism for airway epithelia. 25 Therefore, adenovirus-based vectors are particularly preferred for respiratory gene therapy applications such as gene therapy for cystic fibrosis.

30 In one embodiment, the adenovirus-based gene therapy vector comprises an adenovirus 2 serotype genome in which the Ela and Elb regions of the genome, which are involved in early stages of viral replication have been deleted and replaced by genetic material of interest (e.g., DNA encoding the cystic fibrosis transmembrane regulator protein).

35 In another embodiment, the adenovirus-based therapy vector is a pseudo-adenovirus (PAV). PAVs contain no potentially harmful viral genes, have a theoretical capacity for foreign material of nearly 36 kb, may be produced in reasonably high titers and maintain the tropism of the parent adenovirus for dividing and non-dividing human target cell types. PAVs comprise adenovirus inverted terminal repeats and the minimal sequences of a wild-type adenovirus type 2 genome necessary for efficient replication and packaging by a helper virus and genetic material of interest. In a preferred embodiment, the PAV contains adenovirus 2 sequences.

In a further embodiment, the adenovirus-based gene therapy vector contains the open reading frame 6 (ORF6) of adenoviral early region 4 (E4) from the E4 promoter and is deleted for all other E4 open reading frames. Optionally, this vector can include deletions in the E1 and/or E3 regions. Alternatively, the adenovirus-based gene therapy vector contains the open reading frame 3 (ORF3) of adenoviral E4 from the E4 promoter and is deleted for all other E4 open reading frames. Again, optionally, this vector can include deletions in the E1 and/or E3 regions. The deletion of non-essential open reading frames of E4 increases the cloning capacity by approximately 2 kb without significantly reducing the viability of the virus in cell culture. In combination with deletions in the E1 and/or E3 regions of adenovirus vectors, the theoretical insert capacity of the resultant vectors is increased to 8-9 kb.

The invention also relates to methods of gene therapy using the disclosed vectors and genetically engineered cells produced by the method.

Brief Description of the Tables and Drawings

Further understanding of the invention may be had by reference to the tables and figures wherein:

Table I shows CFTR mutants wherein the known association with CF (Y, yes or N, no), exon localization, domain location and presence (+) or absence (-) of bands A, B, and C of mutant CFTR species is shown. TM6, indicates transmembrane domain 6; NBD nucleotide binding domain; ECD, extracellular domain and Term, termination at 21 codons past residue 1337;

Table II shows the nucleotide sequence of Ad2/CFTR-1;

Table III depicts a nucleotide analysis of Ad2-ORF6/PGK-CFTR;

The convention for naming mutants is first the amino acid normally found at the particular residue, the residue number (Riordan, T.R. et al. (1989) *Science* 245:1066-1073). and the amino acid to which the residue was converted. The single letter amino acid code is used: D, aspartic acid; F, phenylalanine; G, glycine; I, isoleucine; K, lysine; M, methionine; N, asparagine; Q, glutamine; R, arginine; S, serine; W, tryptophan. Thus G551D is a mutant in which glycine 551 is converted to aspartic acid;

Figure 1 shows alignment of CFTR partial cDNA clones used in construction of cDNA containing complete coding sequence of the CFTR, only restriction sites relevant to the DNA constructions described below are shown;

Figure 2 depicts plasmid construction of the CFTR cDNA clone pKK-CFTR1;

Figure 3 depicts plasmid construction of the CFTR cDNA clone pKK-CFTR2;

Figure 4 depicts plasmid construction of the CFTR cDNA clone pSC-CFTR2;

5

Figure 5 shows a plasmid map of the CFTR cDNA clone pSC-CFTR2;

Figure 6 shows the DNA sequence of synthetic DNAs used for insertion of an intron into the CFTR cDNA sequence, with the relevant restriction endonuclease sites and nucleotide positions noted;

10

Figures 7A and 7B depict plasmid construction of the CFTR cDNA clone pKK-CFTR3;

15

Figure 8 shows a plasmid map of the CFTR cDNA pKK-CFTR3 containing an intron between nucleotides 1716 and 1717;

Figure 9 shows treatment of CFTR with glycosidases;

20

Figures 10A and 10B show an analysis of CFTR expressed from COS-7 transfected cells;

Figures 11A and 11B show pulse-chase labeling of wild type and $\Delta F508$ mutant CFTR in COS-7 transfected cells;

25

Figures 12A-12D show immunolocalization of wild type and $\Delta F508$ mutant CFTR; and COS-7 cells transfected with pMT-CFTR or pMT-CFTR- $\Delta F508$;

Figure 13 shows an analysis of mutant forms of CFTR;

30

Figure 14 shows a map of the first generation adenovirus based vector encoding CFTR (Ad2/CFTR-1);

Figure 15 shows the plasmid construction of the Ad2/CFTR-1 vector;

35

Figure 16 shows an example of UV fluorescence from an agarose gel electrophoresis of products of nested RT-PCR from lung homogenates of cotton rats which received Ad2/CFTR-1. The gel demonstrates that the homogenates were positive for virally-encoded CFTR mRNA;

Figure 17 shows an example of UV fluorescence from an agarose gel electrophoresis of products of nested RT-PCR from organ homogenates of cotton rats. The gel demonstrates that all organs of the infected rats were negative for Ad2/CFTR with the exception of the small bowel;

Figures 18A and 18B show differential cell analyses of bronchoalveolar lavage specimens from control and infected rats. These data demonstrate that none of the rats treated with Ad2/CFTR-1 had a change in the total or differential white blood cell count 4, 10, and 14 days after infection (Figure 18A) and 3, 7, and 14 days after infection (Figure 18B);

Figure 19 shows hematoxylin and eosin stained sections of cotton rat tracheas from both treated and control rats sacrificed at different time points after infection with Ad2/CFTR-1. The sections demonstrate that there were no observable differences between the treated and control rats;

Figures 20A and 20B show examples of UV fluorescence from an agarose gel electrophoresis, stained with ethidium bromide, of products of RT-PCR from nasal brushings of Rhesus monkeys after application of Ad2/CFTR-1 or Ad2/ β -Gal;

Figure 21 shows lights microscopy and immunocytochemistry from monkey nasal brushings. The microscopy revealed that there was a positive reaction when nasal epithelial cells from monkeys exposed to Ad2/CFTR-1 were stained with antibodies to CFTR;

Figure 22 shows immunocytochemistry of monkey nasal turbinate biopsies. This microscopy reveals increased immunofluorescence at the apical membrane of the surface epithelium from biopsies obtained from monkeys treated with Ad2/CFTR-1 over that seen at the apical membrane of the surface epithelium from biopsies obtained from control monkeys;

Figures 23A-23D show serum antibody titers in Rhesus monkeys after three vector administrations. These graphs demonstrate that all three monkeys treated with Ad2/CFTR-1 developed antibodies against adenovirus;

Figure 24 shows hematoxylin and eosin stained sections from monkey medial turbinate biopsies. These sections demonstrate that turbinate biopsy specimens from control monkeys could not be differentiated from those from monkeys treated with Ad2/CFTR-1 when reviewed by an independent pathologist;

Figures 25A-25I show photomicrographs of human nasal mucosa immediately before, during, and after Ad2/CFTR-1 application. These photomicrographs demonstrate that inspection of the nasal mucosa showed mild to moderate erythema, edema, and exudate in patients treated with Ad2/CFTR-1 (Figures 25A-25C) and in control patients (Figures 25G-25I). These changes were probably due to local anesthesia and vasoconstriction because when an additional patient was exposed to Ad2/CFTR in a method which did not require the use of local anesthesia or vasoconstriction, there were no symptoms and the nasal mucosa appeared normal (Figures 25D-25F);

Figure 26 shows a photomicrograph of a hematoxylin and eosin stained biopsy of human nasal mucosa obtained from the third patient three days after Ad2/CFTR-1 administration. This section shows a morphology consistent with CF, i.e., a thickened basement membrane and occasional morphonuclear cells in the submucosa, but no abnormalities that could be attributed to the adenovirus vector;

Figure 27 shows transepithelial voltage (V_t) across the nasal epithelium of a normal human subject. Amiloride (μM) and terbutaline (μM) were perfused onto the mucosal surface beginning at the times indicated. Under basal conditions (V_t) was electrically negative. Perfusion of amiloride onto the mucosal surface inhibited (V_t) by blocking apical Na^+ channels;

Figures 28A and 28B show transepithelial voltage (V_t) across the nasal epithelium of normal human subjects (Figure 28A) and patients with CF (Figure 28B). Values were obtained under basal conditions, during perfusion with amiloride (μM), and during perfusion of amiloride plus terbutaline (μM) onto the mucosal surface. Data are from seven normal subjects and nine patients with CF. In patients with CF, (V_t) was more electrically negative than in normal subjects (Figure 28B). Amiloride inhibited (V_t) in CF patients, as it did in normal subjects. However, V_t failed to hyperpolarize when terbutaline was perfused onto the epithelium in the presence of amiloride. Instead, (V_t) either did not change or became less negative, a result very different from that observed in normal subjects;

Figures 29A and 29B show transepithelial voltage (V_t) across the nasal epithelium of a third patient before (Figure 29A) and after (Figure 29B) administration of approximately 25 MOI of Ad2/CFTR-1. Amiloride and terbutaline were perfused onto the mucosal surface beginning at the times indicated. Figure 29A shows an example from the third patient before treatment. Figure 29B shows that in contrast to the response before Ad2/CFTR-1 was applied, after virus replication, in the presence of amiloride, terbutaline stimulated V_t ;

Figures 30A-30F show the time of course changes in transepithelial electrical properties before and after administration of Ad2/CFTR-1. Figures 30A and 30B are from the first patient who received approximately 1 MOI; Figures 30C and 30D are from the second patient who received approximately 3 MOI; and Figures 30E and 30F are from the third patient who received approximately 25 MOI. Figures 30A, 30C, and 30E show values of basal transepithelial voltage (V_t) and Figures 30B, 30D, and 30F show the change in transepithelial voltage (ΔV_t) following perfusion of terbutaline in the presence of amiloride. Day zero indicates the day of Ad2/CFTR-1 administration. Figures 30A, 30C, and 30E show the time course of changes in basal V_t for all three patients. The decrease in basal V_t suggests that application of Ad2/CFTR-1 corrected the CF electrolyte transport defect in nasal epithelium of all three patients. Additional evidence came from an examination of the response to terbutaline. Figures 30B, 30D, and 30F show the time course of the response. These data indicate that Ad2/CFTR-1 corrected the CF defect in Cl^- transport;

Figure 31 shows the time course of changes in transepithelial electrical properties before and after administration of saline instead of Ad2/CFTR-1 to CF patients. Day zero indicates the time of mock administration. The top graph shows basal transepithelial voltage (V_t) and the bottom graph shows the change in transepithelial voltage following perfusion with terbutaline in the presence of amiloride (ΔV_t). Closed symbols are data from two patients that received local anesthetic/vasoconstriction and placement of the applicator for thirty minutes. Open symbol is data from a patient that received local anesthetic/vasoconstriction, but not placement of the applicator. Symptomatic changes and physical findings were the same as those observed in CF patients treated with a similar administration procedure and Ad2/CFTR-1;

Figure 32 shows a map of the second generation adenovirus based vector, PAV;

Figure 33 shows the plasmid construction of a second generation adenoviral vector 6 (Ad E4 ORF6);

Figure 34 is a schematic of Ad2-ORF6/PGK-CFTR which differs from Ad2/CFTR in that the latter utilized the endogenous Ela promoter, had no poly A addition signal directly downstream of CFTR and retained an intact E4 region;

Figure 35 shows short-circuit currents from human CF nasal polyp epithelial cells infected with Ad2-ORF6/PGK-CFTR at multiplicities of 0.3, 3, and 50. At the indicated times: (1) 10 μM amiloride, (2) cAMP agonists (10 μM forskolin and 100 μM IBMX, and (3) 1 mM diphenylamine-2-carboxylate were added to the mucosal solution;

Figures 36A-36D show immunocytochemistry of nasal brushings by laser scanning microscopy of the Rhesus monkey C, before infection (36A) and on 7 days (36B); 24 (36C); and 38 (36D) after the first infection with Ad2-ORF6/PGK-CFTR;

Figures 37A-37D show immunocytochemistry of nasal brushings by laser scanning microscopy of Rhesus monkey D, before infection (37A) and on days 7 (37B); 24 (37C); and 48 (37D) after the first infection with Ad2-ORF6/PGK-CFTR;

Figures 38A-38D show immunocytochemistry of nasal brushings by laser scanning microscopy of the Rhesus monkey E, before infection (38A) and on days 7 (38B); 24 (38C); and 48 (38D) after the first infection with Ad2-ORF6/PGK-CFTR;

Figures 39A-39C show summaries of the clinical signs (or lack thereof) of infection with Ad2-ORF6/PGK-CFTR;

Figures 40A-40C shows a summary of blood counts, sedimentation rate, and clinical chemistries after infection with Ad2-ORF6/PGK-CFTR for monkeys C, D, and E. There was no evidence of a systemic inflammatory response or other abnormalities of the clinical chemistries;

Figure 41 shows summaries of white blood cells counts in monkeys C, D, and E after infection with Ad2-ORF6/PGK-CFTR. These data indicate that the administration of Ad2-ORF6/PGK-CFTR caused no change in the distribution and number of inflammatory cells at any of the time points following viral administration;

Figure 42 shows histology of submucosal biopsy performed on Rhesus monkey C on day 4 after the second viral instillation of Ad2-ORF6/PGK-CFTR. Hematoxylin and eosin stain revealed no evidence of inflammation or cytopathic changes;

Figure 43 shows histology of submucosal biopsy performed on Rhesus monkey D on day 11 after the second viral instillation of Ad2-ORF6/PGK-CFTR. Hematoxylin and eosin stain revealed no evidence of inflammation or cytopathic changes;

Figure 44 shows histology of submucosal biopsy performed on Rhesus monkey E on day 18 after the second viral instillation of Ad2-ORF6/PGK-CFTR. Hematoxylin and eosin stain revealed no evidence of inflammation or cytopathic changes; and

- 10.1 -

Figures 45A-45C show antibody titers to adenovirus prior to and after the first and second administrations of Ad2-ORF6/PGK-CFTR. Prior to administration of Ad2-ORF6/PGK-

CFTR, the monkeys had received instillations of Ad2/CFTR-1. Antibody titers measured by ELISA rose within one week after the first and second administrations of Ad2-ORF6/PGK-CFTR. Serum neutralizing antibodies also rose within a week after viral administration and peaked at day 24. No anti-adenoviral antibodies were detected by ELISA or neutralizing assay in nasal washings of any of the monkeys.

Detailed Description and Best Mode

Gene Therapy

As used herein, the phrase "gene therapy" refers to the transfer of genetic material (e.g., DNA or RNA) of interest into a host to treat or prevent a genetic or acquired disease or condition. The genetic material of interest encodes a product (e.g., a protein polypeptide, peptide or functional RNA) whose production *in vivo* is desired. For example, the genetic material of interest can encode a hormone, receptor, enzyme or (poly) peptide of therapeutic value. Examples of genetic material of interest include DNA encoding: the cystic fibrosis transmembrane regulator (CFTR), Factor VIII, low density lipoprotein receptor, beta-galactosidase, alpha-galactosidase, beta-glucocerebrosidase, insulin, parathyroid hormone, and alpha-1-antitrypsin.

Although the potential for gene therapy to treat genetic diseases has been appreciated for many years, it is only recently that such approaches have become practical with the treatment of two patients with adenosine deaminase deficiency. The protocol consists of removing lymphocytes from the patients, stimulating them to grow in tissue culture, infecting them with an appropriately engineered retrovirus followed by reintroduction of the cells into the patient (Kantoff, P. et al. (1987) *J. Exp. Med.* 166:219). Initial results of treatment are very encouraging. With the approval of a number of other human gene therapy protocols for limited clinical use, and with the demonstration of the feasibility of complementing the CF defect by gene transfer, gene therapy for CF appears a very viable option.

The concept of gene replacement therapy for cystic fibrosis is very simple; a preparation of CFTR coding sequences in some suitable vector in a viral or other carrier delivered directly to the airways of CF patients. Since disease of the pulmonary airways is the major cause of morbidity and is responsible for 95% of mortality, airway epithelial cells are preferred target cells for CF gene therapy. The first generation of CF gene therapy is likely to be transient and to require repeated delivery to the airways. Eventually, however, gene therapy may offer a cure for CF when the identity of the precursor or stem cell to air epithelial cells becomes known. If DNA were incorporated into airway stem cells, all subsequent generations of such cells would make authentic CFTR from the integrated sequences and would correct the physiological defect almost irrespective of the biochemical basis of the action of CFTR.

Although simple in concept, scientific and clinical problems face approaches to gene therapy, not least of these being that CF requires an *in vivo* approach while all gene therapy treatments in humans to date have involved *ex vivo* treatment of cells taken from the patient followed by reintroduction.

- 5 One major obstacle to be overcome before gene therapy becomes a viable treatment approach for CF is the development of appropriate vectors to infect tissue manifesting the disease and deliver the therapeutic CFTR gene. Since viruses have evolved very efficient means to introduce their nucleic acid into cells, many approaches to gene therapy make use of engineered defective viruses. However, the use of viruses *in vivo* raises safety concerns.
- 10 Although potentially safer, the use of simple DNA plasmid constructs containing minimal additional DNA, on the other hand, is often very inefficient and can result in transient protein expression.

- The integration of introduced DNA into the host chromosome has advantages in that such DNA will be passed to daughter cells. In some circumstances, integrated DNA may
- 15 also lead to high or more sustained expression. However, integration often, perhaps always, requires cellular DNA replication in order to occur. This is certainly the case with the present generation of retroviruses. This limits the use of such viruses to circumstances where cell division occurs in a high proportion of cells. For cells cultured *in vitro*, this is seldom a problem, however, the cells of the airway are reported to divide only infrequently
- 20 (Kawanami, O. et al. (1979) *An. Rev. Respir. Dis.* 120:595). The use of retroviruses in CF will probably require damaging the airways (by agents such as SO₂ or O₃) to induce cell division. This may prove impracticable in CF patients.

- Even if efficient DNA integration could be achieved using viruses, the human genome contains elements involved in the regulation of cellular growth only a small fraction of which
- 25 are presently identified. By integrating adjacent to an element such as a proto-oncogene or an anti-oncogene, activation or inactivation of that element could occur leading to uncontrolled growth of the altered cell. It is considered likely that several such activation/inactivation steps are usually required in any one cell to induce uncontrolled proliferation (R.A. Weinberg (1989) *Cancer Research* 49:3713), which may reduce somewhat the potential risk. On the
- 30 other hand, insertional mutagenesis leading to tumor formation is certainly known in animals with some nondefective retroviruses (R.A. Weinberg, *supra*; Payne, G.S. et al. (1982) *Nature* 295:209), and the large numbers of potential integrations occurring during the lifetime of a patient treated repeatedly *in vivo* with retroviruses must raise concerns on the safety of such a procedure.

- 35 In addition to the potential problems associated with viral DNA integration, a number of additional safety issues arise. Many patients may have preexisting antibodies to some of the viruses that are candidates for vectors, for example, adenoviruses. In addition, repeated use of such vectors might induce an immune response. The use of defective viral vectors

may alleviate this problem somewhat, because the vectors will not lead to productive viral life cycles generating infected cells, cell lysis or large numbers of progeny viruses.

Other issues associated with the use of viruses are the possibility of recombination with related viruses naturally infecting the treated patient, complementation of the viral defects by simultaneous expression of wild type virus proteins and containment of aerosols of the engineered viruses.

Gene therapy approaches to CF will face many of the same clinical challenges at protein therapy. These include the inaccessibility of airway epithelium caused by mucus build-up and the hostile nature of the environment in CF airways which may inactivate viruses/vectors. Elements of the vector carriers may be immunogenic and introduction of the DNA may be inefficient. These problems, as with protein therapy, are exacerbated by the absence of a good animal model for the disease nor a simple clinical end point to measure the efficacy of treatment.

15 CF Gene Therapy Vectors - Possible Options

Retroviruses - Although defective retroviruses are the best characterized system and so far the only one approved for use in human gene therapy (Miller, A.D. (1990) *Blood* 76:271), the major issue in relation to CF is the requirement for dividing cells to achieve DNA integration and gene expression. Were conditions found to induce airway cell division, the *in vivo* application of retroviruses, especially if repeated over many years, would necessitate assessment of the safety aspects of insertional mutagenesis in this context.

Adeno-Associated Virus - (AAV) is a naturally occurring defective virus that requires other viruses such as adenoviruses or herpes viruses as helper viruses (Muzyczka, N. (1992) in *Current Topics in Microbiology and Immunology* 158:97). It is also one of the few viruses that may integrate its DNA into non-dividing cells, although this is not yet certain. Vectors containing as little as 300 base pairs of AAV can be packaged and can integrate, but space for exogenous DNA is limited to about 4.5 kb. CFTR DNA may be towards the upper limit of packaging. Furthermore, the packaging process itself is presently inefficient and safety issues such as immunogenicity, complementation and containment will also apply to AAV. Nevertheless, this system is sufficiently promising to warrant further study.

Plasmid DNA - Naked plasmid can be introduced into muscle cells by injection into the tissue. Expression can extend over many months but the number of positive cells is low (Wolff, J. et al. (1989) *Science* 247:1465). Cationic lipids aid introduction of DNA into some cells in culture (Felgner, P. and Ringold, G.M. (1989) *Nature* 337:387). Injection of cationic lipid plasmid DNA complexes into the circulation of mice has been shown to result in expression of the DNA in lung (Brigham, K. et al. (1989) *Am. J. Med. Sci.* 298:278).

Instillation of cationic lipid plasmid DNA into lung also leads to expression in epithelial cells but the efficiency of expression is relatively low and transient (Hazinski, T.A. et al. (1991) *Am. J. Respir., Cell Mol. Biol.* 4:206). One advantage of the use of plasmid DNA is that it can be introduced into non-replicating cells. However, the use of plasmid DNA in the CF airway environment, which already contains high concentrations of endogenous DNA may be problematic.

Receptor Mediated Entry - In an effort to improve the efficiency of plasmid DNA uptake, attempts have been made to utilize receptor-mediated endocytosis as an entry mechanisms and to protect DNA in complexes with polylysine (Wu, G. and Wu, C.H. (1988) *J. Biol. Chem.* 263:14621). One potential problem with this approach is that the incoming plasmid DNA enters the pathway leading from endosome to lysosome, where much incoming material is degraded. One solution to this problem is the use of transferrin DNA-polylysine complexes linked to adenovirus capsids (Curiel, D.T. et al. (1991) *Proc. Natl. Acad. Sci. USA* 88:8850). The latter enter efficiently but have the added advantage of naturally disrupting the endosome thereby avoiding shuttling to the lysosome. This approach has promise but at present is relatively transient and suffers from the same potential problems of immunogenicity as other adenovirus based methods.

Adenovirus - Defective adenoviruses at present appear to be a promising approach to CF gene therapy (Berkner, K.L. (1988) *BioTechniques* 6:616). Adenovirus can be manipulated such that it encodes and expresses the desired gene product, (e.g., CFTR), and at the same time is inactivated in terms of its ability to replicate in a normal lytic viral life cycle. In addition, adenovirus has a natural tropism for airway epithelia. The viruses are able to infect quiescent cells as are found in the airways, offering a major advantage over retroviruses. Adenovirus expression is achieved without integration of the viral DNA into the host cell chromosome, thereby alleviating concerns about insertional mutagenesis. Furthermore, adenoviruses have been used as live enteric vaccines for many years with an excellent safety profile (Schwartz, A.R. et al. (1974) *Am. Rev. Respir. Dis.* 109:233-238). Finally, adenovirus mediated gene transfer has been demonstrated in a number of instances including transfer of alpha-1-antitrypsin and CFTR to the lungs of cotton rats (Rosenfeld, M.A. et al. (1991) *Science* 252:431-434; Rosenfeld et al., (1992) *Cell* 68:143-155). Furthermore, extensive studies to attempt to establish adenovirus as a causative agent in human cancer were uniformly negative (Green, M. et al. (1979) *Proc. Natl. Acad. Sci. USA* 76:6606).

The following properties would be desirable in the design of an adenovirus vector to transfer the gene for CFTR to the airway cells of a CF patient. The vector should allow sufficient expression of the CFTR, while producing minimal viral gene expression. There should be minimal viral DNA replication and ideally no virus replication. Finally,

recombination to produce new viral sequences and complementation to allow growth of the defective virus in the patient should be minimized. A first generation adenovirus vector encoding CFTR (Ad2/CFTR), made as described in the following Example 7, achieves most of these goals and was used in the human trials described in Example 10.

5 Figure 14 shows a map of Ad2/CFTR-1. As can be seen from the figure, this first generation virus includes viral DNA derived from the common relatively benign adenovirus 2 serotype. The Ela and Elb regions of the viral genome, which are involved in early stages of viral replication have been deleted. Their removal impairs viral gene expression and viral replication. The protein products of these genes also have immortalizing and transforming
10 function in some non-permissive cells.

 The CFTR coding sequence is inserted into the viral genome in place of the Ela/Elb region and transcription of the CFTR sequence is driven by the endogenous Ela promoter. This is a moderately strong promoter that is functional in a variety of cells. In contrast to some adenovirus vectors (Rosenfeld, M. et al. (1992) *Cell* 68:143), this adenovirus retains
15 the E3 viral coding region. As a consequence of the inclusion of E3, the length of the adenovirus-CFTR DNA is greater than that of the wild-type adenovirus. The greater length of the recombinant viral DNA renders it more difficult to package. This means that the growth of the Ad2/CFTR virus is impaired even in permissive cells that provide the missing Ela and Elb functions.

20 The E3 region of the Ad2/CFTR-1 encodes a variety of proteins. One of these proteins, gp19, is believed to interact with and prevent presentation of class I proteins of the major histocompatibility complex (MHC) (Gooding, C.R. and Wold, W.S.M. (1990) *Crit. Rev. Immunol.* 10:53). This property prevents recognition of the infected cells and thus may allow viral latency. The presence of E3 sequences, therefore, has two useful attributes; first,
25 the large size of the viral DNA renders it doubly defective for replication (i.e., it lacks early functions and is packaged poorly) and second, the absence of MHC presentation could be useful in later applications of Ad2/CFTR-1 in gene therapy involving multiple administrations because it may avoid an immune response to recombinant virus containing cells.

30 Not only are there advantages associated with the presence of E3; there may be disadvantages associated with its absence. Studies of E3 deleted virus in animals have suggested that they result in a more severe pathology (Gingsberg, H.S. et al. (1989) *Proc. Natl. Acad. Sci. (USA)* 86:3823). Furthermore, E3 deleted virus, such as might be obtained by recombination of an E1 plus E3 deleted virus with wild-type virus, is reported to outgrow
35 wild-type in tissue culture (Barkner, K.L. and Sharp, P. (1983) *Nucleic Acids Research* 11:6003). By contrast, however, a recent report of an E3 replacement vector encoding hepatitis B surface antigen, suggests that when delivered as a live enteric vaccine, such a virus replicates poorly in human compared to wild-type.

The adenovirus vector (Ad2/CFTR-1) and a related virus encoding the marker β -galactosidase (Ad2/ β -gal) have been constructed and grown in human 293 cells. These cells contain the E1 region of adenovirus and constitutively express Ela and Elb, which complement the defective adenoviruses by providing the products of the genes deleted from the vector. Because the size of its genome is greater than that of wild-type virus, Ad2/CFTR is relatively difficult to produce.

The Ad2/CFTR-1 virus has been shown to encode CFTR by demonstrating the presence of the protein in 293 cells. The Ad2/ β -gal virus was shown to produce its protein in a variety of cell lines grown in tissue culture including a monkey bronchiolar cell line (4MBR-5), primary hamster tracheal epithelial cells, human HeLa, human CF PAC cells (see Example 8) and airway epithelial cells from CF patients (Rich, O. et al. (1990) *Nature* 347:358).

Ad2/CFTR-1 is constructed from adenovirus 2 (Ad2) DNA sequences. Other varieties of adenovirus (e.g., Ad3, Ad5, and Ad7) may also prove useful as gene therapy vectors. This may prove essential if immune response against a single serotype reduces the effectiveness of the therapy.

Second Generation Adenoviral Vectors

Adenoviral vectors currently in use retain most ($\geq 80\%$) of the parental viral genetic material leaving their safety untested and in doubt. Second-generation vector systems containing minimal adenoviral regulatory, packaging and replication sequences have therefore been developed.

Pseudo-Adenovirus Vectors (PAV)-PAVs contain adenovirus inverted terminal repeats and the minimal adenovirus 5' sequences required for helper virus dependent replication and packaging of the vector. These vectors contain no potentially harmful viral genes, have a theoretical capacity for foreign material of nearly 36 kb, may be produced in reasonably high titers and maintain the tropism of the parent virus for dividing and non-dividing human target cell types.

The PAV vector can be maintained as either a plasmid-borne construct or as an infectious viral particle. As a plasmid construct, PAV is composed of the minimal sequences from wild type adenovirus type 2 necessary for efficient replication and packaging of these sequences and any desired additional exogenous genetic material, by either a wild-type or defective helper virus.

Specifically, PAV contains adenovirus 2 (Ad2) sequences as shown in Figure 17, from nucleotide (nt) 0-356 forming the 5' end of the vector and the last 109 nt of Ad2 forming the 3' end of the construct. The sequences includes the Ad2 flanking inverted terminal repeats (5'ITR) and the 5' ITR adjoining sequences containing the known packaging signal and Ela enhancer. Various convenient restriction sites have been incorporated into the

fragments, allowing the insertion of promoter/gene cassettes which can be packaged in the PAV virion and used for gene transfer (e.g. for gene therapy). The construction and propagation of PAV is described in detail in the following Example 11. By not containing most native adenoviral DNA, the PAVs described herein are less likely to produce a patient immune response or to replicate in a host.

In addition, the PAV vectors can accommodate foreign DNA up to a maximum length of nearly 36 kb. The PAV vectors therefore, are especially useful for cloning larger genes (e.g., CFTR (7.5 kb)); Factor VIII (8 kb); Factor IX (9 kb)), which, traditional vectors have difficulty accommodating. In addition, PAV vectors can be used to transfer more than one gene, or more than one copy of a particular gene. For example, for gene therapy of cystic fibrosis, PAVs can be used to deliver CFTR in conjunction with other genes such as anti proteases (e.g., antiprotease alpha-1-antitrypsin) tissue inhibitor of metalloproteinase, antioxidants (e.g., superoxide dismutase), enhancers of local host defense (e.g., interferons), mucolytics (e.g., DNase); and proteins which block inflammatory cytokines.

Ad2-E4/ORF6 Adenovirus Vectors

An adenoviral construct expressing only the open reading frame 6 (ORF6) of adenoviral early region 4 (E4) from the E4 promoter and which is deleted for all other known E4 open reading frames was constructed as described in detail in Example 12. Expression of E4 open reading frame 3 is also sufficient to provide E4 functions required for DNA replication and late protein synthesis. However, it provides these functions with reduced efficiency compared to expression of ORF6, which will likely result in lower levels of virus production. Therefore expressing ORF6, rather than ORF3, appears to be a better choice for producing recombinant adenovirus vectors.

The E4 region of adenovirus is suspected to have a role in viral DNA replication, late mRNA synthesis and host protein synthesis shut off, as well as in viral assembly (Falgout, B. and G. Ketner (1987) *J. Virol.* 61:3759-3768). Adenovirus early region 4 is required for efficient virus particle assembly. Adenovirus early region 4 encodes functions required for efficient DNA replication, late gene expression, and host cell shutoff. Halbert, D.N. et al. (1985) *J. Virol.* 56:250-257.

The deletion of non-essential open reading frames of E4 increases the cloning capacity of recombinant adenovirus vectors by approximately 2 kb of insert DNA without significantly reducing the viability of the virus in cell culture. When placed in combination with deletions in the E1 and/or E3 regions of adenovirus vectors, the theoretical insert capacity of the resultant vectors is increased to 8-9 kb. An example of where this increased cloning capacity may prove useful is in the development of a gene therapy vector encoding CFTR. As described above, the first generation adenoviral vector approaches the maximum packaging capacity for viral DNA encapsidation. As a result, this virus grows poorly and may occasionally give rise to defective progeny. Including an E4 deletion in the adenovirus

vector should alleviate these problems. In addition, it allows flexibility in the choice of promoters to drive CFTR expression from the virus. For example, strong promoters such as the adenovirus major late promoter, the cytomegalovirus immediate early promoter or a cellular promoter such as the CFTR promoter, which may be too large for first-generation
5 adenovirus can be used to drive expression.

In addition, by expressing only ORF6 of E4, these second generation adenoviral vectors may be safer for use in gene therapy. Although ORF6 expression is sufficient for viral DNA replication and late protein synthesis in immortalized cells, it has been suggested that ORF6/7 of E4 may also be required in non-dividing primary cells (Hemstrom, C. et al.
10 (1991) *J. Virol.* 65:1440-1449). The 19 kD protein produced from open reading frame 6 and 7 (ORF6/7) complexes with and activates cellular transcription factor E2F, which is required for maximal activation of early region 2. Early region 2 encodes proteins required for viral DNA replication. Activated transcription factor E2F is present in proliferating cells and is involved in the expression of genes required for cell proliferation (e.g., DHFR, c-myc),
15 whereas activated E2F is present in lower levels in non-proliferating cells. Therefore, the expression of only ORF6 of E4 should allow the virus to replicate normally in tissue culture cells (e.g., 293 cells), but the absence of ORF6/7 would prevent the potential activation of transcription factor E2F in non-dividing primary cells and thereby reduce the potential for viral DNA replication.

20

Target Tissue

Because 95% of CF patients die of lung disease, the lung is a preferred target for gene therapy. The hallmark abnormality of the disease is defective electrolyte transport by the epithelial cells that line the airways. Numerous investigators (reviewed in Quinton, F. (1990)
25 *FASEB J.* 4:2709) have observed: a) a complete loss of cAMP-mediated transepithelial chloride secretion, and b) a two to three fold increase in the rate of Na⁺ absorption. cAMP-stimulated chloride secretion requires a chloride channel in the apical membrane (Welsh, M.J. (1987) *Physiol Rev.* 67:1143-1184). The discovery that CFTR is a phosphorylation-regulated chloride channel and that the properties of the CFTR chloride channel are the same as those
30 of the chloride channels in the apical membrane, indicate that CFTR itself mediates transepithelial chloride secretion. This conclusion was supported by studies localizing CFTR in lung tissue: CFTR is located in the apical membrane of airway epithelial cells (Denning, G.M. et al. (1992) *J. Cell Biol.* 118:551) and has been reported to be present in the submucosal glands (Taussig et al., (1973) *J. Clin. Invest.* 89:339). As a consequence of loss
35 of CFTR function, there is a loss of cAMP-regulated transepithelial chloride secretion. At this time it is uncertain how dysfunction of CFTR produces an increase in the rate of Na⁺ absorption. However, it is thought that the defective chloride secretion and increased Na⁺ absorption lead to an alteration of the respiratory tract fluid and hence, to defective mucociliary clearance, a normal pulmonary defense mechanism. As a result, clearance of

inhaled material from the lung is impaired and repeated infections ensue. Although the presumed abnormalities in respiratory tract fluid and mucociliary clearance provide a plausible explanation for the disease, a precise understanding of the pathogenesis is still lacking.

5 Correction of the genetic defect in the airway epithelial cells is likely to reverse the CF pulmonary phenotype. The identity of the specific cells in the airway epithelium that express CFTR cannot be accurately determined by immunocytochemical means, because of the low abundance of protein. However, functional studies suggest that the ciliated epithelial cells and perhaps nonciliated cells of the surface epithelium are among the main cell types
10 involved in electrolyte transport. Thus, in practical terms, the present preferred target cell for gene therapy would appear to be the mature cells that line the pulmonary airways. These are not rapidly dividing cells; rather, most of them are nonproliferating and many may be terminally differentiated. The identification of the progenitor cells in the airway is uncertain. Although CFTR may also be present in submucosal glands (Trezise, A.E. and Buchwald, M.
15 (1991) *Nature* 353:434; Englehardt, J.F. et al. (1992) *J. Clin. Invest.* 90:2598-2607), there is no data as to its function at that site; furthermore, such glands appear to be relatively inaccessible.

 The airway epithelium provides two main advantages for gene therapy. First, access to the airway epithelium can be relatively noninvasive. This is a significant advantage in the
20 development of delivery strategies and it will allow investigators to monitor the therapeutic response. Second, the epithelium forms a barrier between the airway lumen and the interstitium. Thus, application of the vector to the lumen will allow access to the target cell yet, at least to some extent, limit movement through the epithelial barrier to the interstitium and from there to the rest of the body.

25

Efficiency of Gene Delivery Required to Correct The Genetic Defect

 It is unlikely that any gene therapy protocol will correct 100% of the cells that normally express CFTR. However, several observations suggest that correction of a small
30 percent of the involved cells or expression of a fraction of the normal amount of CFTR may be of therapeutic benefit.

a. CF is an autosomal recessive disease and heterozygotes have no lung disease. Thus, 50% of wild-type CFTR would appear sufficient for normal function.

35 b. This issue was tested in mixing experiments using CF cells and recombinant CF cells expressing wild-type CFTR (Johnson, L.G. et al. (1992) *Nature Gen.* 2:21). The data obtained showed that when an epithelium is reconstituted with as few as 6-10% of corrected cells, chloride secretion is comparable to that observed with an epithelium containing 100% corrected cells. Although CFTR expression in the recombinant cells is

probably higher than in normal cells, this result suggests that *in vivo* correction of all CF airway cells may not be required.

5 c. Recent observations show that CFTR containing some CF-associated mutations retains residual chloride channel activity (Sheppard, D.N. et al. (1992) *Pediatr. Pulmon Suppl.* 8:250; Strong, T.V. et al. (1991) *N. Eng. J. Med.* 325:1630). These mutations are associated with mild lung disease. Thus, even a very low level of CFTR activity may at least partly ameliorate the electrolyte transport abnormalities.

10 d. As indicated in experiments described below in Example 8, complementation of CF epithelia, under conditions that probably would not cause expression of CFTR in every cell, restored cAMP stimulated chloride secretion.

15 e. Levels of CFTR in normal human airway epithelia are very low and are barely detectable. It has not been detected using routine biochemical techniques such as immunoprecipitation or immunoblotting and has been exceedingly difficult to detect with immunocytochemical techniques (Denning, G.M. et al. (1992) *J. Cell Biol.* 118:551). Although CFTR has been detected in some cases using laser-scanning confocal microscopy, the signal is at the limits of detection and cannot be detected above background in every case.
20 Despite that minimal levels of CFTR, this small amount is sufficient to generate substantial cAMP-stimulated chloride secretion. The reason that a very small number of CFTR chloride channels can support a large chloride secretory rate is that a large number of ions can pass through a single channel (10^6 - 10^7 ions/sec) (Hille, B. (1984) Sinauer Assoc. Inc., Sunderland, MA 420-426).

25

 f. Previous studies using quantitative PCR have reported that the airway epithelial cells contain at most one to two transcripts per cell (Trapnell, B.C. et al. (1991) *Proc. Natl. Acad. Sci. USA* 88:6565).

30

 Gene therapy for CF would appear to have a wide therapeutic index. Just as partial expression may be of therapeutic value, overexpression of wild-type CFTR appears unlikely to cause significant problems. This conclusion is based on both theoretical considerations and experimental results. Because CFTR is a regulated channel, and because it has a specific function in epithelia, it is unlikely that overexpression of CFTR will lead to uncontrolled
35 chloride secretion. First, secretion would require activation of CFTR by cAMP-dependent phosphorylation. Activation of this kinase is a highly regulated process. Second, even if CFTR chloride channels open in the apical membrane, secretion will not ensue without regulation of the basolateral membrane transporters that are required for chloride to enter the cell from the interstitial space. At the basolateral membrane, the sodium-potassium-chloride

cotransporter and potassium channels serve as important regulators of transepithelial secretion (Welsh, M.J. (1987) *Physiol. Rev.* 67:1143-1184).

Human CFTR has been expressed in transgenic mice under the control of the surfactant protein C (SPC) gene promoter (Whitesett, J.A. et al. (1992) *Nature Gen.* 2:13) and the casein promoter (Ditullio, P. et al (1992) *Bio/Technology* 10:74). In those mice, CFTR was overexpressed in bronchiolar and alveolar epithelial cells and in the mammary glands, respectively. Yet despite the massive overexpression in the transgenic animals, there were no observable morphologic or functional abnormalities. In addition, expression of CFTR in the lungs of cotton rats produced no reported abnormalities (Rosenfeld, M.A. et al. (1992) *Cell* 68:143-155).

The present invention is further illustrated by the following examples which in no way should be construed as being further limiting. The contents of all cited references (including literature references, issued patents, published patent applications, and co-pending patent applications) cited throughout this application are hereby expressly incorporated by reference.

EXAMPLES

Example 1 - Generation of Full Length CFTR cDNAs

Nearly all of the commonly used DNA cloning vectors are based on plasmids containing modified pMB1 replication origins and are present at up to 500 to 700 copies per cell (Sambrook et al. *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor Laboratory Press 1989). The partial CFTR cDNA clones isolated by Riordan et al. were maintained in such a plasmid. It was postulated that an alternative theory to intrinsic clone instability to explain the apparent inability to recover clones encoding full length CFTR protein using high copy number plasmids, was that it was not possible to clone large segments of the CFTR cDNA at high gene dosage in *E. coli*. Expression of the CFTR or portions of the CFTR from regulatory sequences capable of directing transcription and/or translation in the bacterial host cell might result in inviability of the host cell due to toxicity of the transcript or of the full length CFTR protein or fragments thereof. This inadvertent gene expression could occur from either plasmid regulatory sequences or cryptic regulatory sequences within the recombinant CFTR plasmid which are capable of functioning in *E. coli*. Toxic expression of the CFTR coding sequences would be greatly compounded if a large number of copies of the CFTR cDNA were present in cells because a high copy number plasmid was used. If the product was indeed toxic as postulated, the growth of cells containing full length and correct sequence would be actively disfavored. Based upon this novel hypothesis, the following procedures were undertaken. With reference to Figure 2, partial CFTR clone T16-4.5 was cleaved with restriction enzymes Sph 1 and Pst 1 and the resulting 3.9 kb restriction fragment containing exons 11 through most of exon 24 (including

an uncharacterized 119 bp insertion reported by Riordan et al. between nucleotides 1716 and 1717), was isolated by agarose gel purification and ligated between the Sph I and Pst I sites of the pMB1 based vector pkk223-3 (Brosius and Holy, (1984) *Proc. Natl. Acad. Sci.* 81:6929). It was hoped that the pMB1 origin contained within this plasmid would allow it and plasmids constructed from it to replicate at 15-20 copies per host *E. coli* cell (Sambrook et al. *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor Laboratory Press 1989). The resultant plasmid clone was called pkk-4.5.

Partial CFTR clone T11 was cleaved with Eco R1 and Hinc II and the 1.9 kb band encoding the first 1786 nucleotides of the CFTR cDNA plus an additional 100 bp of DNA at the 5' end was isolated by agarose gel purification. This restriction fragment was inserted between the Eco R1 site and Sma I restriction site of the plasmid Bluescript Sk- (Stratagene, catalogue number 212206), such that the CFTR sequences were now flanked on the upstream (5') side by a Sal I site from the cloning vector. This clone, designated T11-R, was cleaved with Sal I and Sph I and the resultant 1.8 kb band isolated by agarose gel purification. Plasmid pkk-4.5 was cleaved with Sal I and Sph I and the large fragment was isolated by agarose gel purification. The purified T11-R fragment and pkk-4.5 fragments were ligated to construct pkk-CFTR1. pkk-CFTR1 contains exons 1 through 24 of the CFTR cDNA. It was discovered that this plasmid is stably maintained in *E. coli* cells and confers no measureably disadvantageous growth characteristics upon host cells.

pkk-CFTR1 contains, between nucleotides 1716 and 1717, the 119 bp insert DNA derived from partial cDNA clone T16-4.5 described above. In addition, subsequent sequence analysis of pkk-CFTR1 revealed unreported differences in the coding sequence between that portion of CFTR1 derived from partial cDNA clone T11 and the published CFTR cDNA sequence. These undesired differences included a 1 base-pair deletion at position 995 and a C to T transition at position 1507.

To complete construction of an intact correct CFTR coding sequence without mutations or insertions and with reference to the construction scheme shown in Figure 3, pkk-CFTR1 was cleaved with Xba I and Hpa I, and dephosphorylated with calf intestinal alkaline phosphatase. In addition, to reduce the likelihood of recovering the original clone, the small unwanted Xba I/Hpa I restriction fragment from pKK-CFTR1 was digested with Sph I. T16-1 was cleaved with Xba I and Acc I and the 1.15 kb fragment isolated by agarose gel purification. T16-4.5 was cleaved with Acc I and Hpa I and the 0.65 kb band was also isolated by agarose gel purification. The two agarose gel purified restriction fragments and the dephosphorylated pKK-CFTR1 were ligated to produce pKK-CFTR2. Alternatively, pKK-CFTR2 could have been constructed using corresponding restriction fragments from the partial CFTR cDNA clone C1-1/5. pKK-CFTR2 contains the uninterrupted CFTR protein coding sequence and conferred slow growth upon *E. coli* host cells in which it was inserted, whereas pKK-CFTR1 did not. The origin of replication of pKK-CFTR2 is derived from pMB1 and confers a plasmid copy number of 15-20 copies per host cell.

Example 2 - Improving Host Cell Viability

An additional enhancement of host cell viability was accomplished by a further reduction in the copy number of CFTR cDNA per host cell. This was achieved by transferring the CFTR cDNA into the plasmid vector, pSC-3Z. pSC-3Z was constructed using the pSC101 replication origin of the low copy number plasmid pLG338 (Stoker *et al.*, Gene 18, 335 (1982)) and the ampicillin resistance gene and polylinker of pGEM-3Z (available from Promega). pLG338 was cleaved with Sph I and Pvu II and the 2.8 kb fragment containing the replication origin isolated by agarose gel purification. pGEM-3Z was cleaved with Alw NI, the resultant restriction fragment ends treated with T4 DNA polymerase and deoxynucleotide triphosphates, cleaved with Sph I and the 1.9 kb band containing the ampicillin resistance gene and the polylinker was isolated by agarose gel purification. The pLG338 and pGEM-3Z fragments were ligated together to produce the low copy number cloning vector pSC-3Z. pSC-3Z and other plasmids containing pSC101 origins of replication are maintained at approximately five copies per cell (Sambrook *et al. supra*).

With additional reference to Figure 4, pKK-CFTR2 was cleaved with Eco RV, Pst I and Sal I and then passed over a Sephacryl S400 spun column (available from Pharmacia) according to the manufacturer's procedure in order to remove the Sal I to Eco RV restriction fragment which was retained within the column. pSC-3Z was digested with Sma I and Pst I and also passed over a Sephacryl S400 spun column to remove the small Sma I/Pst I restriction fragment which was retained within the column. The column eluted fractions from the pKK-CFTR2 digest and the pSC-3Z digest were mixed and ligated to produce pSC-CFTR2. A map of this plasmid is presented in Figure 5. Host cells containing CFTR cDNAs at this and similar gene dosages grow well and have stably maintained the recombinant plasmid with the full length CFTR coding sequence. In addition, this plasmid contains a bacteriophage T7 RNA polymerase promoter adjacent to the CFTR coding sequence and is therefore convenient for *in vitro* transcription/translation of the CFTR protein. The nucleotide sequence of CFTR coding region from pSC-CFTR2 plasmid is presented in Sequence Listing 1 as SEQ ID NO:1. Significantly, this sequence differs from the previously published (Riordan, J.R. et al. (1989) *Science* 245:1066-1073) CFTR sequence at position 1990, where there is C in place of the reported A. See Gregory, R.J. et al. (1990) *Nature* 347:382-386. *E. coli* host cells containing pSC-CFTR2, internally identified with the number pSC-CFTR2/AG1, have been deposited at the American Type Culture Collection and given the accession number: ATCC 68244.

Example 3 - Alternate Method for Improving Host Cell Viability

A second method for enhancing host cell viability comprises disruption of the CFTR protein coding sequence. For this purpose, a synthetic intron was designed for insertion between nucleotides 1716 and 1717 of the CFTR cDNA. This intron is especially

advantageous because of its easily manageable size. Furthermore, it is designed to be efficiently spliced from CFTR primary RNA transcripts when expressed in eukaryotic cells. Four synthetic oligonucleotides were synthesized (1195RG, 1196RG, 1197RG and 1198RG) collectively extending from the Sph I cleavage site at position 1700 to the Hinc II cleavage site at position 1785 and including the additional 83 nucleotides between 1716 and 1717 (see Figure 6). These oligonucleotides were phosphorylated with T4 polynucleotide kinase as described by Sambrook et al., mixed together, heated to 95°C for 5 minutes in the same buffer used during phosphorylation, and allowed to cool to room temperature over several hours to allow annealing of the single stranded oligonucleotides. To insert the synthetic intron into the CFTR coding sequence and with reference to Figures 7A and 7B, a subclone of plasmid T11 was made by cleaving the Sal I site in the polylinker, repairing the recessed ends of the cleaved DNA with deoxynucleotide triphosphates and the large fragment of DNA Polymerase I and religating the DNA. This plasmid was then digested with Eco RV and Nru I and religated. The resulting plasmid T16-Δ5' extended from the Nru I site at position 490 of the CFTR cDNA to the 3' end of clone T16 and contained single sites for Sph I and Hinc II at positions corresponding to nucleotides 1700 and 1785 of the CFTR cDNA. T16-Δ5' plasmid was cleaved with Sph I and Hinc II and the large fragment was isolated by agarose gel purification. The annealed synthetic oligonucleotides were ligated into this vector fragment to generate T16-intron.

T16-intron was then digested with Eco RI and Sma I and the large fragment was isolated by agarose gel purification. T16-4.5 was digested with Eco RI and Sca I and the 790 bp fragment was also isolated by agarose gel purification. The purified T16-intron and T16-4.5 fragments were ligated to produce T16-intron-2. T16-intron-2 contains CFTR cDNA sequences extending from the Nru I site at position 490 to the Sca I site at position 2818, and includes the unique Hpa I site at position 2463 which is not present in T16-1 or T16-intron-1.

T16-intron-2 was then cleaved with Xba I and Hpa I and the 1800 bp fragment was isolated by agarose gel purification. pKK-CFTR1 was digested with Xba I and Hpa I and the large fragment was also isolated by agarose gel purification and ligated with the fragment derived from T16-intron-2 to yield pKK-CFTR3, shown in Figure 8. The CFTR cDNA within pKK-CFTR3 is identical to that within pSC-CFTR2 and pKK-CFTR2 except for the insertion of the 83 bp intron between nucleotides 1716 and 1717. The insertion of this intron resulted in improved growth characteristics for cells harboring pKK-CFTR3 relative to cells containing the unmodified CFTR cDNA in pKK-CFTR2.

Example 4 - In vitro Transcription/Translation

In addition to sequence analysis, the integrity of the CFTR cDNA open reading frame was verified by *in vitro* transcription/translation. This method also provided the initial CFTR protein for identification purposes. 5 micrograms of pSC-CFTR2 plasmid DNA were linearized with Sal I and used to direct the synthesis of CFTR RNA transcripts with T7 RNA

polymerase as described by the supplier (Stratagene). This transcript was extracted with phenol and chloroform and precipitated with ethanol. The transcript was resuspended in 25 microliters of water and varying amounts were added to a reticulocyte lysate *in vitro* translation system (Promega). The reactions were performed as described by the supplier in the presence of canine pancreatic microsomal membranes (Promega), using ^{35}S -methionine to label newly synthesized proteins. *In vitro* translation products were analysed by discontinuous polyacrylamide gel electrophoresis in the presence of 0.1% SDS with 8% separating gels (Laemmli, U.K. (1970) *Nature* 227:680-685). Before electrophoresis, the *in vitro* translation reactions were denatured with 3% SDS, 8 M urea and 5% 2-mercaptoethanol in 0.65 M Tris-HCl, pH 6.8. Following electrophoresis, the gels were fixed in methanol:acetic acid:water (30:10:60), rinsed with water and impregnated with 1 M sodium salicylate. ^{35}S labelled proteins were detected by fluorography. A band of approximately 180 kD was detected, consistent with translation of the full length CFTR insert.

15

Example 5 - Elimination of Cryptic Regulatory Signals

Analysis of the DNA sequence of the CFTR has revealed the presence of a potential *E. coli* RNA polymerase promoter between nucleotides 748 and 778 which conforms well to the derived consensus sequence for *E. coli* promoters (Reznikoff and McClure, Maximizing Gene Expression, 1, Butterworth Publishers, Stoneham, MA). If this sequence functions as a promoter functions in *E. coli*, it could direct synthesis of potentially toxic partial CFTR polypeptides. Thus, an additional advantageous procedure for maintaining plasmids containing CFTR cDNAs in *E. coli* would be to alter the sequence of this potential promoter such that it will not function in *E. coli*. This may be accomplished without altering the amino acid sequence encoded by the CFTR cDNA. Specifically, plasmids containing complete or partial CFTR cDNA's would be altered by site-directed mutagenesis using synthetic oligonucleotides (Zoller and Smith, (1983) *Methods Enzymol.* 100:468). More specifically, altering the nucleotide sequence at position 908 from a T to C and at position 774 from an A to a G effectively eliminates the activity of this promoter sequence without altering the amino acid coding potential of the CFTR open reading frame. Other potential regulatory signals within the CFTR cDNA for transcription and translation could also be advantageously altered and/or deleted by the same method.

Futher analysis has identified a sequence extending from nucleotide 908 to 936 which functions efficiently as a transcriptional promoter element in *E. coli* (Gregory, R.J. et al. (1990) *Nature* 347:382-386). Mutation at position 936 is capable of inactivating this promoter and allowing the CFTR cDNA to be stably maintained as a plasmid in *E. coli* (Cheng, S.H. et al. (1990) *Cell* 63:827-834). Specifically position 936 has been altered from a C to a T residue without the amino acid sequence encoded by the cDNA being altered. Other mutations within this regulatory element described in Gregory, R.J. et al. (1990)

Nature 347:382-386 could also be used to inactivate the transcriptional promoter activity. Specifically, the sequence from 908 to 913 (TTGTGA) and from 931 to 936 (GAAAAT) could be altered by site directed mutagenesis without altering the amino acid sequence encoded by the cDNA.

5

Example 6 - Cloning of CFTR in Alternate Host Systems

Although the CFTR cDNA displays apparent toxicity in *E. coli* cells, other types of host cells may not be affected in this way. Alternative host systems in which the entire CFTR cDNA protein encoding region may be maintained and/or expressed include other bacterial species and yeast. It is not possible *a priori* to predict which cells might be resistant and which might not. Screening a number of different host/vector combinations is necessary to find a suitable host tolerant of expression of the full length protein or potentially toxic fragments thereof.

15

Example 7 - Generation of Adenovirus Vector Encoding CFTR (Ad2/CFTR)

1. DNA preparation - Construction of the recombinant Ad2/CFTR-1 virus (the sequence of which is shown in Table II and as SEQ ID NO:3) was accomplished as follows: The CFTR cDNA was excised from the plasmid pCMV-CFTR-936C using restriction enzymes SpeI and EclI361. pCMV-CFTR-936C consists of a minimal CFTR cDNA encompassing nucleotides 123-4622 of the published CFTR sequence cloned into the multiple cloning site of pRC/CMV (Invitrogen Corp.) using synthetic linkers. The CFTR cDNA within this plasmid has been completely sequenced. The SpeI/EclI361 restriction fragment contains 47 bp of 5' sequence derived from synthetic linkers and the multiple cloning site of the vector.

25 The CFTR cDNA (the sequence of which is shown as SEQ ID NO:1 and the amino acid sequence encoded by the CFTR cDNA is shown as SEQ ID NO:2) was inserted between the NheI and SnaBI restriction sites of the adenovirus gene transfer vector pBR-Ad2-7. pBR-Ad2-7 is a pBR322 based plasmid containing an approximately 7 kb insert derived from the 5' 10680 bp of Ad2 inserted between the Clal and BamHI sites of pBR322. From this Ad2 fragment, the sequences corresponding to Ad2 nucleotides 546-3497 were deleted and replaced with a 12 bp multiple cloning site containing an NheI site, an MluI site, and a SnaBI site. The construct also contains the 5' inverted terminal repeat and viral packaging signals, the Ela enhancer and promoter, the Elb 3' intron and the 3' untranslated region and polyadenylation sites. The resulting plasmid was called pBR-Ad2-7/CFTR. Its use to assemble virus is described below.

35

2. Virus Preparation from DNA - To generate the recombinant Ad2/CFTR-1 adenovirus, the vector pBR-Ad2-7/CFTR was cleaved with BstBI at the site corresponding to the unique BstBI site at 10670 in Ad2. The cleaved plasmid DNA was ligated to BstBI restricted Ad2

DNA. Following ligation, the reaction was used to transfect 293 cells by the calcium phosphate procedure. Approximately 7-8 days following transfection, a single plaque appeared and was used to reinfect a dish of 293 cells. Following development of cytopathic effect (CPE), the medium was removed and saved. Total DNA was prepared from the infected cells and analyzed by restriction analysis with multiple enzymes to verify the integrity of the construct. Viral supernatant was then used to infect 293 cells and upon development of CPE, expression of CFTR was assayed by the protein kinase A (PKA) immunoprecipitation assay (Gregory, R.J. et al. (1990) *Nature* 347:382). Following these verification procedures, the virus was further purified by two rounds of plaque purification.

Plaque purified virus was grown into a small seed stock by inoculation at low multiplicities of infection onto 293 cells grown in monolayers in 925 medium supplemented with 10% bovine calf serum. Material at this stage was designated a Research Viral Seed Stock (RVSS) and was used in all preliminary experiments.

3. Virus Host Cell - Ad2/CFTR-1 is propagated in human 293 cells (ATCC CRL 1573). These cells are a human embryonal kidney cell line which were immortalized with sheared fragments of human Ad5 DNA. The 293 cell line expresses adenovirus early region 1 gene products and in consequence, will support the growth of E1 deficient adenoviruses. By analogy with retroviruses, 293 cells could be considered a packaging cell line, but they differ from usual retrovirus lines in that they do not provide missing viral structural proteins, rather, they provide only some missing viral early functions.

Production lots of virus are propagated in 293 cells derived from the Working Cell Bank (WCB). The WCB is in turn derived from the Master Cell Bank (MCB) which was grown up from a fresh vial of cells obtained from ATCC. Because 293 cells are of human origin, they are being tested extensively for the presence of biological agents. The MCB and WCB are being characterized for identity and the absence of adventitious agents by Microbiological Associates, Rockville, MD.

4. Growth of Production Lots of Virus

Production lots of Ad2/CFTR-1 are produced by inoculation of approximately $5-10 \times 10^7$ pfu of MVSS onto approximately $1-2 \times 10^7$ Wcb 293 cells grown in a T175 flask containing 25 mls of 925 medium. Inoculation is achieved by direct addition of the virus (approximately 2-5 mls) to each flask. Batches of 50-60 flasks constitute a lot.

Following 40-48 hours incubation at 37°C, the cells are shaken loose from the flask and transferred with medium to a 250 ml centrifuge bottle and spun at 1000 xg. The cell pellet is resuspended in 4 ml phosphate buffered saline containing 0.1 g/l CaCl_2 and 0.1g/l MgCl_2 and the cells subjected to cycles of freeze-thaw to release virus. Cellular debris is removed by centrifugation at 1000 xg for 15 min. The supernatant from this centrifugation is layered on top of the CsCl step gradient: 2 ml 1.4g/ml CsCl and 3 ml 1.25g/ml CsCl in 10

mM Tris, 1 mM EDTA (TE) and spun for 1 hour at 35,000 rpm in a Beckman SW41 rotor. Virus is then removed from the interface between the two CsCl layers, mixed with 1.35 g/ml CsCl in TE and then subjected to a 2.5 hour equilibrium centrifugation at 75,000 rpm in a TLN-100 rotor. Virus is removed by puncturing the side of the tube with a hypodermic
5 needle and gently removing the banded virus. To reduce the CsCl concentration, the sample is dialyzed against 2 changes of 2 liters of phosphate buffered saline with 10% sucrose.

Following this procedure, dialyzed virus is stable at 4°C for several weeks or can be stored for longer periods at -80°C. Aliquots of material for human use will be tested and while awaiting the results of these tests, the remainder will be stored frozen. The tests to be
10 performed are described below:

5. Structure and Purity of Virus

SDS polyacrylamide gel electrophoresis of purified virions reveals a number of polypeptides, many of which have been characterized. When preparations of virus were
15 subjected to one or two additional rounds of CsCl centrifugation, the protein profile obtained was indistinguishable. This indicates that additional equilibrium centrifugation does not purify the virus further, and may suggest that even the less intense bands detected in the virus preparations represent minor virion components rather than contaminating proteins. The identity of the protein bands is presently being established by N-terminal sequence analysis.

20

6. Contaminating Materials - The material to be administered to patients will be 2×10^6 pfu, 2×10^7 pfu and 5×10^7 pfu of purified Ad2/CFTR-1. Assuming a minimum particle to pfu ratio of 500, this corresponds to 1×10^9 , 1×10^{10} and 2.5×10^{10} viral particles, these correspond to a dose by mass of 0.25 µg, 2.5 µg and 6.25 µg assuming a molecular mass for
25 adenovirus of 150×10^6 .

The origin of the materials from which a production lot of the purified Ad2/CFTR-1 is derived was described in detail above and is illustrated as a flow diagram in Figure 6. All the starting materials from which the purified virus is made (i.e., MCB, and WCB, and the MVSS) will be extensively tested. Further, the growth medium used will be tested and the
30 serum will be from only approved suppliers who will provide test certificates. In this way, all the components used to generate a production lot will have been characterized. Following growth, the production lot virus will be purified by two rounds of CsCl centrifugation, dialyzed, and tested. A production lot should constitute $1-5 \times 10^{10}$ pfu Ad2/CFTR-1.

As described above, to detect any contaminating material aliquots of the production
35 lot will be analyzed by SDS gel electrophoresis and restriction enzyme mapping. However, these tests have limited sensitivity. Indeed, unlike the situation for purified single chain recombinant proteins, it is very difficult to quantitate the purity of the AD2/CFTR-1 using SDS polyacrylamide gel electrophoresis (or similar methods). An alternative is the immunological detection of contaminating proteins (IDCP). Such an assay utilizes antibodies

raised against the proteins purified in a mock purification run. Development of such an assay has not yet been attempted for the CsCl purification scheme for Ad2/CFTR-1. However, initially an IDCP assay developed for the detection of contaminants in recombinant proteins produced in Chinese hamster ovary (CHO) cells will be used. In addition, to hamster
5 proteins, these assays detect bovine serum albumin (BSA), transferrin and IgG heavy and light chain derived from the serum added to the growth medium. Tests using such reagents to examine research batches of Ad2/CFTR-1 by both ELISA and Western blots are in progress.

Other proteins contaminating the virus preparation are likely to be from the 293 cells - that is, of human origin. Human proteins contaminating therapeutic agents derived from
10 human sources are usually not problematic. In this case, however, we plan to test the production lot for transforming factors. Such factors could be activities of contaminating human proteins or of the Ad2/CFTR-1 vector or other contaminating agents. For the test, it is proposed that 10 dishes of Rat 1 cells containing 2×10^6 cells (the number of target cells in the patient) with 4 times the highest human dose of Ad2/CFTR-1 (2×10^8 pfu) will be
15 infected. Following infection, the cells will be plated out in agar and examined for the appearance of transformed foci for 2 weeks. Wild type adenovirus will be used as a control.

Nucleic acids and proteins would be expected to be separated from purified virus preparations upon equilibrium density centrifugation. Furthermore, the 293 cells are not expected to contain VL30 sequences. Biologically active nucleic cells should be detected.
20

Example 8 - Preliminary Experiments Testing the Ability of Ad2/ β Gal or Ad2/CFTR Virus to Enter Airway Epithelial Cells

a. Hamster Studies

25 Initial studies involving the intratracheal instillation of the Ad- β Gal viral vector into Syrian hamsters, which are reported to be permissive for human adenovirus are being performed. The first study, a time course assessment of the pulmonary and systemic acute inflammatory response to a single intratracheal administration of Ad- β Gal viral vector, has been completed. In this study, a total of 24 animals distributed among three treatment
30 groups, specifically, 8 vehicle control, 8 low dose virus (1×10^{11} particles; 3×10^8 pfu), and 8 high dose virus (1.7×10^{12} particles; 5×10^9 pfu), were used. Within each treatment group, 2 animals were analyzed at each of four time points after viral vector instillation: 6 hrs, 24 hrs, 48 hrs, and 7 days. At the time of sacrifice of each animal, lung lavage and blood samples were taken for analysis. The lungs were fixed and processed for normal light-level
35 histology. Blood and lavage fluid were evaluated for total leukocyte count and leukocyte differential. As an additional measure of the inflammatory process, lavage fluid was also evaluated for total protein. Following embeddings, sectioning and hematoxylin/eosin staining, lung sections were evaluated for signs of inflammation and airway epithelial damage.

With the small sample size, the data from this preliminary study were not amenable to statistical analyses, however, some general trends could be ascertained. In the peripheral blood samples, total leukocyte counts showed no apparent dose- or time- dependent changes. In the blood leukocyte differential counts, there may have been a minor dose-related
5 elevation in percent neutrophil at 6 hours; however, data from all other time points showed no elevation in neutrophil percentages. Taken together, these data suggest little or nor systemic inflammatory response to the viral administration.

From the lung lavage, some elevation in total neutrophil counts were observed at the first three time points (6 hr, 24 hr, 48 hr). By seven days, both total and percent neutrophil
10 values had returned to normal range. The trends in lung lavage protein levels were more difficult to assess due to inter-animal variability; however, no obvious dose- or time- dependent effects were apparent. First, no damage to airway epithelium was observed at any time point or virus dose level. Second, a time- and dose- dependent mild inflammatory response was observed, being maximal at 48 hr in the high virus dose animals. By seven
15 days, the inflammatory response had completely resolved, such that the lungs from animals in all treatment groups were indistinguishable.

In summary, a mild, transient, pulmonary inflammatory response appears to be associated with the intratracheal administration of the described doses of adenoviral vector in the Syrian Hamster.

20 A second, single intratracheal dose, hamster study has been initiated. This study is designed to assess the possibility of the spread of ineffective viral vectors to organs outside of the lung and the antibody response of the animals to the adenoviral vector. In this study, the three treatment groups (vehicle control, low dose virus, high dose virus) each contained 12 animals. Animals will be evaluated at three time points: 1 day, 7 days, and 1 month. In this
25 study, viral vector persistence and possible spread will be evaluated by the assessment of the presence of infective virions in numerous organs including lung, gut, heart, liver, spleen, kidney, brain and gonads. Changes in adenoviral antibody titer will be measured in peripheral blood and lung lavage. Additionally, lung lavage, peripheral blood and lung histology will be evaluated as in the previous study.

30

b. Primate studies.

Studies of recombinant adenovirus are also underway in primates. The goal of these studies is to assess the ability of recombinant adenoviral vectors to deliver genes to the respiratory epithelium *in vivo* and to assess the safety of the construct in primates. Initial
35 studies in primates targeted nasal epithelia as the site of infection because of its similarity to lower airway epithelia, because of its accessibility, and because nasal epithelia was used for the first human studies. The Rhesus monkey (*Macaca mulatta*) has been chosen for studies, because it has a nasal epithelium similar to that of humans.

How expression of CFTR affects the electrolyte transport properties of the nasal epithelium can be studied in patients with cystic fibrosis. But because the primates have normal CFTR function, instead the ability to transfer a reporter gene was assessed. Therefore the Ad- β Gal virus was used. The epithelial cell density in the nasal cavity of the Rhesus monkey is estimated to be 2×10^6 cells/cm (based on an average nasal epithelial cell diameter of 7 μ m) and the surface near 25-50 cm². Thus, there are about 5×10^7 cells in the nasal epithelium of Rhesus monkey. To focus especially on safety, the higher viral doses (20-200 MOI) were used *in vivo*. Thus doses in the range of 10^9 - 10^{10} pfu were used.

In the first pilot study the right nostril of Monkey A was infected with Ad- β -Gal (~1 ml). This viral preparation was purified by CsCl gradient centrifugation and then by gel filtration chromatography one week later. Adenoviruses are typically stable in CsCl at 4°C for one to two weeks. However, this viral preparation was found to be defective (i.e., it did not produce detectable β -galactosidase activity in the permissive 293 cells). Thus, it was concluded that there was no live viral activity in the material. β -galactosidase activity in nasal epithelial cells from Monkey A was also not detected. Therefore, in the next study, two different preparations of Ad- β -Gal virus: one that was purified on a CsCl gradient and then dialyzed against Tris-buffered saline to remove the CsCl, and a crude unpurified one was used. Titers of Ad- β -Gal viruses were $\sim 2 \times 10^{10}$ pfu/ml and $> 1 \times 10^{13}$ pfu/ml, respectively, and both preparations produced detectable β -galactosidase activity in 293 cells.

Monkeys were anesthetized by intramuscular injection of ketamine (15 mg/kg). One week before administration of virus, the nasal mucosa of each monkey was brushed to establish baseline cell differentials and levels of β -galactosidase. Blood was drawn for baseline determination of cell differentials, blood chemistries, adenovirus antibody titers, and viral cultures. Each monkey was also examined for weight, temperature, appetite, and general health prior to infection.

The entire epithelium of one nasal cavity was used in each monkey. A foley catheter (size 10) was inserted through each nasal cavity into the pharynx, inflated with 2-3 ml of air, and then pulled anteriorly to obtain tight posterior occlusion at the posterior choana. Both nasal cavities were then irrigated with a solution (~5 ml) of 5 mM dithiothreitol plus 0.2 U/ml neuraminidase in phosphate-buffered saline (PBS) for five minutes. This solution was used to dissolve any residual mucus overlaying the epithelia. (It was subsequently found that such treatment is not required.) The washing procedure also allowed the determination of whether the balloons were effectively isolating the nasal cavity. The virus (Ad- β -Gal) was then slowly instilled into the right nostril with the posterior balloon inflated. The viral solution remained in contact with the nasal mucosa for 30 minutes. At the end of 30 minutes, the remaining viral solution was removed by suction. The balloons were deflated, the catheters removed, and the monkey allowed to recover from anesthesia. Monkey A received the CsCl-purified virus (~1.5 ml) and Monkey B received the crude virus (~6 ml). (note that this was the second exposure of Monkey A to the recombinant adenovirus).

Both monkeys were followed daily for appearance of the nasal mucosa, conjunctivitis, appetite, activity, and stool consistency. Each monkey was subsequently anesthetized on days 1, 4, 7, 14, and 21 to obtain nasal, pharyngeal, and tracheal cell samples (either by swabs or brushes) as described below. Phlebotomy was performed over the same time course for hematology, ESR, general screen, antibody serology and viral cultures. Stools were collected every week to assess viral cultures.

To obtain nasal epithelial cells from an anesthetized monkey, the nasal mucosa was first impregnated with 5 drops of Afrin (0.05% oxymetazoline hydrochloride, Schering-Plough) and 1 ml of 2% Lidocaine for 5 min. A cytobrush (the kind typically used for Pap smears) was then used to gently rub the mucosa for about 10 seconds. For tracheal brushings, a flexible fiberoptic bronchoscope; a 3 mm cytology brush (Bard) was advanced through the bronchoscope into the trachea, and a small area was brushed for about 10 seconds. This procedure was repeated twice to obtain a total of $\sim 10^6$ cells/ml. Cells were then collected on slides (approximately 2×10^4 cells/slide using a Cytospin 3 (Shandon, PA)) for subsequent staining (see below).

To determine viral efficacy, nasal, pharyngeal, and tracheal cells were stained for β -galactosidase using X-gal (5-bromo-4-chloro-3-indolyl- β -D-galactoside). Cleavage of X-gal by β -galactosidase produces a blue color that can be seen with light microscopy. The Ad- β -gal vector included a nuclear-localization signal (NLS) (from SV40 large T-antigen) at the amino-terminus of the β -galactosidase sequence to direct expression of this protein to the nucleus. Thus, the number of blue nuclei after staining was determined.

RT-PCR (reverse transcriptase-polymerase chain reaction) was also used to determine viral efficacy. This assay indicates the presence of β -galactosidase mRNA in cells obtained by brushings or swabs. PCR primers were used in both the adenovirus sequence and the LacZ sequence to distinguish virally-produced mRNA from endogenous mRNA. PCR was also used to detect the presence of the recombinant adenovirus DNA. Cytospin preparations were used to assess for the presence of virally produced β -galactosidase mRNA in the respiratory epithelial cells using *in-situ* hybridization. This technique has the advantage of being highly specific and will allow assessment which cells are producing the mRNA.

Whether there was any inflammatory response was assessed by visual inspection of the nasal epithelium and by cytological examination of Wright-stained cells (cytospin). The percentage of neutrophils and lymphocytes were compared to that of the control nostril and to the normal values from four control monkeys. Systemic responses by white blood cell counts, sedimentation rate, and fever were also assessed.

Viral replication at each of the time points was assessed by testing for the presence of live virus in the supernatant of the cell suspension from swabs or brushes. Each supernatant was used to infect (at several dilutions) the virus-sensitive 293 cell line. Cytopathic changes in the 293 cells were monitored for 1 week and then the cells were fixed and stained for β -galactosidase. Cytopathic effects and blue-stained cells indicated the presence of live virus.

Positive supernatants will also be subjected to analysis of nonintegrating DNA to identify (confirm) the contributing virus(es).

Antibody titers to type 2 adenovirus and to the recombinant adenovirus were determined by ELISA. Blood/serum analysis was performed using an automated chemistry analyzer Hitachi 737 and an automated hematology analyzer Technicom H6. The blood buffy coat was cultured in A549 cells for wild type adenovirus and was cultured in the permissive 293 cells.

Results: Both monkeys tolerated the procedure well. Daily examination revealed no evidence of coryza, conjunctivitis or diarrhea. For both monkeys, the nasal mucosa was mildly erythematous in both the infection side and the control side; this was interpreted as being due to the instrumentation. Appetites and weights were not affected by virus administered in either monkey. Physical examination on days 1, 4, 7, 14 and 21 revealed no evidence of lymphadenopathy, tachypnea, or tachycardia. On day 21, monkey B had a temperature 39.1°C (normal for Rhesus monkey 38.8°C) but had no other abnormalities on physical exam or in laboratory data. Monkey A had a slight leukocytosis on day 1 post infection which returned to normal by day 4; the WBC was 4,920 on the day of infection, 8,070 on day 1, and 5,200 on day 4. The ESR did not change after the infection. Electrolytes and transaminases were normal throughout.

Wright stains of cells from nasal brushing were performed on days 4, 7, 14, and 21. They revealed less than 5% neutrophils and lymphocytes. There was no difference between the infected and the control side.

X-Gal stains of the pharyngeal swabs revealed blue-stained cells in both monkeys on days 4, 7, and 14; only a few of the cells had clear nuclear localization of the pigment and some pigment was seen in extracellular debris. On day 7 post infection, X-Gal stains from the right nostril of monkey A, revealed a total of 135 ciliated cells with nuclear-localized blue stain. The control side had only 4 blue cells. Monkey B had 2 blue cells from the infected nostril and none from the control side. Blue cells were not seen on day 7, 14, or 21.

RT-PCR on day 3 post infection revealed a band of the correct size that hybridized with a β -Gal probe, consistent with β -Gal mRNA in the samples from Monkey A control nostril and Monkey B infected nostril. On day 7 there was a positive band in the sample from the infected nostril of Monkey A, the same specimen that revealed blue cells.

Fluid from each nostril, the pharynx, and trachea of both monkeys was placed on 293 cells to check for the presence of live virus by cytopathic effect and X-Gal stain. In Monkey A, live virus was detected in both nostrils on day 3 after infection; no live virus was detected at either one or two weeks post-infection. In Monkey B, live virus was detected in both nostrils, pharynx, and trachea on day 3, and only in the infected nostril on day 7 after infection. No live virus was detected 2 weeks after the infection.

c. Human Explant Studies

In a second type of experiment, epithelial cells from a nasal polyp of a CF patient were cultured on permeable filter supports. These cells form an electrically tight epithelial monolayer after several days in culture. Eight days after seeding, the cells were exposed to the Ad2/CFTR virus for 6 hours. Three days later, the short-circuit current (I_{sc}) across the monolayer was measured. cAMP agonists did not increase the I_{sc}, indicating that there was no change in chloride secretion. However, this defect was corrected after infection with recombinant Ad2/CFTR. Cells infected with Ad2/CFTR (MOI=5; MOI refers to multiplicity of infection; 1 MOI indicates one pfu/cell) express functional CFTR; cAMP agonists stimulated I_{sc}, indicating stimulation of Cl⁻ secretion. Ad2/CFTR also corrected the CF chloride channel defect in CF tracheal epithelial cells. Additional studies indicated that Ad2/CFTR was able to correct the chloride secretory defect without altering the transepithelial electrical resistance; this result indicates that the integrity of the epithelial cells and the tight junctions was not disrupted by infection with Ad2/CFTR. Application of 1 MOI of Ad2/CFTR was also found to be sufficient to correct the CF chloride secretory defect.

The experiments using primary cultures of human airway epithelial cells indicate that the Ad2/CFTR virus is able to enter CF airway epithelial cells and express sufficient CFTR to correct the defect in chloride transport.

20 Example 9 -In Vivo Delivery to and Expression of CFTR in Cotton Rat and Rhesus Monkey Epithelium

MATERIALS AND METHODS

Adenovirus vector

25 Ad2/CFTR-1 was prepared as described in Example 7. The DNA construct comprises a full length copy of the Ad2 genome of approximately 37.5 kb from which the early region 1 genes (nucleotides 546 to 3497) have been replaced by cDNA for CFTR (nucleotides 123 to 4622 of the published CFTR sequence with 53 additional linker nucleotides). The viral Ela promoter was used for CFTR cDNA. Termination/polyadenylation occurs at the site normally used by the Elb and protein IX transcripts. The recombinant virus E3 region was conserved. The size of the Ad2-CFTR-1 vector is approximately 104.5% that of wild-type adenovirus. The recombinant virus was grown in 293 cells that complement the E1 early viral promoters. The cells were frozen and thawed three times to release the virus and the preparation was purified on a CsCl gradient, then dialyzed against Tris-buffered saline (TBS) to remove the CsCl, as described.

Animals

Rats. Twenty two cotton rats (6-8 weeks old, weighing between 80-100 g) were used for this study. Rats were anesthetized by inhaled methoxyflurane (Pitman Moore, Inc., Mundelen, Ill). Virus was applied to the lungs by nasal instillation during inspiration.

5 Two cotton rat studies were performed. In the first study, seven rats were assigned to a one time pulmonary infection with 100 μ l solution containing 4.1×10^9 plaque forming units (pfu) of the Ad2/CFTR-1 virus and 3 rats served as controls. One control rat and either two or three experimental rats were sacrificed with methoxyflurane and studies at each of three time points: 4, 11, or 15 days after infection.

10 The second group of rats was used to test the effect of repeat administration of the recombinant virus. All 12 rats received 2.1×10^8 pfu of the Ad2/CFTR-1 virus on day 0 and 9 of the rats received a second dose of 3.2×10^8 pfu of Ad2/CFTR-1 14 days later. Groups of one control rat and three experimental rats were sacrificed at 3, 7, or 14 days after the second administration of virus. Before necropsy, the trachea was cannulated and
15 brochoaveolar lavage (BAL) was performed with 3 ml aliquots of phosphate-buffered saline. A median sternotomy was performed and the right ventricle cannulated for blood collection. The right lung and trachea were fixed in 4% formaldehyde and the left lung was frozen in liquid nitrogen and kept at -70°C for evaluation by immunochemistry, reverse transcriptase polymerase chain reaction (RT-PCR), and viral culture. Other organs were removed and
20 quickly frozen in liquid nitrogen for evaluation by polymerase chain reaction (PCR).

Monkeys. Three female Rhesus monkeys were used for this study; a fourth female monkey was kept in the same room, and was used as control. For application of the virus, the monkeys were anesthetized by intramuscular injection of ketamine (15 mg/kg). The entire epithelium of one nasal cavity in each monkey was used for virus application. A foley
25 catheter (size 10) was inserted through each nasal cavity into the pharynx, the balloon was inflated with 2-3 ml of air, and then pulled anteriorly to obtain a tight occlusion at the posterior choana. The Ad2/CFTR-1 virus was then instilled slowly in the right nostril with the posterior balloon inflated. The viral solution remained in contact with the nasal mucosa for 30 min. The balloons were deflated, the catheters were removed, and the monkeys were
30 allowed to recover from anesthesia. A similar procedure was performed on the left nostril, except that TBS solution was instilled as a control. The monkeys received a total of three doses of the virus over a period of 5 months. The total dose given was 2.5×10^9 pfu the first time, 2.3×10^9 pfu the second time, and 2.8×10^9 pfu the third time. It was estimated that the cell density of the nasal epithelia to be 2×10^6 cells/cm² and a surface area of 25 to 50
35 cm². This corresponds to a multiplicity of infection (MOI) of approximately 25.

The animals were evaluated 1 week before the first administration of virus, on the day of administration, and on days 1, 3, 6, 13, 21, 27, and 42 days after infection. The second administration of virus occurred on day 55. The monkeys were evaluated on day 55 and then on days 56, 59, 62, 69, 76, 83, 89, 96, 103, and 111. For the third administration, on day 134,

only the left nostril was cannulated and exposed to the virus. The control monkey received instillations of PBS instead of virus. Biopsies of the left medial turbinate were carried out on day 135 in one of the infected monkeys, on day 138 on the second infected monkey, and on day 142 on the third infected monkey and on the control monkey.

5 For evaluations, monkeys were anesthetized by intramuscular injection of ketamine (15 mg/kg). To obtain nasal epithelial cells, the nasal mucosa was first impregnated with 5 drops of Afrin (0.05% oxymetazoline hydrochloride, Schering-Plough) and 1 ml of 2% Lidocaine for 5 minutes. A cytobrush was then used to gently rub the mucosa for about 3 sec. To obtain pharyngeal epithelial swabs, a cotton-tipped applicator was rubbed over the
10 back of the pharynx 2-3 times. The resulting cells were dislodged from brushes or applicators into 2 ml of sterile PBS. Biopsies of the medial turbinate were performed using cupped forceps under direct endoscopic control.

Animals were evaluated daily for evidence of abnormal behavior or physical signs. A record of food and fluid intake was used to assess appetite and general health. Stool
15 consistency was also recorded to check for the possibility of diarrhea. At each of the evaluation time points, rectal temperature, respiratory rate, and heart rate were measured. The nasal mucosa, conjunctivas, and pharynx were visually inspected. The monkeys were also examined for lymphadenopathy.

Venous blood from the monkeys was collected by standard venipuncture technique.
20 Blood/serum analysis was performed in the clinical laboratory of the University of Iowa Hospitals and Clinics using a Hitachi 737 automated chemistry analyzer and a Technicom H6 automated hematology analyzer.

Serology

25 Sera were obtained and anti-adenoviral antibody titers were measured by an enzyme-linked immunoabsorbant assay (ELISA). For the ELISA, 50 ng/well of filled adenovirus (Lee Biomolecular Research Laboratories, San Diego, Ca) in 0.1M NaHCO₃ were coated on 96 well plates at 4°C overnight. The test samples at appropriate dilutions were added, starting at a dilution of 1/50. The samples were incubated for 1 hour, the plates washed, and
30 a goat anti-human IgG HRP conjugate (Jackson ImmunoResearch Laboratories, West Grove, PA) was added and incubated for 1 hour. The plates were washed and O-Phenylenediamine (Sigma Chemical Co., St. Louis, MO) was added for 30 min. at room temperature. The assay was stopped with 4.5 M H₂SO₄ and read at 490 nm on a Molecular Devices microplate reader. The titer was calculated as the product of the reciprocal of the initial dilution and the
35 reciprocal of the dilution in the last well with an OD>0.100.

Neutralizing antibodies measure the ability of the monkey serum to prevent infection of 293 cells by adenovirus. Monkey serum (1:25 dilution) [or nasal washings (1:2 dilutions)] was added in two-fold serial dilutions to a 96 well plate. Adenovirus (2.5 x 10⁵ pfu) was added and incubated for 1 hour at 37°C. The 293 cells were then added to all wells and the

plates were incubated until the serum-free control wells exhibited >95% cytopathic effect. The titer was calculated as the product of the reciprocal of the initial dilution times the reciprocal of the dilution in the last well showing >95% cytopathic effect.

5 Bronchoalveolar lavage and nasal brushings for cytology

Bronchoalveolar lavage (BAL) was performed by cannulating the trachea with a silastic catheter and injecting 5 ml of PBS. Gentle suction was applied to recover the fluid. The BAL sample was spun at 5000 rpm for 5 min. and cells were resuspended in 293 media at a concentration of 10^6 cells/ml. Cells were obtained from the monkey's nasal epithelium by gently rubbing the nasal mucosa for about 3 sec. with a cytobrush. The resulting cells were dislodged from the brushes into 2 ml of PBS. Forty microliters of the cell suspension were cytocentrifuged onto slides and stained with Wright's stain. Samples were examined by light microscopy.

15

Histology of lung sections and nasal biopsies

The right lung of each cotton rat was removed, inflated with 4% formaldehyde, and embedded in paraffin for sectioning. Nasal biopsies from the monkeys were also fixed with 4% formaldehyde. Histologic sections were stained with hematoxylin and eosin (H&E). Sections were reviewed by at least one of the study personnel and by a pathologist who was unaware of the treatment each rat received.

20

Immunocytochemistry

Pieces of lung and trachea of the cotton rats and nasal biopsies were frozen in liquid nitrogen on O.C.T. compound. Cryosections and paraffin sections of the specimens were used for immunofluorescence microscopy. Cytospin slides of nasal brushings were prepared on gelatin coated slides and fixed with paraformaldehyde. The tissue was permeabilized with Triton X-100, then a pool of monoclonal antibodies to CFTR (M13-1, M1-4) (Denning, G.M. et al. (1992) *J. Clin. Invest.* 89:339-349) was added and incubated for 12 hours. The primary antibody was removed and an anti-mouse biotinylated antibody (Biomed, Foster City, CA) was added. After removal of the secondary antibody, streptavidin FITC (Biomed, Foster City, Ca) was added and the slides were observed under a laser scanning confocal microscope. Both control animal samples and non-immune IgG stained samples were used as controls.

35

PCR

PCR was performed on pieces of small bowel, brain, heart, kidney, liver, ovaries, and spleen from cotton rats. Approximately 1 g of the rat organs was mechanically ground and mixed with 50 μ l sterile water, boiled for 5 min., and centrifuged. A 5 μ l aliquot of the

supernatant was removed for further analysis. Monkey nasal brushings suspensions were also used for PCR.

Nested PCR primer sets were designed to selectively amplify Ad2/CFTR-1 DNA over endogenous CFTR by placing one primer from each set in the adenovirus sequence and the other primer in the CFTR sequence. The first primer set amplifies a 723 bp fragment and is shown below:

Ad2 5' ACT CTT GAG TGC CAG CGA GTA GAG TTT TCT CCT CCG 3' (SEQ ID NO:4)

CFTR 5' GCA AAG GAG CGA TCC ACA CGA AAT GTG CC 3' (SEQ ID NO:5)

The nested primer set amplifies a 506 bp fragment and is shown below:

Ad2 5' CTC CTC CGA GCC GCT CCG AGC TAG 3' (SEQ ID NO:6)

CFTR 5' CCA AAA ATG GCT GGG TGT AGG AGC AGT GTC C 3' (SEQ ID NO:7)

A PCR reaction mix containing 10mM Tris-Cl (pH 8.3), 50mM KCl, 1.5 mM MgCl₂, 0.001% (w/v) gelatin, 400 µM each dNTP, 0.6 µM each primer (first set), and 2.5 units AmpliTaq (Perkin Elmer) was aliquoted into separate tubes. A 5 µl aliquot of each sample prep was then added and the mixture was overlaid with 50 µl of light mineral oil. The samples were processed on a Barnstead/Thermolyne (Dubuque, IA) thermal cycler programmed for 1 min. at 94°C, 1 min. at 65°C, and 2 min. at 72°C for 40 cycles. Post-run dwell was for 7 min. at 72°C. A 5 µl aliquot was removed and added to a second PCR reaction using the nested set of primers and cycled as above. A 10 µl aliquot of the final amplification reaction was analyzed on a 1% agarose gel and visualized with ethidium bromide.

To determine the sensitivity of this procedure, a PCR mix containing control rat liver supernatant was aliquoted into several tubes and spiked with dilutions of Ad2/CFTR-1. Following the amplification protocols described above, it was determined that the nested PCR procedure could detect as little as 50 pfu of viral DNA.

RT-PCR

RT-PCR was used to detect vector-generated mRNA in cotton rat lung tissue and samples from nasal brushings from monkeys. A 200 µl aliquot of guanidine isothiocyanate solution (4 M guanidine isothiocyanate, 25 mM sodium citrate pH 7.0, 0.5% sarcosyl, and 0.1 M β-mercaptoethanol) was added to a frozen section of each lung and pellet from nasal brushings and the tissue was mechanically ground. Total RNA was isolated utilizing a single-step method (Chomczynski, P. and Sacchi, N. et al. (1987) *Analytical Biochemistry* 162:156-159; Hanson, C.A. et al. (1990) *Am. J. Pathol.* 137:1-6). The RNA was incubated with 1 unit RQ1 RNase-free DNase (Promega Corp., Madison WI) at 37°C for 20 min., denatured at 99°C for 5 min., precipitated with ammonium acetate and ethanol, and redissolved in 4 µl diethylpyrocarbonate treated water containing 20 units RNase Block 1 (Stratagene, La Jolla CA). A 2 µl aliquot of the purified RNA was reverse transcribed using

the GeneAmp RNA PCR kit (Perkin Elmer Cetus) and the downstream primer from the first primer set described in the previous section. Reverse transcriptase was omitted from the reaction with the remaining 2 µl of the purified RNA prep, as a control in which preparations (both +/- RT) were then amplified using nested primer sets and the PCR protocols described above. A 10 µl aliquot of the final amplification reaction was analyzed on a 1% agarose gel and visualized with ethidium bromide.

Southern analysis.

To verify the identity of the PCR products, Southern analysis was performed. The DNA was transferred to a nylon membrane as described (Sambrook *et al.*, *supra*). A fragment of CFTR cDNA (amino acids #1-525) was labeled with [³²P]-dCTP (ICN Biomedicals, Inc. Irvine CA) using an oligolabeling kit (Pharmacia, Piscataway, NJ) and purified over a NICK column (Pharmacia Piscataway, NJ) for use as a hybridization probe. The labeled probe was denatured, cooled, and incubated with the prehybridized filter for 15 hours at 42°C. The hybridized filter was then exposed to film (Kodak XAR-5) for 10 min.

Culture of Ad2/CFTR-1

Viral cultures were performed on the permissive 293 cell line. For culture of virus from lung tissue, 1 g of lung was frozen/thawed 3-6 times and then mechanically disrupted in 200 µl of 293 media. For culture of BAL and monkey nasal brushings, the cell suspension was spun for 5 min and the supernatant was collected. Fifty µl of the supernatant was added in duplicate to 293 cells grown in 96 well plates at 50% confluence. The 293 cells were incubated for 72 hr at 37°C, then fixed with a mixture of equal parts of methanol and acetone for 10 min. and incubated with FITC-labeled anti-adenovirus monoclonal antibodies (Chemicon, Light Diagnostics, Temecuca, CA) for 30 min. Positive nuclear immunofluorescence was interpreted as positive culture. The sensitivity of the assay was evaluated by adding dilutions of Ad2/CFTR-1 to 50 µl of the lung homogenate from one of the control rats. Viral replication was detected when as little as 1 pfu was added.

RESULTS

Efficacy of Ad2/CFTR-1 in the lungs of cotton rats.

To test the ability of Ad2/CFTR-1 to transfer CFTR cDNA to the intrapulmonary airway epithelium, several studies were performed. 4 x 10⁶ pfu - IU of Ad2/CFTR-1 in 100 µl was administered to seven cotton rats; three control rats received 100 µl of TBS (the vehicle for the virus). The rats were sacrificed 4, 10 or 14 days later. To detect viral transcripts encoding CFTR, reverse transcriptase was used to prepare cDNA from lung homogenates. The cDNA was amplified with PCR using primers that span adenovirus and CFTR-encoded

sequences. Thus, the procedure did not detect endogenous rat CFTR. Figure 16 shows that the lungs of animals which received Ad2/CFTR-1 were positive for virally-encoded CFTR mRNA. The lungs of all control rats were negative.

To detect the protein, lung sections were immunostained with antibodies specific to CFTR. CFTR was detected at the apical membrane of bronchial epithelium from all rats exposed to Ad2/CFTR-1, but not from control rats. The location of recombinant CFTR at the apical membrane is consistent with the location of endogenous CFTR in human airway epithelium. Recombinant CFTR was detected above background levels because endogenous levels of CFTR in airway epithelia are very low and thus, difficult to detect by immunocytochemistry (Trapnell, B. et al. (1991) *Proc. Natl. Acad. Sci. USA* 88:6565-6569; Denning, G.M. et al. (1992) *J. Cell Biol.* 118:551-59).

These results show that Ad2/CFTR-1 directs the expression of CFTR mRNA in the lung of the cotton rat and CFTR protein in the intrapulmonary airways.

15 Safety of Ad2/CFTR-1 in cotton rats.

Because the E1 region of Ad2 is deleted in the Ad2/CFTR-1 virus, the vector was expected to be replication-impaired (Berkner, K.L. (1988) *BioTechniques* 6:616-629) and that it would be unable to shut off host cell protein synthesis (Basuss, L.E. et al. (1989) *J. Virol.* 50:202-212). Previous *in vitro* studies have suggested that this is the case in a variety of cells including primary cultures of human airway epithelial cells (Rich, D.P. et al. (1993) *Human Gene Therapy* 4:461-476). However, it is important to confirm this *in vivo* in the cotton rat, which is the most permissive animal model for human adenovirus infection (Ginsberg, H.S. et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:3823-3827; Prince, G.A. et al. (1993) *J. Virol* 67:101-111). Although dose of virus of 4.1×10^{10} pfus per kg was used, none of the rats died. More importantly, extracts from lung homogenates from each of the cotton rats were cultured in the permissive 293 cell line. With this assay 1 pfu of recombinant virus was detected in lung homogenate. However, virus was not detected by culture in the lungs of any of the treated animals. Thus, the virus did not appear to replicate *in vivo*.

It is also possible that administration of Ad2/CFTR-1 could cause an inflammatory response, either due to a direct effect of the virus or as a result of administration of viral particles. Several studies were performed to test this possibility. None of the rats had a change in the total or differential white blood cell count, suggesting that there was no major systemic inflammatory response. To assess the pulmonary inflammatory response more directly, bronchoalveolar lavage was performed on each of the rats (Figures 17A and 17B). Figure 17A shows that there was no change in the total number of cells recovered from the lavage or in the differential cell count.

Sections of the lung stained by H&E were also prepared. There was no evidence of viral inclusions or any other changes characteristic of adenoviral infection (Prince, G.A. et al. (1993) *J. Virol.* 67:101-111). When coded lung sections were evaluated by a skilled reader

who was unaware of which sections were treated, she was unable to distinguish between sections from the treated and untreated lungs.

It seemed possible that the recombinant adenovirus could escape from the lung into other tissues. To test for this possibility, other organs from the rats were evaluated using nested PCR to detect viral DNA. All organs tested from infected rats were negative, with the exception of small bowel which was positive in 3 of 7 rats. Figure 18 shows the results of 2 infected rats and one control rat sacrificed on day 4 after infection. The organ homogenates from the infected rats sacrificed were negative for Ad2/CFTR-1 with the exception of the small bowel. Organ homogenates from control rats sacrificed on day 4 after infection were negative for Ad2/CFTR-1. The presence of viral DNA in the small bowel suggests that the rats may have swallowed some of the virus at the time of instillation or, alternatively, the normal airway clearance mechanisms may have resulted in deposition of viral DNA in the gastrointestinal tract. Despite the presence of viral DNA in homogenates of small intestine, none of the rats developed diarrhea. This result suggests that if the virus expressed CFTR in the intestinal epithelium, there was no obvious adverse consequence.

Repeat administration of Ad2/CFTR-1 to cotton rats

Because adenovirus DNA integration into chromosomal DNA is not necessary for gene expression and only occurs at very low frequency, expression following any given treatment was anticipated to be finite and that repeated administration of recombinant adenovirus would be required for treatment of CF airway disease. Therefore, the effect of repeated administration of Ad2/CFTR-1 cotton rats was examined. Twelve cotton rats received 50 μ l of Ad2/CFTR-1. Two weeks later, 9 of the rats received a second dose of 50 μ l of Ad2/CFTR-1 and 3 rats received 50 μ l of TBS. Rats were sacrificed on day 3, 7, or 14 after virus administration. At the time of the second vector administration all cotton rats had an increased antibody titer to adenovirus.

After the second intrapulmonary administration of virus, none of the rats died. Moreover, the results of studies assessing safety and efficacy were similar to results obtained in animals receiving adenovirus for the first time. Viral cultures of rat lung homogenates on 293 cells were negative at all time points, suggesting that there was no virus replication. There was no difference between treated and control rats in the total or differential white blood count at any of the time points. The lungs were evaluated by histologic sections stained with H&E; and found no observable differences between the control and treated rats when sections were read by us or by a blinded skilled reader. Examples of some sections are shown in Figure 19. When organs were examined for viral DNA using PCR, viral DNA was found only in the small intestine of 2 rats. Despite seropositivity of the rats at the time of the second administration, expression of CFTR (as assessed by RT-PCR and by immunocytochemistry of sections stained with CFTR antibodies) similar to that seen in animals that received a single administration was observed.

These results suggest that prior administration of Ad2/CFTR-1 and the development of an antibody response did not cause an inflammatory response in the rats nor did it prevent virus-dependent production of CFTR.

5 Evidence that Ad2/CFTR-1 expresses CFTR in primate airway epithelium

The cells lining the respiratory tract and the immune system of primates are similar to those of humans. To test the ability of Ad2/CFTR-1 to transfer CFTR to the respiratory epithelium of primates, Ad2/CFTR was applied on three occasions as described in the methods to the nasal epithelium of three Rhesus monkeys. To obtain cells from the
10 respiratory epithelium, the epithelium was brushed using a procedure similar to that used to sample the airway epithelium of humans during fiberoptic bronchoscopy.

To assess gene transfer, RT-PCR was used as described above for the cotton rats. RT-PCR was positive on cells brushed from the right nostril of all three monkeys, although it was only detectable for 18 days after virus administration. An example of the results are
15 shown in Figure 20A. The presence of a positive reaction in cells from the left nostril most likely represents some virus movement to the left side due to drainage, or possibly from the monkey moving the virus from one nostril to the other with its fingers after it recovered from anesthesia.

The specificity of the RT-PCR is shown in Figure 20B. A Southern blot with a probe
20 to CFTR hybridized with the RT-PCR product from the monkey infected with Ad2/CFTR-1. As a control, one monkey received a different virus (Ad2/ β Gal-1) which encodes β -galactosidase. When different primers were used to reverse transcribe the β -galactosidase mRNA and amplify the cDNA, the appropriate PCR product was detected. However, the PCR product did not hybridize to the CFTR probe on Southern blot. This result shows the
25 specificity of the reaction for amplification of the adenovirus-directed CFTR transcript.

The failure to detect evidence of adenovirus-encoded CFTR mRNA at 18 days or beyond suggests that the sensitivity of the RT-PCR may be low because of limited efficacy of the reverse transcriptase or because RNases may have degraded RNA after cell acquisition. Viral DNA, however, was detected by PCR in brushings from the nasal epithelium for
30 seventy days after application of the virus. This result indicates that although mRNA was not detected after 2 weeks, viral DNA was present for a prolonged period and may have been transcriptionally active.

To assess the presence of CFTR proteins directly, cells obtained by brushing were plated onto slides by cytopspin and stained with antibodies to CFTR. Figure 21 shows an
35 example of the immunocytochemistry of the brushed cells. A positive reaction is clearly evident in cells exposed to Ad2/CFTR-1. The cells were scored as positive by immunocytochemistry when evaluated by a reader uninformed to the identity of the samples. Immunocytochemistry remained positive for five to six weeks for the three monkeys, even after the second administration of Ad2/CFTR-1. On occasion, a few positive staining cells

were observed from the contralateral nostril of the monkeys. However, this was of short duration, lasting at most one week.

Sections of nasal turbinate biopsies obtained within a week after the third infection were also examined. In sections from the control monkey, little if any immunofluorescence from the surface epithelium was observed, but the submucosal glands showed significant staining of CFTR (Fig. 22). These observations are consistent with results of previous studies (Engelhardt, J.F. and Wilson, J.M. (1992) *Nature Gen.* 2:240-248.) In contrast, sections from monkeys that received Ad2/CFTR-1 revealed increased immunofluorescence at the apical membrane of the surface epithelium. The submucosal glands did not appear to have greater immunostraining than was observed under control conditions. These results indicate that Ad2/CFTR-1 can transfer the CFTR cDNA to the airway epithelium of Rhesus monkeys, even in seropositive animals (see below).

Safety of Ad2/CFTR-1 administered to monkeys

Figure 23 shows that all three treated monkeys developed antibodies against adenovirus. Antibody titers measured by ELISA rose within two weeks after the first infection. With subsequent infections the titer rose within days. The sentinel monkey had low antibody titers throughout the experiment. Tests for the presence of neutralizing antibodies were also performed. After the first administration, neutralizing antibodies were not observed, but they were detected after the second administration and during the third viral administration (Fig. 23).

To detect virus, supernatants from nasal brushings and swabs were cultured on 293 cells. All monkeys had positive cultures on day 1 and on day 3 or 4 from the infected nostril. Cultures remained positive in one of the monkeys at seven days after administration, but cultures were never positive beyond 7 days. Live virus was occasionally detected in swabs from the contra lateral nostril during the first 4 days after infection. The rapid loss of detectable virus suggests that there was not viral replication. Stools were routinely cultured, but virus was never detected in stools from any of the monkeys.

None of the monkeys developed any clinical signs of viral infection or inflammation. Visual inspection of the nasal epithelium revealed slight erythema in all three monkeys in both nostrils on the first day after infection; but similar erythema was observed in the control monkey and likely resulted from the instrumentation. There was no visible abnormalities at days 3 or 4, or on weekly inspection thereafter. Physical examination revealed no fever, lymphadenopathy, conjunctivitis, tachypnea, or tachycardia at any of the time points. No abnormalities were found in a complete blood count or sedimentation rate, nor were abnormalities observed in serum electrolytes, transaminases, or blood urea nitrogen and creatinine.

Examination of Wright-stained cells from the nasal brushings showed that neutrophils and lymphocytes accounted for less than 5% of total cells in all three monkeys.

Administration of the Ad2/CFTR-1 caused no change in the distribution or number of inflammatory cells at any of the time points following virus administration. H&E stains of the nasal turbinate biopsies specimens from the control monkey could not be differentiated from that of the experimental monkey when the specimens were reviewed by an independent pathologist. (Fig. 24)

These results demonstrate the ability of a recombinant adenovirus encoding CFTR (Ad2/CFTR-1) to express CFTR cDNA in the airway epithelium of cotton rats and monkeys during repeated administration. They also indicate that application of the virus involves little if any risk. Thus, they suggest that such a vector may be of value in expressing CFTR in the airway epithelium of humans with cystic fibrosis.

Two methods were used to show that Ad2/CFTR-1 expresses CFTR in the airway epithelium of cotton rats and primates: CFTR mRNA was detected using RT-PCR and protein was detected by immunocytochemistry. Duration of expression as assessed immunocytochemically was five to six weeks. Because very little protein is required to generate Cl^- secretion (Welsh, M.J. (1987) *Physiol. Rev.* 67:1143-1184; Trapnell, B.C. et al. (1991) *Proc. Natl. Acad. Sci. USA* 88:6565-6569; Denning, G.M. et al. (1992) *J. Cell Biol.* 118:551-559), it is likely that functional expression of CFTR persists substantially longer than the period of time during which CFTR was detected by immunocytochemistry. Support for this evidence comes from two considerations: first, it is very difficult to detect CFTR immunocytochemically in the airway epithelium, yet the expression of an apical membrane Cl^- permeability due to the presence of CFTR Cl^- channels is readily detected. The ability of a minimal amount of CFTR to have important functional effects is likely a result of the fact that a single ion channel conducts a very large number of ions ($10^6 - 10^7$ ions/sec). Thus, ion channels are not usually abundant proteins in epithelia. Second, previous work suggests that the defective electrolyte transport of CF epithelia can be corrected when only 6-10% of cells in a CF airway epithelium overexpress wild-type CFTR (Johnson, L.G. et al. (1992) *Nature Gen.* 2:21-25). Thus, correction of the biologic defect in CF patients may be possible when only a small percent of the cells express CFTR. This is also consistent with our previous studies *in vitro* showing that Ad2/CFTR-1 at relatively low multiplicities of infection generated a cAMP-stimulated Cl^- secretory response in CF epithelia (Rich, D.P. et al. (1993) *Human Gene Therapy* 4:461-476).

This study also provides the first comprehensive data on the safety of adenovirus vectors for gene transfer to airway epithelium. Several aspects of the studies are encouraging. There was no evidence of viral replication, rather infectious viral particles were rapidly cleared from both cotton rats and primates. These data, together with our previous *in vitro* studies, suggest that replication of recombinant virus in humans will likely not be a problem. The other major consideration for safety of an adenovirus vector in the treatment of CF is the possibility of an inflammatory response. The data indicate that the virus generated an antibody response in both cotton rats and monkeys. Despite this, no evidence of a

systemic or local inflammatory response was observed. The cells obtained by bronchoalveolar lavage and by brushing and swabs were not altered by virus application. Moreover, the histology of epithelia treated with adenovirus was indistinguishable from that of control epithelia. These data suggest that at least three sequential exposures of airway epithelium to adenovirus does not cause a detrimental inflammatory response.

These data suggest that Ad2/CFTR-1 can effectively transfer CFTR cDNA to airway epithelium and direct the expression of CFTR. They also suggest that transfer is relatively safe in animals. Thus, they suggest that Ad2/CFTR-1 may be a good vector for treating patients with CF. This was confirmed in the following example.

10

Example 10 - CFTR Gene Therapy in Nasal Epithelia from Human CF Subjects

EXPERIMENTAL PROCEDURES

15 Adenovirus vector

The recombinant adenovirus Ad2/CFTR-1 was used to deliver CFTR cDNA. The construction and preparation of Ad2/CFTR-1, and its use *in vitro* and *in vivo* in animals, has been previously described (Rich, D.P. et al. (1993) *Human Gene Therapy* 4:461-476; Zabner, J. et al. (1993) *Nature Gen.* (in press)). The DNA construct comprises a full length copy of the Ad2 genome from which the early region 1 genes (nucleotides 546 to 3497) have been replaced by cDNA for CFTR. The viral E1a promoter was used for CFTR cDNA; this is a low to moderate strength promoter. Termination/polyadenylation occurs at the site normally used by E1b and protein IX transcripts. The E3 region of the virus was conserved.

25 Patients

Three patients with CF were studied. Genotype was determined by IG Labs (Framingham, MA). All three patients had mild CF as defined by an NIH score > 70 (Taussig, L.M. et al. (1973) *J. Pediatr.* 82:380-390), a normal weight for height ratio, a forced expiratory volume in one second (FEV1) greater than 50% of predicted and an arterial PO₂ greater than 72. All patients were seropositive for type 2 adenovirus, and had no recent viral illnesses. Pretreatment cultures of nasal swabs, pharyngeal swabs, sputum, urine, stool, and blood leukocytes were negative for adenovirus. PCR of pretreatment nasal brushings using primers for the adenovirus E1 region were negative. Patients were evaluated at least twice by FEV1, cytology of nasal mucosa, visual inspection, and measurement of V_t before treatment. Prior to treatment, a coronal computed tomographic scan of the paranasal sinuses and a chest X-ray were obtained.

The first patient was a 21 year old woman who was diagnosed at 3 months after birth. She had pancreatic insufficiency, a positive sweat chloride test (101 mEq/l), and is homozygous for the $\Delta F508$ mutation. Her NIH score was 90 and her FEV1 was 83%

predicted. The second patient was a 36 year old man who was diagnosed at the age of 13 when he presented with symptoms of pancreatic insufficiency. A sweat chloride test revealed a chloride concentration of 70 mEq/l. He is a heterozygote with the $\Delta F508$ and G551D mutations. His NIH score was 88 and his FEV1 was 66% predicted. The third patient was a 50 year old woman, diagnosed at the age of 9 with a positive sweat chloride test (104 mEq/l). She has pancreatic insufficiency and insulin dependent diabetes mellitus. She is homozygous for the $\Delta F508$ mutation. Her NIH score was 73 and her FEV1 was 65% predicted.

Transepithelial voltage

The transepithelial electric potential difference across the nasal epithelium was measured using techniques similar to those previously described (Alton, E.W.F.W. et al (1987) *Thorax* 42:815-817; Knowles, M. et al. (1981) *N. Eng. J. Med.* 305:1489-1495). A 23 gauge subcutaneous needle connected with sterile normal saline solution to a silver/silver chloride pellet (E.W. Wright, Guilford, CT) was used as a reference electrode. The exploring electrode was a size 8 rubber catheter (modified Argyle^R Foley catheter, St. Louis, MO) with one side hole at the tip. The catheter was filled with Ringer's solution containing (in mM), 135 NaCl, 2.4 KH₂PO₄, K₂HPO₄, 1.2CaCl₂, 1.2 MgCl₂ and 10 Hepes (titrated to pH 7.4 with NaOH) and was connected to a silver/silver chloride pellet. Voltage was measured with a voltmeter (Keithley Instruments Inc., Cleveland, OH) connected to a strip chart recorder (Servocorder, Watanabe Instruments, Japan). Prior to the measurements, the silver/silver chloride pellets were connected in series with the Ringer's solution; the pellets were changed if the recorded V_t was greater than ± 4 mV. The rubber catheter was introduced into the nostril under telescopic guidance (Hopkins Telescope, Karl Storz, Tuttlingen West Germany) and the side hole of the catheter was placed next to the study area in the medical aspect of the inferior nasal turbinate. The distance from the anterior tip of the inferior turbinate and the spatial relationship with the medial turbinate, the maxillary sinus ostium, and in one patient a small polyp, were used to locate the area of Ad2/CFTR-1 administration for measurements. Photographs and video recorder images were also used. Basal V_t was recorded until no changes in V_t were observed after slow intermittent 100 μ l/min infusion of the Ringer's solution. Once a stable baseline was achieved, 200 μ l of a Ringer's solution containing 100 μ M amiloride (Merck and Co. Inc., West Point, PA) was instilled through the catheter and changes in V_t were recorded until no further change were observed after intermittent instillations. Finally, 200 μ l Ringer's solution containing 100 μ M amiloride plus 10 μ M terbutaline (Geigy Pharmaceuticals, Ardsley, NY) was instilled and the changes in V_t were recorded.

Measurements of basal V_t were reproducible over time: in the three treated patients, the coefficients of variation before administration of Ad2/CFTR-1 were 3.6%, 12%, and 12%. The changes induced by terbutaline were also reproducible. In 30 measurements in 9 CF patients, the terbutaline-induced changes in V_t (ΔV_t) ranged from 0 mV to +4 mV;

hyperpolarization of V_t was never observed. In contrast, in 7 normal subjects ΔV_t ranged from -1 mV to -5 mV; hyperpolarization was always observed.

Ad2/CFTR-1 application and cell acquisition

5 The patients were taken to the operating room and monitoring was commenced using continuous EKG and pulse oximetry recording as well as automatic intermittent blood pressure measurement. After mild sedation, the nasal mucosa was anesthetized by atomizing 0.5 ml of 5% cocaine. The mucosa in the area of the inferior turbinate was then packed with cotton pledgets previously soaked in a mixture of 2 ml of 0.1% adrenaline and 8 ml of 1%
10 tetracaine. The pledgets remained in place for 10-40 min. Using endoscopic visualization with a television monitoring system, the applicator was introduced through the nostril and positioned on the medial aspect of the inferior turbinate, at least three centimeters from its anterior tip (Figures 25A-25I). The viral suspension was infused into the applicator through connecting catheters. The position of the applicator was monitored endoscopically to ensure
15 that it did not move and that enough pressure was applied to prevent leakage. After the virus was in contact with the nasal epithelium for thirty minutes, the viral suspension was removed, and the applicator was withdrawn. In the third patient's right nasal cavity, the virus was applied using the modified Foley catheter used for V_t measurements. The catheter was introduced without anesthetic under endoscopic guidance until the side hole of the catheter
20 was in contact with the area of interest in the inferior turbinate. The viral solution was infused slowly until a drop of solution was seen with the telescope. The catheter was left in place for thirty minutes and then removed.

Cells were obtained from the area of virus administration approximately 2 weeks before treatment and then at weekly intervals after treatment. The inferior turbinate was
25 packed for 10 minutes with cotton pledgets previously soaked in 1 ml of 5% cocaine. Under endoscopic control, the area of administration was gently brushed for 5 seconds. The brushed cells were dislodged in PBS. Swabs of the nasal epithelia were collected using cotton tipped applicators without anesthesia. Cytospin slides were prepared and stained with Wright's stain. Light microscopy was used to assess the respiratory epithelial cells and inflammatory
30 cells. For biopsies, sedatives/anesthesia was administered as described for the application procedure. After endoscopic inspection, and identification of the site to be biopsied, the submucosa was injected with 1% xylocaine, with 1/100,000 epinephrine. The area of virus application on the inferior turbinate was removed. The specimen was fixed in 4% formaldehyde and stained.

35

RESULTS

On day one after Ad2/CFTR-1 administration and at all subsequent time points, Ad2/CFTR-1 from the nasal epithelium, pharynx, blood, urine, or stool could not be cultured. As a control for the sensitivity of the culture assay, samples were routinely spiked with 10

and 100 IU Ad2/CFTR-1. In every case, the spiked samples were positive, indicating that, at a minimum, 10 IU of Ad2/CFTR should have been detected. No evidence of a systemic response as assessed by history, physical examination, serum chemistries or cell counts, chest and sinus X-rays, pulmonary function tests, or arterial blood gases performed before and after Ad2/CFTR-1 administration. An increase in antibodies to adenovirus was not detectable by ELISA or by neutralization for 35 days after treatment.

Three to four hours after Ad2/CFTR-1 administration, at the time that local anesthesia and localized vasoconstriction abated, all patients began to complain of nasal congestion and in one case, mild rhinorrhea. These were isolated symptoms that diminished by 18 hours and resolved by 28 to 42 hours. Inspection of the nasal mucosa showed mild to moderate erythema, edema, and exudate (Figures 25A-25C). These physical findings followed a time course similar to the symptoms. The physical findings were not limited to the site of virus application, even though preliminary studies using the applicator showed that marker methylene blue was limited to the area of application. In two additional patients with CF, the identical anesthesia and application procedure were used, but saline was applied instead of virus, yet the same symptoms and physical findings were observed in these patients (Figures 25G-25I). Moreover, the local anesthesia and vasoconstriction generated similar changes even when the applicator was not used, suggesting that the anesthesia/vasoconstriction caused some, if not all the injury. Twenty-four hours after the application procedure, analysis of cells removed from nasal swabs revealed an equivalent increase in the percent neutrophils in patients treated with Ad2/CFTR-1 or with saline. One week after application, the neutrophilia had resolved in both groups. Respiratory epithelial cells obtained by nasal brushing appeared normal at one week and at subsequent time points, and showed no evidence of inclusion bodies. To further evaluate the mucosa, the epithelium was biopsied on day three in the first patient and day one in the second patient. Independent evaluation by two pathologists not otherwise associated with the study suggested changes consistent with mild trauma and possible ischemia (probably secondary to the anesthetic/vasoconstrictors used before virus administration), but there were no abnormalities suggestive of virus-mediated damage.

Because the application procedure produced some mild injury in the first two patients, the method of administration was altered in the third patient. The method used did not require the use of local anesthesia or vasoconstriction and which was thus less likely to cause injury, but which was also less certain in its ability to constrain Ad2/CFTR-1 in a precisely defined area. On the right side, Ad2/CFTR-1 was administered as in the first two patients, and on the left side, the virus was administered without anesthesia or the applicator, instead using a small Foley catheter to apply and maintain Ad2/CFTR-1 in a relatively defined area by surface tension (Figure 25E). On the right side, the symptoms and physical findings were the same as those observed in the first two patients. By contrast, on the left side there were no symptoms and on inspection the nasal mucosa appeared normal (Figures 25D-25F). Nasal

swabs obtained from the right side showed neutrophilia similar to that observed in the first two patients. In contrast, the left side which had no anesthesia and minimal manipulation, did not develop neutrophilia. Biopsy of the left side on day 3 after administration (Figure 26), showed morphology consistent with CF-- a thickened basement membrane and
 5 occasional polymorphonuclear cells in the submucosa-- but no abnormalities that could be attributed to the adenovirus vector.

The first patient developed symptoms of a sore throat and increased cough that began three weeks after treatment and persisted for two days. Six weeks after treatment she developed an exacerbation of her bronchitis/bronchiectasis and hemoptysis that required
 10 hospitalization. The second patient had a transient episode of minimal hemoptysis three weeks after treatment; it was not accompanied by any other symptoms before or after the episode. The third patient has an exacerbation of bronchitis three weeks after treatment for which she was given oral antibiotics. Based on each patient's pretreatment clinical history, evaluation of the episodes, and viral cultures, no evidence could be discerned that linked
 15 these episodes to administration of Ad2/CFTR-1. Rather the episodes appeared consistent with the normal course of disease in each individual.

The loss of CFTR Cl^- channel function causes abnormal ion transport across affected epithelia, which in turn contributes to the pathogenesis of CF-associated airway disease (Boat, T.F. et al. in *The Metabolic Basis of Inherited Diseases* (Scriver, C.R. et al. eds.,
 20 McGraw-Hill, New York (1989); Quinton, P.M. (1990) *FASEB J.* 4:2709-2717). In airway epithelia, ion transport is dominated by two electrically conductive processes: amiloride-sensitive absorption of Na^+ from the mucosal to the submucosal surface and cAMP-stimulated Cl^- secretion in the opposite direction. (Quinton, P.M. (1990) *FASEB J.* 4:2709-2717; Welsh, M.J. (1987) *Physiol. Rev.* 67:1143-1184). These two transport processes can be
 25 assessed noninvasively by measuring the voltage across the nasal epithelium (V_t) *in vivo* (Knowles, M. et al (1981) *N. Eng. J. Med.* 305:1489-1495; Alton, E.W.F.W. et al.(1987) *Thorax* 42:815-817). Figure 27 shows an example from a normal subject. Under basal conditions, V_t was electrically negative (lumen referenced to the submucosal surface). Perfusion of amiloride (100 μM) onto the mucosal surface inhibited V_t by blocking apical
 30 Na^+ channels (Knowles, M. et al (1981) *N. Eng. J. Med.* 305:1489-1495; Quinton, P.M. (1990) *FASEB J.* 4:2709-2717; Welsh, M.J. (1992) *Neuron* 8:821-829). Subsequent perfusion of terbutaline (10 μM) a β -adrenergic agonist, hyperpolarized V_t by increasing cellular levels of cAMP, opening CFTR Cl^- channels, and stimulating chloride secretion (Quinton, P.M. (1990) *FASEB J.* 4:2709-2717; Welsh, M.J. et al. (1992) *Neuron* 8:821-829).
 35 Figure 28A shows results from seven normal subjects: basal V_t was $-10.5 \pm 1.0\text{mV}$, and in the presence of amiloride, terbutaline hyperpolarized V_t by $-2.3 \pm 0.5\text{mV}$.

In patients with CF, V_t was more electrically negative than in normal subjects (Figure 28B), as has been previously reported (Knowles, M. et al. (1981) *N. Eng. J. Med.* 305:1489-1495). Basal V_t was $-37.0 \pm 2.4\text{mV}$, much more negative than values in normal subjects ($P <$

0.001). (Note the difference in scale in Figure 28A and Figure 28B). Amiloride inhibited V_t , as it did in normal subjects. However, V_t failed to hyperpolarize when terbutaline was perfused onto the epithelium in the presence of amiloride. Instead, V_t either did not change or became less negative: on average V_t depolarized by $+1.8 \pm 0.6$ mV, a result very different from that observed in normal subjects. ($P < 0.001$).

After Ad2/CFTR-1 was applied, basal V_t became less negative in all three CF patients: Figure 29A shows an example from the third patient before (Figure 29A) and after (Figure 29B) treatment and Figures 30A, 30C, and 30E show the time course of changes in basal V_t for all three patients. The decrease in basal V_t suggests that application of Ad2/CFTR-1 corrected the CF electrolyte transport defect in nasal epithelium of all three patients. Additional evidence came from an examination of the response to terbutaline. Figure 30B shows that in contrast to the response before Ad2/CFTR-1 was applied, after virus replication, in the presence of amiloride, terbutaline stimulated V_t . Figures 30B, 30D, and 30F show the time course of the response. These data indicate that Ad2/CFTR-1 corrected the CF defect in Cl^- transport. Correction of the Cl^- transport defect cannot be attributed to the anesthesia/application procedure because it did not occur in patients treated with saline instead of Ad2/CFTR-1 (Figure 31). Moreover, the effects of the anesthesia were generalized on the nasal mucosa, but basal V_t decreased only in the area of virus administration. Finally, similar changes were observed in the left nasal mucosa of the third patient (Figures 30E and 30F), which had no symptomatic or physical response after the modified application procedure.

Unsuccessful attempts were made to detect CFTR transcripts by reverse transcriptase-PCR and by immunocytochemistry in cells from nasal brushings and biopsies. Although similar studies in animals have been successful (Zabner, J. et al. (1993) *Nature Gen.* (in press)), those studies used much higher doses of Ad2/CFTR-1. The lack of success in the present case likely reflects the small amount of available tissue, the low MOI, the fact that only a fraction of cells may have been corrected, and the fact that Ad2/CFTR-1 contains a low to moderate strength promoter (Ela) which produces much less mRNA and protein than comparable constructs using a much stronger CMV promoter (unpublished observation). The Ela promoter was chosen because CFTR normally expressed at very low levels in airway epithelial cells (Trapnell, B.C. et al. (1991) *Proc. Natl. Acad. Sci. USA* 88:6565-6569). It is also difficult to detect CFTR protein and mRNA in normal human airway epithelia, although function is readily detected because a single ion channel can conduct a very large number of ions per second and thus efficiently support Cl^- transport.

With time, the electrical changes that indicate correction of the CF defect reverted toward pretreatment values. However, the basal V_t appeared to revert more slowly than did the change in V_t produced by terbutaline. The significance of this difference is unknown, but it may reflect the relative sensitivity of the two measurements to expression of normal CFTR. In any case, this study was not designed to test the duration of correction because the treated

area was removed by biopsy on one side and the nasal mucosa on the other side was brushed to obtain cells for analysis at 7 to 10 days after virus administration, and then at approximately weekly intervals. Brushing the mucosa removes cells, disrupts the epithelium, and reduces basal V_t to zero for at least two days afterwards, thus preventing an accurate assessment of duration of the effect of Ad2/CFTR-1.

Efficacy of adenovirus-mediated gene transfer.

The major conclusion of this study is that *in vivo* application of a recombinant adenovirus encoding CFTR can correct the defect in airway epithelial Cl^- transport that is characteristic of CF epithelia.

Complementation of the Cl^- channel defect in human nasal epithelium could be measured as a change in basal voltage and as a change in the response to cAMP agonists. Although the protocol was not designed to establish duration, changes in these parameters were detected for at least three weeks. These results represent the first report that administration of a recombinant adenovirus to humans can correct a genetic lesion as measured by a functional assay. This study contrasts with most earlier attempts at gene transfer to humans, in that a recombinant viral vector was administered directly to humans, rather than using a *in vitro* protocol involving removal of cells from the patient, transduction of the cells in culture, followed by reintroduction of the cells into the patient.

Evidence that the CF Cl^- transport defect was corrected at all three doses of virus, corresponding to 1, 3, and 25 MOI, was obtained. This result is consistent with earlier studies showing that similar MOIs reversed the CF fluid and electrolyte transport defects in primary cultures of CF airway cells grown as epithelia on permeable filter supports (Rich, D.P. et al. (1993) *Human Gene Therapy* 4:461-476 and Zabner et al. submitted for publication): at an MOI of less than 1, cAMP-stimulated Cl^- secretion was partially restored, and after treatment with 1 MOI Ad2/CFTR-1 cAMP agonists stimulated fluid secretion that was within the range observed in epithelia from normal subjects. At an MOI of 1, a related adenovirus vector produced β -galactosidase activity in 20% of infected epithelial cells as assessed by fluorescence-activated cell analysis (Zabner et al. submitted for publication). Such data would imply that pharmacologic dose of adenovirus in CF airways might correspond to an MOI of one. If it is estimated that there are 2×10^6 cells/cm² in the airway (Mariassy, A.T. in *Comparative Biology of the Normal Lung* (CRC Press, Boca Raton 1992), and that the airways from the trachea to the respiratory bronchioles have a surface area of 1400 cm² (Weibel, E.R. *Morphometry of the Human Lung* (Springer Verlag, Heidelberg, 1963) then there would be approximately 3×10^9 potential target cells. Assuming a particle to IU ratio of 100, this would correspond to approximately 3×10^{11} particles of adenovirus with a mass of approximately 75 μ g. While obviously only a crude estimate, such information is useful in designing animal experiments to establish the likely safety profile of a human dose.

It is possible that an efficacious MOI of recombinant adenovirus could be less than the lowest MOI tested here. Some evidence suggests that not all cells in an epithelial monolayer need to express CFTR to correct the CF electrolyte transport defects. Mixing experiments showed that when perhaps 5-10% of cells overexpress CFTR, the monolayer exhibits wild-type electrical properties (Johnson, L.G. et al. (1992) *Nature Gen.* 2:21-25). Studies using liposomes to express CFTR in mice bearing a disrupted CFTR gene also suggest that only a small proportion of cells need to be corrected (Hyde, S.C. et al. (1993) *Nature* 362:250-255). The results referred to above using airway epithelial monolayers and multiplicities of Ad2/CFTR-1 as low as 0.1 showed measurable changes in Cl^- secretion (Rich, D.P. et al. (1993) *Human Gene Therapy* 4:461-476 and Zabner et al. submitted for publication).

Given the very high sensitivity of electrolyte transport assays (which result because a single Cl^- channel is capable of transporting large numbers of ions/sec) and the low activity of the E1a promoter used to transcribe CFTR, the inability to detect CFTR protein and CFTR mRNA are perhaps not surprising. Although CFTR mRNA could not be detected by reverse transcriptase-PCR, Ad2/CFTR-1 DNA could be detected in the samples by standard PCR, demonstrating the presence of input DNA and suggesting that the reverse transcriptase reaction may have been suboptimal. This could have occurred because of factors in the tissue that inhibit the reverse transcriptase. Although there is little doubt that the changes in electrolyte transport measured here result from expression of CFTR, it remains to be seen whether this will lead to measurable clinical changes in lung function.

Safety considerations.

Application of the adenovirus vector to the nasal epithelium in these three patients was well-tolerated. Although mild inflammation was observed in the nasal epithelium of all three patients following administration of Ad2/CFTR-1, similar changes were observed in two volunteers who underwent a sham procedure using saline rather than the viral vector. Clearly a combination of anesthetic- and procedure-related trauma resulted in the changes in the nasal mucosa. There is insufficient evidence to conclude that no inflammation results from virus administration. However, using a modified administration of the highest MOI of virus tested (25 MOI) in one patient, no inflammation was observed under conditions that resulted in evidence of biophysical efficacy that lasted until the area was removed by biopsy at three days.

There was no evidence of replication of Ad2/CFTR-1. Earlier studies had established that replication of Ad2/CFTR-1 in tissue culture and experimental animals is severely impaired (Rich, D.P. et al. (1993) *Human Gene Therapy* 4:461-476; Zabner, J. et al. (1993) *Nature Gen.* (in press)). Replication only occurs in cells that supply the missing early proteins of the E1 region of adenovirus, such as 293 cells, or under conditions where the E1 region is provided by coinfection with or recombination with an E1-containing adenovirus

(Graham, F.L. and Prevec, L. Vaccines: New Approaches to Immunological Problems (R.W. Ellis, ed., Boston, Butterworth-Heinemann, 1992); Berkner, K.L. (1988) *Biotechniques* 6:616-629). The patients studied here were seropositive for adenovirus types 2 and 5 prior to the study were negative for adenovirus upon culture of nasal swabs prior to administration of Ad2/CFTR-1, and were shown by PCR methods to lack endogenous E1 DNA sequences such as have been reported in some human subjects (Matsuse T. et al. (1992) *Am. Rev. Respir. Dis.* 146:177-184).

Example 11 - Construction and Packaging of Pseudo Adenoviral Vector (PAV)

10 With reference to Figure 32, the PAV construct was made by inserting the Ad2 packaging signal and E1 enhancer region (0-358 nt) in Bluescript II SK- (Stratagene, LaJolla, CA). A variation of this vector, known as PAV II was constructed similarly, except the Ad2 packaging signal and E1 enhancer region contained 0-380 nt. The addition of nucleotides at the 5' end results in larger PAVs, which may be more efficiently packaged, yet would include
15 more adenoviral sequences and therefore could potentially be more immunogenic or more capable of replicating.

To allow ease of manipulation for either the insertion of gene coding regions or complete excision and use in transfections for the purpose of generating infectious particles, a complementary plasmid was also built in pBluescript SKII-. This complementary plasmid
20 contains the Ad2 major late promoter (MLP) and tripartite leader (TPL) DNA and an SV40 T-antigen nuclear localization signal (NLS) and polyadenylation signal (SVpA). As can be seen in Figure 32, this plasmid contains a convenient restriction site for the insertion of genes of interest between the MLP/TPL and SV40 poly A. This construct is engineered such that the entire cassette may be excised and inserted into the former PAV I or PAV II construct.

25 Generation of PAV infectious particles was performed by excision of PAV from the plasmid with the Apa I and Sac II restriction endonucleases and co-transfection into 293 cells (an Ela/Elb expressing cell line) (Graham, F.L. et al, (1977) *J. Gen Virol* 36:59-74) with either wild-type Ad2, or packaging/replication deficient helper virus. Purification of PAV from helper can be accompanied by CsCl gradient isolation as PAV viral particles will be of a
30 lower density and will band at a higher position in the gradient.

For gene therapy, it is desirable to generate significant quantities of PAV virion free from contaminating helper virus. The primary advantage of PAV over standard adenoviral vectors is the ability to package large DNA inserts into virion (up to about 36 kb). However, PAV requires a helper virus for replication and packaging and this helper virus will be the
35 predominant species in any PAV preparation. To increase the proportion of PAV in viral preparation several approaches can be employed. For example, one can use a helper virus which is partially defective for packaging into virions (either by virtue of mutations in the packaging sequences (Grable, M. and Hearing P. (1992) *J. Virol.* 66: 723-731)) or by virtue of its size -viruses with genome sizes greater than approximately 37.5 kb package

inefficiently. In mixed infections with packaging defective virus, PAV would be expected to be represented at higher levels in the virus mixture than would occur with non-packaging defective helper viruses.

Another approach is to make the helper virus dependent upon PAV for its own replication. This may most easily be accomplished by deleting an essential gene from the helper virus (e.g. IX or a terminal protein) and placing that gene in the PAV vector. In this way neither PAV nor the helper virus is capable of independent replication - PAV and the helper virus are therefore co-dependent. This should result in higher PAV representation in the resulting virus preparation.

A third approach is to develop a novel packaging cell line, which is capable of generating significant quantities of PAV virion free from contaminating helper virus. A novel protein IX, (pIX) packaging system has been developed. This system exploits several documented features of adenovirus molecular biology. The first is that adenoviral defective particles are known to comprise up to 30% or more of standard wild-type adenoviral preparations. These defective or incomplete particles are stable and contain 15-95% of the adenoviral genome, typically 15-30%. Packaging of a PAV genome (15-30% of wild-type genome) should package comparably. Secondly, stable packaging of full-length Ad genome but not genomes <95% required the presence of the adenoviral gene designated pIX.

The novel packaging system is based on the generation of an Ad protein pIX expressing 293 cell line. In addition, an adenoviral helper virus engineered such that the E1 region is deleted but enough exogenous material is inserted to equal or slightly exceed the full length 36 kb size. Both of these two constructs would be introduced into the 293/pIX cell line as purified DNA. In the presence of pIX, yields of both predicted progeny viruses as seen in current PAV/Ad2 production experiments can be obtained. Virus containing lysates from these cells can then be titered independently (for the marker gene activity specific to either vector) and used to infect standard 293 (lacking pIX) at a multiplicity of infection of 1 relative to PAV. Since research with this line as well as from incomplete or defective particle research indicates that full length genomes have a competitive packaging advantage, it is expected that infection with an MOI of 1 relative to PAV will necessarily equate to an effective MOI for helper of greater than 1. All cells will presumably contain both PAV (at least 1) and helper (greater than 1). Replication and viral capsid production in this cell should occur normally but only PAV genomes should be packaged. Harvesting these 293/pIX cultures is expected to yield essentially helper-free PAV.

Example 12 - Construction of Ad2-E4/ORF 6

Ad2-E4/ORF6 (Figure 33 shows the plasmid construction of Ad2-E4/ORF6) which is an adenovirus 2 based vector deleted for all Ad2 sequences between nucleotides 32815 and 35577. This deletion removes all open reading frames of E4 but leaves the E4 promoter and first 32-37 nucleotides of the E4 mRNA intact. In place of the deleted sequences, a DNA

fragment encoding ORF6 (Ad2 nucleotides 34082-33178) which was derived by polymerase chain reaction of Ad2 DNA with ORF6 specific DNA primers (Genzyme oligo. # 2371 - CGGATCCTTTATTATAGGGGAAGTCCACGCCTAC (SEQ. ID NO:8) and oligo. #2372 - CGGGATCCATCGATGAAATATGACTACGTCCG (SEQ. ID NO:9) were inserted). Additional sequences supplied by the oligonucleotides included a cloning site at the 5' and 3' ends of the PCR fragment (Clal and BamHI respectively) and a polyadenylation sequence at the 3' end to ensure correct polyadenylation of the ORF6 mRNA. As illustrated in Figure 33, the PCR fragment was first ligated to a DNA fragment including the inverted terminal repeat (ITR) and E4 promoter region of Ad2 (Ad2 nucleotides 35937-35577) and cloned in the bacterial plasmid pBluescript (Stratagene) to create plasmid ORF6. After sequencing to verify the integrity of the ORF6 reading frame, the fragment encompassing the ITR and ORF6 was subcloned into a second plasmid, pAd Δ E4, which contains the 3' end of Ad2 from a Sac I site to the 3' ITR (Ad2 nucleotides 28562-35937) and is deleted for all E4 sequences (promoter to poly A site Ad2 positions 32815-35641) using flanking restriction sites. In this second plasmid, virus expressing only E4 ORF6, pAdORF6 was cut with restriction enzyme PacI and ligated to Ad2 DNA digested with PacI. This PacI site corresponds to Ad2 nucleotide 28612. 293 cells were transfected with the ligation and the resulting virus was subjected to restriction analysis to verify that the Ad2 E4 region had been substituted with the corresponding region of pAdORF6 and that the only remaining E4 open reading frame was ORF6.

A cell line could in theory be established that would fully complement E4 functions deleted from a recombinant virus. The problem with this approach is that E4 functions in the regulation of host cell protein synthesis and is therefore toxic to cells. The present recombinant adenoviruses are deleted for the E1 region and must be grown in 293 cells which complement E1 functions. The E4 promoter is activated by the Ela gene product, and therefore to prevent inadvertent toxic expression of E4 transcription of E4 must be tightly regulated. The requirements of such a promoter or transactivating system is that in the uninduced state expression must be low enough to avoid toxicity to the host cell, but in the induced state must be sufficiently activated to make enough E4 gene product to complement the E4 deleted virus during virus production.

Example 13

An adenoviral vector is prepared as described in Example 7 while substituting the phosphoglycerate kinase (PGK) promoter for the Ela promoter.

Example 14

An adenoviral vector is prepared as described in Example 11 while substituting the PGK promoter for the Ad2 major late promoter (MLP).

Example 15: Generation of Ad2-ORF6/PGK-CFTR

This protocol uses a second generation adenovirus vector named Ad2-ORF6/PGK-CFTR. This virus lacks E1 and in its place contains a modified transcription unit with the PGK promoter and a poly A addition site flanking the CFTR cDNA. The PGK promoter is of only moderate strength but is long lasting and not subject to shut off. The E4 region of the vector has also been modified in that the whole coding sequence has been removed and replaced by ORF6, the only E4 gene essential for growth of Ad in tissue culture. This has the effect of generating a genome of 101% the size of wild type Ad2.

The DNA construct comprises a full length copy of the Ad2 genome from which the early region 1 (E1) genes (present at the 5' end of the viral genome) have been deleted and replaced by an expression cassette encoding CFTR. The expression cassette includes the promoter for phosphoglycerate kinase (PGK) and a polyadenylation (poly A) addition signal from the bovine growth hormone gene (BGH). In addition, the E4 region of Ad2 has been deleted and replaced with only open reading frame 6 (ORF6) of the Ad2 E4 region. The adenovirus vector is referred to as AD2-ORF6/PGK-CFTR and is illustrated schematically in Figure 34. The entire wild-type Ad2 genome has been previously sequenced (Roberts, R.J., (1986) In Adenovirus DNA, W. Oberfler, editor, Martinus Nihoff Publishing, Boston) and the existing numbering system has been adopted here when referring to the wild type genome. Ad2 genomic regions flanking E1 and E4 deletions, and insertions into the genome are being completely sequenced.

The Ad2-ORF6/PGK-CFTR construct differs from the one used in our earlier protocol (Ad2/CFTR-1) in that the latter utilized the endogenous E1a promoter, had no poly A addition signal directly downstream of CFTR and retained an intact E4 region. The properties of Ad2/CFTR-1 in tissue culture and in animal studies have been reported (Rich et al., (1993) *Human Gene Therapy* 4:461-467; and Zabner et al. (1993) *Nature Genetics* (in Press)).

At the 5' end of the genome, nucleotides 357 to 3328 of Ad2 have been deleted and replaced with (in order 5' to 3') 22 nucleotides of linker, 534 nucleotides of the PGK promoter, 86 nucleotides of linker, nucleotides 123-4622 of the published CFTR sequence (Riordan et al. (1989) *Science* 245:1066-1073), 21 nucleotides of linker, and a 32 nucleotide synthetic BGH poly A addition signal followed by a final 11 nucleotides of linker. The topology of the 5' end of the recombinant molecule is illustrated in Figure 34.

At the 3' end of the genome of Ad2-ORF6/PGK-CFTR, Ad2 sequences between nucleotides 32815 and 35577 have been deleted to remove all open reading frames of E4 but retain the E4 promoter, the E4 cap sites and first 32-37 nucleotides of E4 mRNA. The deleted sequences were replaced with a fragment derived by PCR which contains open reading frame 6 of Ad2 (nucleotides 34082-33178) and a synthetic poly A addition signal. The topology of the 3' end of the molecule is shown in Figure 34. The sequence of this segment of the molecule will be confirmed. The remainder of the Ad2 viral DNA sequence is

published in Roberts, R.J. in Adenovirus DNA. (W. Oberfler, Martinus Nihoff Publishing, Boston, 1986). The overall size of the Ad2-ORF6/PGK-CFTR vector is 36,336 bp which is 101.3% of full length Ad2. See Table III for the sequence of Ad2-ORF6/PGK-CFTR.

The CFTR transcript is predicted to initiate at one of three closely spaced transcriptional start sites in the cloned PGK promoter (Singer-Sam et al. (1984) *Gene* 32:409-417) at nucleotides 828, 829 and 837 of the recombinant vector (Singer-Sam et al. (1984) *Gene* 32:409-417). A hybrid 5' untranslated region is comprised of 72, 80 or 81 nucleotides of PGK promoter region, 86 nucleotide of linker sequence, and 10 nucleotides derived from the CFTR insert. Transcriptional termination is expected to be directed by the BGH poly A addition signal at recombinant vector nucleotide 5530 yielding an approximately 4.7 kb transcript. The CFTR coding region comprises nucleotides 1010-5454 of the recombinant virus and nucleotides 182, 181 or 173 to 4624, 4623, or 4615 of the PGK-CFTR-BGH mRNA respectively, depending on which transcriptional initiation site is used. Within the CFTR cDNA there are two differences from the published (Riordan et al, *cited supra*) cDNA sequence. An A to C change at position 1990 of the CFTR cDNA (published CFTR cDNA coordinates) which was an error in the original published sequence, and a T to C change introduced at position 936. The change at position 936 is translationally silent but increases the stability of the cDNA when propagated in bacterial plasmids (Gregory et al. (1990) *Nature* 347:382-386; and Cheng et al. (1990) *Cell* 63:827-834). The 3' untranslated region of the predicted CFTR transcript comprises 21 nucleotides of linker sequence and approximately 10 nucleotides of synthetic BGH poly A additional signal.

Although the activity of CFTR can be measured by electrophysiological methods, it is relatively difficult to detect biochemically or immunocytochemically, particularly at low levels of expression (Gregory et al., *cited supra*; and Denning et al. (1992) *J. Cell Biol.* 118:551-559). A high expression level reporter gene encoding the *E. coli* β galactosidase protein fused to a nuclear localization signal derived from the SV40 T-antigen was therefore constructed. Reporter gene transcription is driven by the powerful CMV early gene constitutive promoter. Specifically, the E1 region of wild type Ad2 between nucleotides 357-3498 has been deleted and replaced it with a 515 bp fragment containing the CMV promoter and a 3252 bp fragment encoding the β galactosidase gene.

Regulatory Characteristics of the Elements of the AD2-ORF6/PGK-CFTR

In general terms, the vector is similar to several earlier adenovirus vectors encoding CFTR but it differs in three specific ways from the Ad2/CFTR-1 construct.

PGK Promoter

Transcription of CFTR is from the PGK promoter. This is a promoter of only moderate strength but because it is a so-called house keeping promoter we considered it more likely to be capable of long term albeit perhaps low level expression. It may also be less

likely to be subject to "shut-down" than some of the very strong promoters used in other studies especially with retroviruses. Since CFTR is not an abundant protein longevity of expression is probably more critical than high level expression. Expression from the PGK promoter in a retrovirus vector has been shown to be long lasting (Apperley et al. (1991)

5 *Blood* 78:310-317).

Polyadenylation Signal

Ad2-ORG6/PGK-CFTR contains an exogenous poly A addition signal after the CFTR coding region and prior to the protein IX coding sequence of the Ad2 E1 region. Since
10 protein is believed to be involved in packaging of virions, this coding region was retained. Furthermore, since protein IX is synthesized from a separate transcript with its own promoter, to prevent possible promoter occlusion at the protein IX promoter, the BGH poly A addition signal was inserted. There is indirect evidence that promoter occlusion can be problematic in that Ad2/CMV β Gal grows to lower viral titers on 293 cells than does Ad2/ β gal-1. These
15 constructs are identical except for the promoter used for β galactosidase expression. Since the CMV promoter is much stronger than the E1a promoter it is probable that abundant transcription from the CMV promoter through the β galactosidase DNA into the protein IX coding region reduces expression of protein IX from its own promoter by promoter occlusion and that this is responsible for the lower titer of Ad2/CMV- β gal obtained.

20

Alterations of the E4 Region

A large portion of the E4 region of the Ad2 genome has been deleted for two reasons. The first reason is to decrease the size of the vector used or expression of CFTR. Adenovirus vectors with genomes much larger than wild type are packaged less efficiently and are
25 therefore difficult to grow to high titer. The combination of the deletions in the E1 and E4 regions in Ad2-ORG6/PGK-CFTR reduce the genome size to 101% of wild type. In practice it is straightforward to prepare high titer lots of this virus.

The second reason to remove E4 sequences relates to the safety of adenovirus vectors. A goal of these studies is to remove as many viral genes as possible to inactivate the Ad2 virus backbone in as many ways as possible. The OF 6/7 gene of the E4 region encodes a protein
30 that is involved in activation of the cellular transcription factor E2-F which is in turn implicated in the activation of the E2 region of adenovirus (Hemstrom et al. (1991) *J. Virol.* 65:1440-1449). Therefore removal of ORF6/7 from adenovirus vectors may provide a further margin of safety at least when grown in non-proliferating cells. The removal of the E1 region already renders such vectors disabled, in part because E1a, if present, is able to displace E2-F from the retinoblastoma gene product, thereby also contributing to the stimulation of E2
35 transcription. The ORF6 reading frame of Ad2 was added back to the E1-E4 backbone of the Ad2-ORG6/PGK-CFTR vector because ORF6 function is essential for production of the recombinant virus in 293 cells. ORF6 is believed to be involved in DNA replication, host

cell shut off and late mRNA accumulation in the normal adenovirus life cycle. The E1-E4-ORF6⁺ backbone Ad2 vector does replicate in 293 cells.

The promoter/enhancer used to drive transcription of ORF6 of E4 is the endogenous E4 promoter. This promoter requires E1a for activation and contains E1a core enhancer elements and SP1 transcription factor binding sites (reviewed in Berk, A.J. (1986) *Ann. Rev. Genet.* 20:75-79).

Replication Origin

The only replication origins present in Ad2-ORF6/PGK-CFTR are those present in the Ad2 parent genome. Replication of Ad2-ORF6/PGK-CFTR sequences has not been detected except when complemented with wild type E1 activity.

Steps Used to Derive the DNA Construct

Construction of the recombinant Ad2-ORF6/PGK-CFTR virus was accomplished by *in vivo* recombination of Ad2-ORF6 DNA and a plasmid containing the 5' 10.7 kb of adenovirus engineered to have an expression cassette encoding the human CFTR cDNA driven by the PGK promoter and a BGH poly A signal in place of the E1 coding region.

The generation of the plasmid, pBRAd2/PGK-CFTR is described here. The starting plasmid contains an approximately 7.5 kb insert cloned into the ClaI and BamHI sites of pBR322 and comprises the first 10,680 nucleotides of Ad2 with a deletion of the Ad2 sequences between nucleotides 356 and 3328. This plasmid contains a CMV promoter inserted into the ClaI and SpeI sites at the region of the E1 deletion and is designated pBRAd2/CMV. The plasmid also contains the Ad2 5' ITR, packaging and replication sequences and E1 enhancer. The E1 promoter, E1a and most of E1b coding region has been deleted. The 3' terminal portion of the E1b coding region coincides with the pIX promoter which was retained. The CMV promoter was removed and replaced with the PGK promoter as a ClaI and SpeI fragment from the plasmid PGK-GCR. The resulting plasmid, pBRAd2/PGK, was digested with AvrII and BstBI and the excised fragment replaced with the SpeI to BstBI fragment from the plasmid construct pAd2E1a/CFTR. This transferred a fragment containing the CFTR cDNA, BGH poly A signal and the Ad2 genomic sequences from 3327 to 10,670. The resulting plasmid is designated pBRAd2/PGK-CFTR. The CFTR cDNA fragment was originally derived from the plasmid pCMV-CFTR-936C using restriction enzymes SpeI and Ecl136II. pCMV-CFTR-936C consists of a minimal CFTR cDNA encompassing nucleotides 123-4622 of the published CFTR sequence cloned into the multiple cloning site of pRC/CMV (Invitrogen Corp.) using synthetic linkers. The CFTR cDNA within this plasmid has been completely sequenced.

The Ad2 backbone virus with the E4 region that expresses only open reading frame 6 was constructed as follows. A DNA fragment encoding ORF6 (Ad2 nucleotides 34082-33178) was derived by PCR with ORF6 specific DNA primers. Additional sequences

supplied by the oligonucleotides include cloning sites at the 5' and 3' ends of the PCR fragment. (ClaI and BamHI respectively) and a poly A addition sequence AATAAA at the 3' end to ensure correct polyadenylation of ORF6 mRNA. The PCR fragment was cloned into pBluescript (Stratagene) along with an Ad2 fragment (nucleotides 35937-35577) containing the inverted terminal repeat, E4 promoter, E4 mRNA cap sites and first 32-37 nucleotides of E4 mRNA to create pORF6. A Sall-BamHI fragment encompassing the ITR and ORF6 was used to replace the Sall-BamHI fragment encompassing the ITR and E4 deletion in pAdΔE4 contains the 3' end of Ad2 from a SpeI site to the 3' ITR (nucleotides 27123-35937) and is deleted for all E4 sequences including the promoter and poly A signal (nucleotides 32815-35641). The resulting construct, pAdE4ORF6 was cut with PacI and ligated to Ad2 DNA digested with PacI nucleotide 28612). 293 cells were transfected with the ligation reaction to generate virus containing only open reading frame 6 from the E4 region.

In Vitro Studies with Ad2-ORF6/PGK-CFTR

The ability of Ad2-ORF6/PGK-CFTR to express CFTR in several cell lines, including human HeLa cells, human 293 cells, and primary cultures of normal and CF human airway epithelia was tested. As an example, the results from the human 293 cells is related here. When human 293 cells were grown on culture dishes, the vector was able to transfer CFTR cDNA and express CFTR as assessed by immunoprecipitation and by functional assays of halide efflux. Gregory, R.J. et al. (1990) *Nature* 347:382-386; Cheng, S.H. et al. (1990) *Cell* 63:827-834. More specifically, procedures for preparing cell lysates, immunoprecipitation of proteins using anti-CFTR antibodies, one-dimensional peptide analysis and SDS-polyacrylamide gel electrophoresis were as described by Cheng et al. Cheng, S.H. et al. (1990) *Cell* 63:827-834. Halide efflux assays were performed as described by Cheng, S.H. et al. (1991) *Cell* 66:1027-1036. cAMP-stimulated CFTR chloride channel activity was measured using the halide sensitive fluorophore SPQ in 293 cells treated with 500 IU/cell Ad2-ORF6/PGK-CFTR. Stimulation of the infected cells with forskolin (20 μM) and IBMX (100 μM) increased SPQ fluorescence indicating the presence of functional chloride channels produced by the vector.

Additional studies using primary cultures of human airway (nasal polyp) epithelial cells (from CF patients) infected with Ad2-ORF6/PGK-CFTR demonstrated that Ad2-ORF6/PGK-CFTR infection of the nasal polyp epithelial cells resulted in the expression of cAMP dependent Cl⁻ channels. Figure 35 is an example of the results obtained from such studies. Primary cultures of CF nasal polyp epithelial cells were infected with Ad2-ORF6/PGK-CFTR at multiplicities of 0.3, 3, and 50. Three days post infection, monolayers were mounted in Ussing chambers and short-circuit current was measured. At the indicated times: (1) 10 μM amiloride, (2) cAMP agonists (10 μM forskolin and 100 μM IBMX), and (3) 1 mM diphenylamine-2-carboxylate were added to the mucosal solution.

In Vivo Studies with Ad2-ORF6/PGK-CFTR

Virus preparation

Two preparations of Ad2-ORF6/PGK-CFTR virus were used in this study. Both were prepared at Genzyme Corporation, in a Research Laboratory. The preparations were purified on a CsCl gradient and then dialyzed against tris-buffered saline to remove the CsCl. The preparation for the first administration (lot #2) had a titer of 2×10^{10} IU/ml. The preparation for the second administration (lot #6) had a titer of 4×10^{10} IU/ml.

Animals

Three female Rhesus monkeys, *Macaca mulatta*, were used for this study. Monkey C (#20046) weighed 6.4 kg. Monkey D (#20047) weighed 6.25 kg. Monkey E (#20048) weighed 10 kg. The monkeys were housed in the University of Iowa at least 360 days before the start of the study. The animals were maintained with free access to food and water throughout the study. The animals were part of a safety study and efficacy study for a different viral vector (Ad2/CFTR-1) and they were exposed to 3 nasal viral instillation throughout the year. The previous instillation of Ad2/CFTR-1 was performed 116 days prior to the initiation of this study. All three Rhesus monkeys had an anti-adenoviral antibody response as detected by ELISA after each viral instillation. There are no known contaminants that are expected to interfere with the outcome of this study. Fluorescent lighting was controlled to automatically provide alternate light/dark cycles of approximately 12 hours each. The monkeys were housed in an isolation room in separate cages. Strict respiratory and body fluid isolation precautions were taken.

Virus administration

For application of the virus, the monkeys were anesthetized by intramuscular injection of ketamine (15 mg/kg). The entire epithelium of one nasal cavity in each monkey was used for this study. A foley catheter (size 10) was inserted through each nasal cavity into the pharynx, the balloon was inflated with a 2-3 ml of air, and then pulled anteriorly to obtain a tight occlusion at the posterior choana. The Ad2-ORF6/PGK-CFTR virus was then instilled slowly into the right nostril with the posterior balloon inflated. The viral solution remained in contact with the nasal mucosa for 30 min. The balloons were deflated, the catheters were removed, and the monkeys were allowed to recover from anesthesia.

On the first administration, the viral preparation had a titer of 2×10^{10} IU/ml and each monkey received approximately 0.3 ml. Thus the total dose applied to each monkey was approximately 6.5×10^9 IU. This total dose is approximately half the highest dose proposed for the human study. When considered on a IU/kg basis, a 6 kg monkey received a dose approximately 3 times greater than the highest proposed dose for a 60 kg human.

Timing of evaluations.

The animals were evaluated on the day of administration, and on days 3, 7, 24, 38, and 44 days after infection. The second administration of virus occurred on day 44. The monkeys were evaluated on day 48 and then on days 55, 62, and 129.

- 5 For evaluations, monkeys were anesthetized by intramuscular injection of ketamine (15 mg/kg). To obtain nasal epithelial cells after the first viral administration, the nasal mucosa was first impregnated with 5 drops of Afrin (0.05% oxymetazoline hydrochloride, Schering-Plough) and 1 ml of 2% Lidocaine for 5 minutes. A cytobrush was then used to gently rub the mucosa for about 3 sec. To obtain pharyngeal epithelial swabs, a cotton-tipped
- 10 applicator was rubbed over the back of the pharynx 2-3 times. The resulting cells were dislodged from brushes or applicators into 2 ml of sterile PBS. After the second administration of Ad2-ORF6/PGK-CFTR, the monkeys were followed clinically for 3 weeks, and mucosal biopsies were obtained from the monkeys medial turbinate at days 4, 11 and 18.

15 Animal evaluation.

- Animals were evaluated daily for evidence of abnormal behavior or physical signs. A record of food and fluid intake was used to assess appetite and general health. Stool consistency was also recorded to check for the possibility of diarrhea. At each of the evaluation time points, rectal temperature, respiratory rate, and heart rate were measured.
- 20 The nasal mucosa, conjunctivas and pharynx were visually inspected. The monkeys were also examined for lymphadenopathy.

Hematology and serum chemistry

- Venous blood from the monkeys was collected by standard venipuncture technique.
- 25 Blood/serum analysis was performed in the clinical laboratory of the University of Iowa Hospitals and Clinics using a Hitachi 737 automated chemistry analyzer and a Technicom H6 automated hematology analyzer.

Serology

- 30 Sera from the monkeys were obtained and anti-adenoviral antibody titers were measured by ELISA. For the ELISA, 50 ng/well of killed adenovirus (Lee Biomolecular Research Laboratories, San Diego, Ca) was coated in 0.1M NaHCO₃ at 4° C overnight on 96 well plates. The test samples at appropriate dilutions were added, starting at a dilution of 1/50. The samples were incubated for 1 hour, the plates washed, and a goat anti-human IgG
- 35 HRP conjugate (Jackson ImmunoResearch Laboratories, West Grove, PA) was added for 1 hour. The plates were washed and O-Phenylenediamine (OPD) (Sigma Chemical Co., St. Louis, MO) was added for 30 min. at room temperature. The assay was stopped with 4.5 M H₂SO₄ and read at 490 nm on a Molecular Devices microplate reader. The titer was calculated as the product of the reciprocal of the initial dilution and the reciprocal of the

dilution in the last well with an OD>0.100. Nasal washings from the monkeys were obtained and anti-adenoviral antibody titers were measured by ELISA, starting at a dilution of 1/4.

Nasal Washings.

5 Nasal washings were obtained to test for the possibility of secretory antibodies that could act as neutralizing antibodies. Three ml of sterile PBS was slowly instilled into the nasal cavity of the monkeys, the fluid was collected by gravity. The washings were centrifuged at 1000 RPM for 5 minutes and the supernatant was used for anti-adenoviral, and neutralizing antibody measurement.

10

Cytology

Cells were obtained from the monkey's nasal epithelium by gently rubbing the nasal mucosa for about 3 seconds with a cytobrush. The resulting cells were dislodged from the brushes into 2 ml of PBS. The cell suspension was spun at 5000 rpm for 5 min. and
15 resuspended in 293 media at a concentration of 10^6 cells/ml. Forty μ l of the cell suspension was placed on slides using a Cytospin. Cytospin slides were stained with Wright's stain and analyzed for cell differential using light microscopy.

Culture for Ad2-ORF6/PFK-CFTR

20 To assess for the presence of infectious viral particles, the supernatant from the nasal brushings and pharyngeal swabs of the monkeys were used. Twenty-five μ l of the supernatant was added in duplicate to 293 cells. 293 cells were used at 50% confluence and were seeded in 96 well plates. 293 cells were incubated for 72 hours at 37°C, then fixed with a mixture of equal parts of methanol and acetone for 10 min and incubated with an FITC
25 label anti-adenovirus monoclonal antibodies (Chemicon, Light Diagnostics, Temecuca, Ca) for 30 min. Positive nuclear immunofluorescence was interpreted as positive culture.

Immunocytochemistry for the detection of CFTR.

Cells were obtained by brushing. Eighty μ l of cell suspension were spun onto gelatin-coated slides. The slides were allowed to air dry, and then fixed with 4% paraformaldehyde.
30 The cells were permeabilized with 0.2 Triton-X (Pierce, Rockford, Il) and then blocked for 60 minutes with 5% goat serum (Sigma, Mo). A pool of monoclonal antibodies (M13-1, M1-4, and M6-4) (Gregory et al., (1990) *Nature* 347:382-386); Denning et al., (1992) *J. Cell Biol.* 118:(3) 551-559; Denning et al., (1992) *Nature* 358:761-764) were added and incubated for
35 12 hours. The primary antibody was washed off and an antimouse biotinylated antibody (Biomeda, Foster City, Ca) was added. After washing, the secondary antibody, streptavidin FITC (Biomeda, Foster City, Ca) was added and the slides were observed with a laser scanning confocal microscope.

Biopsies

To assess for histologic evidence of safety, nasal medial turbinate biopsies were obtained on day 4, 11 and 18 after the second viral administration as described before (Zabner et al (1993) Human Gene Therapy, in press). Nasal biopsies were fixed in 4% formaldehyde and H&E stained sections were reviewed.

RESULTS

Studies of efficacy.

To directly assess the presence of CFTR, cells obtained by brushing were plated onto slides by cytospin and stained with antibodies to CFTR. A positive reaction is clearly evident in cells exposed to Ad2-ORF6/PGK-CFTR. The cells were scored as positive by immunocytochemistry when evaluated by a reader blinded to the identity of the samples. Cells obtained prior to infection and from other untreated monkeys were used as negative controls. Figures 36A-36D, 37A-37D, and 38A-38D show examples from each monkey.

Studies of safety

None of the monkeys developed any clinical signs of viral infections or inflammation. There were no visible abnormalities at days 3, 4, 7 or on weekly inspection thereafter. Physical examination revealed no fever, lymphadenopathy, conjunctivitis, coryza, tachypnea, or tachycardia at any of the time points. There was no cough, sneezing or diarrhea. The monkeys had no fever. Appetites and weights were not affected by virus administration in either monkey. The data are summarized in Figures 39A-39C.

The presence of live virus was tested in the supernatant of cell suspensions from swabs and brushes from each nostril and the pharynx. Each supernatant was used to infect the virus-sensitive 293 cell line. Live virus was never detected at any of the time points. The rapid loss of live virus suggests that there was no viral replication.

The results of complete blood counts, sedimentation rate, and clinical chemistries are shown in Figure 40A-40C. There was no evidence of a systemic inflammatory response or other abnormalities of the clinical chemistries.

Epithelial inflammation was assessed by cytological examination of Wright-stained cells (cytospin) obtained from brushings of the nasal epithelium. The percentage of neutrophils and lymphocytes from the infected nostrils were compared to those of the control nostrils and values from four control monkeys. Wright stains of cells from nasal brushing were performed on each of the evaluation days. Neutrophils and lymphocytes accounted for less than 5% of total cells at all time points. The data are shown in Figure 41. The data indicate that administration of Ad2-ORF6/PGK-CFTR caused no change in the distribution or number of inflammatory cells at any of the time points following virus administration,

even during a second administration of the virus. The biopsy slides obtained after the second Ad2-ORF6/PGK-CFTR administration were reviewed by an independent pathologist, who found no evidence of inflammation or any other cytopathic effects. Figures 42 to 44 show an example from each monkey.

5 Figures 45A-45C shows that all three monkeys had developed antibody titers to adenovirus prior to the first infection with Ad2-ORF6/PGK-CFTR (Zabner et al. (1993) *Human Gene Therapy* (in press)). Antibody titers measured by ELISA rose within one week after the first and second administration and peaked at day 24. No anti-adenoviral antibodies were detected by ELISA or neutralizing assay in nasal washings of any of the monkeys.

10 These results combined with demonstrate the ability of a recombinant adenovirus encoding CFTR (Ad2-ORF6/PGK-CFTR) to express CFTR cDNA in the airway epithelium of monkeys. These monkeys have been followed clinically for 12 months after the first viral administration and no complications have been observed.

15 The results of the safety studies are encouraging. No evidence of viral replication was found; infectious viral particles were rapidly cleared. The other major consideration for safety of an adenovirus vector in the treatment of CF is the possibility of an inflammatory response. The data indicate that the virus generated an antibody response, but despite this, no evidence of a systemic or local inflammatory response was observed. The cells obtained by brushings and swabs were not altered by virus application. Since these Monkeys had been
20 previously exposed three times to Ad2/CFTR-1, these data suggest that at least five sequential exposures of airway epithelium to adenovirus does not cause a detrimental inflammatory response.

25 These data indicate that Ad2-ORF6/PGK-CFTR can effectively transfer CFTR cDNA to airway epithelium and direct the expression of CFTR. They also indicate that transfer and expression is safe in primates.

Equivalents

30 Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents of the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

TABLE I

Mutant	CF	Exon	CFTR Domain	A	B
Wild Type				-	+
R334W	Y	7	TM6	-	+
K464M	N	9	NBD1	-	+
Δ 1507	Y	10	NBD1	-	+
Δ F508	Y	10	NBD1	-	+
F508R	N	10	NBD1	-	+
S549I	Y	11	NBD1	-	+
G551D	Y	11	NBD1	-	+
N894,900Q	N	15	ECD4	+	-
K1250M	N	20	NBD2	-	+
Tth111	N	22	NB-Term	-	+

Table II.

10	20	30	40	50	60
CATCATCAAT AATATACCTT ATTTTGGATT GAAGCCAATA TGATAATGAG GGGGTGGAGT					
GTAGTAGTTA TTATATGGAA TAAACCTAA CTTCGGTTAT ACTATTACTC CCCCACCTCA					
____INVERTED TERMINAL REPETITION-ORIGIN OF REPLICATION____					60>
70	80	90	100	110	120
TTGTGACGTG GCGCGGGGCG TGGGAACGGG GCGGGTGACG TAGTAGTGTG GCGGAAGTGT					
AACACTGCAC CGCGCCCGCG ACCCTTGCCC CGCCCACTGC ATCATCACAC CGCCTTCACA					
____INVERTED TERMINAL REPETITION-ORIGIN OF R____>					
130	140	150	160	170	180
GATGTTGCAA GTGTGGCGGA ACACATGTAA GCGCCGGATG TGGTAAAAGT GACGTTTTTG					
CTACAACGTT CACACCGCCT TGTGTACATT CGCGGCCTAC ACCATTTTCA CTGCAAAAAC					
190	200	210	220	230	240
GTGTGCGCCG GTGTATACGG GAAGTGACAA TTTTCGCGCG GTTTTAGGCG GATGTTGTAG					
CACACGCGGC CACATATGCC CTTCACGTG AAAAGCGCGC CAAAATCCGC CTACAACATC					
____b____ELA ENHANCER AND VIRAL PACKAGING DOMAIN____					50>
250	260	270	280	290	300
TAAATTTGGG CGTAACCAAG TAATGTTTGG CCATTTTCGC GGGAAACTG AATAAGAGGA					
ATTTAAACCC GCATTGGTTC ATTACAAACC GGTAAAAGCG CCCTTTGGAC TTATTCTCCT					
____60_b____ELA ENHANCER AND VIRAL PACKAGING DOMAIN_0_b____					110>
310	320	330	340	350	360
AGTGAAATCT GAATAATTCT GTGTACTCA TAGCGCGTAA TATTTGTCTA GGGCCGCGGG					
TCACTTTAGA CTTATTAAGA CACAATGAGT ATCGCGCATT ATAAACAGAT CCCGGCGCCC					
____120_b____ELA ENHANCER AND VIRAL PACKAGING DOMAIN_0_b____					170>
370	380	390	400	410	420
GACTTTGACC GTTTACGTGG AGACTCGCCC AGGTGTTTTT CTCAGGTGTT TTCCGCGTTC					
CTGAAACTGG CAAATGCACC TCTGAGCGGG TCCACAAAAA GAGTCCACAA AAGGCGCAAG					
____ELA ENHANCER A_90>					
____c____10_EL A PROMOTER REGION_0_c____					40>
430	440	450	460	470	480
CGGGTCAAAG TTGGCGTTTT ATTATTATAG TCAGCTGACG CGCAGTGTAT TTATACCCGG					
GCCCACTTTC AACCGCAAAA TAATAATATC AGTCGACTGC GCGTCACATA AATATGGGCC					
____50_c____60_EL A PROMOTER REGION_c____					90_c____100>
490	500	510	520	530	540
TGAGTTCCTC AAGAGGCCAC TCTTGAGTGC CAGCGAGTAG AGTTTTCTCC TCCGAGCGGC					
ACTCAAGGAG TTCTCCGGTG AGAAGTCACG GTCGCTCATC TCAAAAGAGG AGGCTCGGCG					
____h____HYBRID ELA-CFTR-ELB MESSAGE____>					
____ELA PROMOTER_120>					
____o____EL A RNA 5' UNTRANSLATED_o____					40>
550	560	570	580	590	600
TCCGAGCTAG TAACGGCCGC CAGTGTCTTG CAGATATCAA AGTCGACGGT ACCCGAGAGG					
AGGCTCGATC ATTGCGCGCG GTCACACGAC GTCTATAGTT TCAGGTGCCA TGGGCTCTCT					

_____h_____HYBRID ELA-CFTR-ELB MESSAGE _____h_____
 _____>
 _____e_____10_____SYNTHETIC LINKER SEQUENCES_____40_____e_____
 _____130>

610 620 630 640 650 660
 CCATGCAGAG GTCGCCTCTG GAAAAGGCCA GCGTTGTCTC CAAACTTTTT TTCAGCTGGA
 GGTACGTCTC CAGCGGAGAC CTTTCCGGT CGCAACAGAG GTTTGAAAAA AAGTCGACCT
 M Q R S P L E K A S V V S K L F F S W>
 _____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON_____
 _____h_____HYBRID ELA-CFTR-ELB MESSAGE _____h_____
 _____140i_____123 TO 4622 OF HUMAN CFTR CDNA_____180i_____190>

670 680 690 700 710 720
 CCAGACCAAT TTTGAGGAAA GGATACAGAC AGCGCCTGGA ATTGTCAGAC ATATACCAAA
 GGTCTGGTTA AAACCTCTTT CCTATGTCTG TCGCGGACCT TAACAGTCTG TATATGGTTT
 T R P I L R K G Y R Q R L E L S D I Y Q>
 _____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON_____
 _____h_____HYBRID ELA-CFTR-ELB MESSAGE _____h_____
 _____200i_____123 TO 4622 OF HUMAN CFTR CDNA_____240i_____250>

730 740 750 760 770 780
 TCCCTTCTGT TGATTCTGCT GACAATCTAT CTGAAAATT GGAAAGAGAA TGGGATAGAG
 AGGGAAGACA ACTAAGACGA CTGTTAGATA GACTTTTTAA CCTTCTCTT ACCCTATCTC
 I P S V D S A D N L S E K L E R E W D R>
 _____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON_____
 _____h_____HYBRID ELA-CFTR-ELB MESSAGE _____h_____
 _____260i_____123 TO 4622 OF HUMAN CFTR CDNA_____300i_____310>

790 800 810 820 830 840
 AGCTGGCTTC AAAGAAAAT CCTAACTCA TTAATGCCCT TCGGCGATGT TTTTCTGGA
 TCGACCGAAG TTTCTTTTAA GGATTGAGT ATTACGGGA AGCCGCTACA AAAAGACCT
 E L A S K K N P K L I N A L R R C F F W>
 _____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON_____
 _____h_____HYBRID ELA-CFTR-ELB MESSAGE _____h_____
 _____320i_____123 TO 4622 OF HUMAN CFTR CDNA_____360i_____370>

850 860 870 880 890 900
 GATTATGTT CTATGGATC TTTTATATT TAGGGGAAGT CACCAAGCA GTACAGCCTC
 CTAATACAA GATACCTTAG AAAATATAA ATCCCCTTCA GTGGTTTCGT CATGTCGGAG
 R F M F Y G I F L Y L G E V T K A V Q P>
 _____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON_____
 _____h_____HYBRID ELA-CFTR-ELB MESSAGE _____h_____
 _____380i_____123 TO 4622 OF HUMAN CFTR CDNA_____420i_____430>

910 920 930 940 950 960
 TCTTACTGGG AAGATCATA GCTTCCTATG ACCCGGATAA CAAGGAGGAA CGCTCTATCG
 AAGATGACCC TTCTTAGTAT CGAAGGATAC TGGGCCTATT GTTCCTCCTT GCGAGATAGC
 L L L G R I I A S Y D P D N K E E R S I>
 _____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON_____
 _____h_____HYBRID ELA-CFTR-ELB MESSAGE _____h_____
 _____440i_____123 TO 4622 OF HUMAN CFTR CDNA_____480i_____490>

970 980 990 1000 1010 1020
 CGATTTATCT AGGCATAGGC TTATGCCCTC TCTTTATTGT GAGGACACTG CTCCTACACC

GCTAAATAGA TCCGTATCCG AATACGGAAG AGAAATAACA CTCCTGTGAC GAGGATGTGG
 A I Y L G I G L C L L F I V R T L L L H>
 ____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____>
 ____h____HYBRID ELA-CFTR-ELB MESSAGE____h____>
 ____500i____123 TO 4622 OF HUMAN CFTR CDNA____540i____550>

1030 1040 1050 1060 1070 1080

CAGCCATTTT TGGCCITCAT CACATTGGAA TGCAGATGAG AATAGCTATG TTTAGTTTGA
 GTCGGTAAAA ACCGGAAGTA GTGTAACCTT ACGTCTACTC TTATCGATAC AAATCAAAC
 P A I F G L H H I G M Q M R I A M F S L>
 ____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____>
 ____h____HYBRID ELA-CFTR-ELB MESSAGE____h____>
 ____560i____123 TO 4622 OF HUMAN CFTR CDNA____600i____610>

1090 1100 1110 1120 1130 1140

TTTATAAGAA GACTTTAAAG CTGTCAAGCC GTGTTCTAGA TAAAATAAGT ATTGGACAAC
 AAATATTCTT CTGAAATTTC GACAGTTCGG CACAAGATCT ATTTTATTC TAACCTGTG
 I Y K K T L K L S S R V L D K I S I G Q>
 ____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____>
 ____h____HYBRID ELA-CFTR-ELB MESSAGE____h____>
 ____620i____123 TO 4622 OF HUMAN CFTR CDNA____660i____670>

1150 1160 1170 1180 1190 1200

TTGTTAGTCT CCTTTCCAAC AACCTGAACA AATTTGATGA AGGACTTGCA TTGGCACATT
 AACAATCAGA GGAAAGGTTG TTGGACTTGT TTAAGTACT TCCTGAACGT AACCGTGTA
 L V S L L S N N L N K F D E G L A L A H>
 ____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____>
 ____h____HYBRID ELA-CFTR-ELB MESSAGE____h____>
 ____680i____123 TO 4622 OF HUMAN CFTR CDNA____720i____730>

1210 1220 1230 1240 1250 1260

TCGTGTGGAT CGCTCCTTTG CAAGTGGCAC TCCTCATGGG GCTAATCTGG GAGTTGTTAC
 AGCACACCTA GCGAGGAAAC GTTCACCGTG AGGAGTACCC CGATTAGACC CTCAACAATG
 F V W I A P L Q V A L L M G L I W E L L>
 ____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____>
 ____h____HYBRID ELA-CFTR-ELB MESSAGE____h____>
 ____740i____123 TO 4622 OF HUMAN CFTR CDNA____780i____790>

1270 1280 1290 1300 1310 1320

AGGCGTCTGC CTTCTGTGGA CTTGGTTTCC TGATAGTCTT TGCCCTTTTT CAGGCTGGGC
 TCCGCAGACG GAAGACACCT GAACCAAAGG ACTATCAGGA ACGGGAAAA GTCCGACCCG
 Q A S A F C G L G F L I V L A L F Q A G>
 ____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____>
 ____h____HYBRID ELA-CFTR-ELB MESSAGE____h____>
 ____800i____123 TO 4622 OF HUMAN CFTR CDNA____840i____850>

1330 1340 1350 1360 1370 1380

TAGGGAGAAT GATGATGAAG TACAGAGATC AGAGAGCTGG GAAGATCAGT GAAAGACTTG
 ATCCCTCTTA CTACTACTTC ATGTCTCTAG TCTCTCGACC CTTCTAGTCA CTTTCTGAAAC
 L G R M M M K Y R D Q R A G K I S E R L>
 ____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____>
 ____h____HYBRID ELA-CFTR-ELB MESSAGE____h____>
 ____860i____123 TO 4622 OF HUMAN CFTR CDNA____900i____910>

1390 1400 1410 1420 1430 1440

TGATTACCTC AGAAATGATT GAAAACATCC AATCTGTAA GGCATACTGC TGGGAAGAAG
 ACTAATGGAG TCTTTACTAA CTTTGTAGG TTAGACAATT CCGTATGACG ACCCTTCTTC
 V I T S E M I E N I Q S V K A Y C W E E
 _____ CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON _____>
 _____ h _____ HYBRID ELA-CFTR-ELB MESSAGE _____ h _____>
 _____ 920i _____ 123 TO 4622 OF HUMAN CFTR CDNA _____ 960i _____ 970>

1450 1460 1470 1480 1490 1500

CAATGGAAAA AATGATTGAA AACTTAAGAC AACAGAACT GAAACTGACT CGGAAGGCAG
 GTTACCTTTT TTACTAACTT TTGAATTCTG TTTGTCTTGA CTTTGACTGA GCCTTCCGTC
 A M E K M I E N L R Q T E L K L T R K A
 _____ CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON _____>
 _____ h _____ HYBRID ELA-CFTR-ELB MESSAGE _____ h _____>
 _____ 980i _____ 123 TO 4622 OF HUMAN CFTR CDNA _____ 1020i _____ 1030>

1510 1520 1530 1540 1550 1560

CCTATGTGAG ATACTTCAAT AGCTCAGCCT TCTTCTTCTC AGGGTTCTTT GTGGTGTTTT
 GGATACACTC TATGAAGTTA TCGAGTCGGA AGAAGAAGAG TCCCAAGAAA CACCACAAAA
 A Y V R Y F N S S A F F F S G F F V V F
 _____ CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON _____>
 _____ h _____ HYBRID ELA-CFTR-ELB MESSAGE _____ h _____>
 _____ 1040i _____ 123 TO 4622 OF HUMAN CFTR CDNA _____ 1080i _____ 1090>

1570 1580 1590 1600 1610 1620

TATCTGTGCT TCCCTATGCA CTAATCAAAG GAATCATCCT CCGGAAAATA TTCACACCA
 ATAGACACGA AGGGATACGT GATTAGTTTC CTAGTAGGA GGCCTTTTAT AAGTGGTGGT
 L S V L P Y A L I K G I I L R K I F T T
 _____ CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON _____>
 _____ h _____ HYBRID ELA-CFTR-ELB MESSAGE _____ h _____>
 _____ 1100i _____ 123 TO 4622 OF HUMAN CFTR CDNA _____ 1140i _____ 1150>

1630 1640 1650 1660 1670 1680

TCTCATTCCTG CATGTGTTCTG CGCATGGCGG TCACTCGGCA ATTTCCCTGG GCTGTACAAA
 AGAGTAAGAC GTAACAAGAC GCGTACCGCC AGTGAGCCGT TAAAGGGACC CGACATGTTT
 I S F C I V L R M A V T R Q F P W A V Q
 _____ CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON _____>
 _____ h _____ HYBRID ELA-CFTR-ELB MESSAGE _____ h _____>
 _____ 1160i _____ 123 TO 4622 OF HUMAN CFTR CDNA _____ 1200i _____ 1210>

1690 1700 1710 1720 1730 1740

CATGSTATGA CTCTCTTGGG GCAATAAACA AATACAGGA TTTCTTACAA AGCAAGAAT
 GTACCATACT GAGAGAACCT CCGTATTGTT TTTATGTCCT AAAGAATGTT TTCCTTCTTA
 T W Y D S L G A I N K I Q D F L Q K Q E
 _____ CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON _____>
 _____ h _____ HYBRID ELA-CFTR-ELB MESSAGE _____ h _____>
 _____ 1220i _____ 123 TO 4622 OF HUMAN CFTR CDNA _____ 1260i _____ 1270>

1750 1760 1770 1780 1790 1800

ATAAGACATT GGAATATAAC TTAACGACTA CAGAAGTAGT GATGGAGAAAT GTAACAGCCT
 TATTCTGTAA CCTTATATTG AATTGCTGAT GTCTTCATCA CTACCTCTTA CATTGTCCGA
 Y K T L E Y N L T T T E V V M E N V T A
 _____ CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON _____>
 _____ h _____ HYBRID ELA-CFTR-ELB MESSAGE _____ h _____>
 _____ 1280i _____ 123 TO 4622 OF HUMAN CFTR CDNA _____ 1320i _____ 1330>

1810 1820 1830 1840 1850 1860

TCTGGGAGGA GGGATTGCGG GAATTATTTG AGAAAGCAAA ACAAAACAAT AACAAATAGAA
 AGACCCTCCT CCCTAAACCC CTTAATAAAC TCTTTCGTTT TGTTTTGTTA TTGTTATCTT
 F W E E G F G E L F E K A K Q N N N N R>
 ____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____>
 ____h____HYBRID ELA-CFTR-ELB MESSAGE____h____>
 ____1340i____123 TO 4622 OF HUMAN CFTR CDNA____1380i____1390>

1870 1880 1890 1900 1910 1920

AAACTTCTAA TGGTGATGAC AGCCTCTTCT TCAGTAATTT CTCACTTCTT GGTACTCCTG
 TTTGAAGATT ACCACTACTG TCGGAGAAGA AGTCATTAAA GAGTGAAGAA CCATGAGGAC
 K T S N G D D S L F F S N F S L L G T P>
 ____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____>
 ____h____HYBRID ELA-CFTR-ELB MESSAGE____h____>
 ____1400i____123 TO 4622 OF HUMAN CFTR CDNA____1440i____1450>

1930 1940 1950 1960 1970 1980

TCCTGAAAGA TATTAATTTT AAGATAGAAA GAGGACAGTT GTTGGCGGTT GCTGGATCCA
 AGGACTTTCT ATAATTAAAG TTCTATCTTT CTCCTGTCAA CAACCGCCAA CGACCTAGGT
 V L K D I N F K I E R G Q L L A V A G S>
 ____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____>
 ____h____HYBRID ELA-CFTR-ELB MESSAGE____h____>
 ____1460i____123 TO 4622 OF HUMAN CFTR CDNA____1500i____1510>

1990 2000 2010 2020 2030 2040

CTGGAGCAGG CAAGACTTCA CTTCTAATGA TGATTATGGG AGAACTGGAG CCTTCAGAGG
 GACCTCGTCC GTTCTGAAGT GAAGATTACT ACTAATACCC TCTTGACCTC GGAAGTCTCC
 T G A G K T S L L M M I M G E L E P S E>
 ____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____>
 ____h____HYBRID ELA-CFTR-ELB MESSAGE____h____>
 ____1520i____123 TO 4622 OF HUMAN CFTR CDNA____1560i____1570>

2050 2060 2070 2080 2090 2100

GTAAAATTAA GCACAGTGGA AGAATTTTCT TCTGTTCTCA GTTTTCCTGG ATTATGCCTG
 CATTTTAATT CGTGTCACCT TCTTAAAGTA AGACAAGAGT CAAAAGGACC TAATACGGAC
 G K I K H S G R I S F C S Q F S W I M P>
 ____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____>
 ____h____HYBRID ELA-CFTR-ELB MESSAGE____h____>
 ____1580i____123 TO 4622 OF HUMAN CFTR CDNA____1620i____1630>

2110 2120 2130 2140 2150 2160

GCACCATTAAG AAAAATATC ATCTTTGGTG TTTCCTATGA TGAATATAGA TACAGAAGCC
 CGTGTAATT TCTTTTATAG TAGAAACCAC AAAGGATACT ACTTATATCT ATGTCTTCGC
 G T I K E N I I F G V S Y D E Y R Y R S>
 ____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____>
 ____h____HYBRID ELA-CFTR-ELB MESSAGE____h____>
 ____1640i____123 TO 4622 OF HUMAN CFTR CDNA____1680i____1690>

2170 2180 2190 2200 2210 2220

TCATCAAGC ATGCCAACTA GAGAGGACA TCTCCAAGTT TGCAGAGAAA GACAATATAG
 AGTAGTTTCG TACGGTTGAT CTTCTCCTGT AGAGTTTCAA ACGTCTCTTT CTGTTATATC
 V I K A C Q L E E D I S K F A E K D N I>
 ____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____>
 ____h____HYBRID ELA-CFTR-ELB MESSAGE____h____>
 ____1700i____123 TO 4622 OF HUMAN CFTR CDNA____1740i____1750>

-72-

2230	2240	2250	2260	2270	2280
TTCTTGGAGA AGGTGGAATC ACACTGAGTG GAGGTCAACG AGCAAGAATT TCTTTAGCAA AAGAACCTCT TCCACCTTAG TGTGACTCAC CTCCAGTTGC TCGTTCTTAA AGAAATCGTT V L G E G G I T L S G G Q R A R I S L A> ____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____> ____h____HYBRID ELA-CFTR-ELB MESSAGE____h____> ____1760i____123 TO 4622 OF HUMAN CFTR CDNA____1800i____1810>					
2290	2300	2310	2320	2330	2340
GAGCAGTATA CAAAGATGCT GATTGTATT TATTAGACTC TCCTTTTGGG TACCTAGATG CTCGTCATAT GTTCTACGA CTAAACATAA ATAATCTGAG AGGAAAACCT ATGGATCTAC R A V Y K D A D L Y L L D S P F G Y L D> ____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____> ____h____HYBRID ELA-CFTR-ELB MESSAGE____h____> ____1820i____123 TO 4622 OF HUMAN CFTR CDNA____1860i____1870>					
2350	2360	2370	2380	2390	2400
TTTTAACAGA AAAAGAAATA TTTGAAAGCT GTGTCTGTAA ACTGATGGCT AACAAAATA AAAATTGTCT TTTTCTTTAT AAACCTTCGA CACAGACATT TGACTACCGA TTGTTTGTAT V L T E K E I F E S C V C K L M A N K T> ____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____> ____h____HYBRID ELA-CFTR-ELB MESSAGE____h____> ____1880i____123 TO 4622 OF HUMAN CFTR CDNA____1920i____1930>					
2410	2420	2430	2440	2450	2460
GGATTTTGGT CACTTCTAAA ATGGAACATT TAAAGAAAGC TGACAAAATA TTAATTTTGC CCTAAAACCA GTGAAGATTT TACCTGTGTA ATTTCTTTTCG ACTGTTTTAT AATTAAAACG R I L V T S K M E H L K K A D K I L I L> ____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____> ____h____HYBRID ELA-CFTR-ELB MESSAGE____h____> ____1940i____123 TO 4622 OF HUMAN CFTR CDNA____1980i____1990>					
2470	2480	2490	2500	2510	2520
ATGAAGGTAG CAGCTATTTT TATGGGACAT TTTCAGAACT CCAAATCTA CAGCCAGACT TACTTCCATC GTCGATAAAA ATACCCCTGA AAAGTCTTGA GGTTTTAGAT GTCGGTCTGA H E G S S Y F Y G T F S E L Q N L Q P D> ____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____> ____h____HYBRID ELA-CFTR-ELB MESSAGE____h____> ____2000i____123 TO 4622 OF HUMAN CFTR CDNA____2040i____2050>					
2530	2540	2550	2560	2570	2580
TTAGCTCAA ACTCATGGGA TGTGATTCTT TCGACCAATT TAGTGCAGAA AGAAGAAATT AATCGAGTTT TGAGTACCCT AACTAAGAA AGCTGGTTAA ATCAGTCTT TCTTCTTTAA F S S K L M G C D S F D Q F S A E R R N> ____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____> ____h____HYBRID ELA-CFTR-ELB MESSAGE____h____> ____2060i____123 TO 4622 OF HUMAN CFTR CDNA____2100i____2110>					
2590	2600	2610	2620	2630	2640
CAATCCTAAC TGAGACCTTA CACCGTTTCT CATTAGAAAG AGATGCTCCT GTCTCCTGGA GTTAGGATTG ACTCTGGAAT GTGGCAAGA GTAATCTTCC TCTACGAGGA CAGAGGACCT S I L T E T L H R F S L E G D A P V S W> ____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____> ____h____HYBRID ELA-CFTR-ELB MESSAGE____h____> ____2120i____123 TO 4622 OF HUMAN CFTR CDNA____2160i____2170>					

-73-

2650 2660 2670 2680 2690 2700
 CAGAAACAAA AAAACAATCT TTAAACAGA CTGGAGAGTT TGGGGAAAA AGGAAGAATT
 GTCTTTGTTT TTTTGTTAGA AAATTTGTCT GACCTCTCAA ACCCCTTTT TCCTTCTTAA
 T E T K K Q S F K Q T G E F G E K R K N>
 _____ CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON _____>
 _____ h _____ HYBRID ELA-CFTR-ELB MESSAGE _____>
 _____ 2180i _____ 123 TO 4622 OF HUMAN CFTR CDNA _____ 2220i _____ 2230>

2710 2720 2730 2740 2750 2760
 CTATTCTCAA TCCAATCAAC TCTATACGAA AATTTTCCAT TGTGCAAAAG ACTCCCTTAC
 GATAAGAGTT AGGTTAGTTG AGATATGCTT TTAAAAGGTA ACACGTTTTC TGAGGGGAATG
 S I L N P I N S I R K F S I V Q K T P L>
 _____ CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON _____>
 _____ h _____ HYBRID ELA-CFTR-ELB MESSAGE _____>
 _____ 2240i _____ 123 TO 4622 OF HUMAN CFTR CDNA _____ 2280i _____ 2290>

2770 2780 2790 2800 2810 2820
 AAATGAATGG CATCGAAGAG GATTCTGATG AGCCTTTAGA GAGAAGGCTG TCCTTAGTAC
 TTTACTTACC GTAGCTTCTC CTAAGACTAC TCGGAAATCT CTCTTCCGAC AGGAATCATG
 Q M N G I E E D S D E P L E R R L S L V>
 _____ CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON _____>
 _____ h _____ HYBRID ELA-CFTR-ELB MESSAGE _____>
 _____ 2300i _____ 123 TO 4622 OF HUMAN CFTR CDNA _____ 2340i _____ 2350>

2830 2840 2850 2860 2870 2880
 CAGATTCTGA GCAGGGAGAG GCGATACTGC CTCGCATCAG CGTGATCAGC ACTGGCCCCA
 GTCTAAGACT CGTCCCTCTC CGCTATGACG GAGCGTAGTC GCACTAGTCG TGACCGGGGT
 P D S E Q G E A I L P R I S V I S T G P>
 _____ CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON _____>
 _____ h _____ HYBRID ELA-CFTR-ELB MESSAGE _____>
 _____ 2360i _____ 123 TO 4622 OF HUMAN CFTR CDNA _____ 2400i _____ 2410>

2890 2900 2910 2920 2930 2940
 CGCTTCAGGC ACCAAGGAGG CAGTCTGTCC TGAACCTGAT GACACACTCA GTTAACCAAG
 GCGAAGTCCG TGCTTCCTCC GTCAGACAGG ACTTGGACTA CTGTGTGAGT CAATTGGTTC
 T L Q A R R R Q S V L N L M T H S V N Q>
 _____ CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON _____>
 _____ h _____ HYBRID ELA-CFTR-ELB MESSAGE _____>
 _____ 2420i _____ 123 TO 4622 OF HUMAN CFTR CDNA _____ 2460i _____ 2470>

2950 2960 2970 2980 2990 3000
 GTCAGACAT TCACCGAAG ACAACAGCAT CCACACGAA AGTGTCACTG GCCCCTCAGG
 CAGTCTTGTA ACTGGCTTTC TGTGTGCGTA GGTGTGCTTT TCACAGTGAC CGGGGAGTCC
 G Q N I H R K T T A S T R K V S L A P Q>
 _____ CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON _____>
 _____ h _____ HYBRID ELA-CFTR-ELB MESSAGE _____>
 _____ 2480i _____ 123 TO 4622 OF HUMAN CFTR CDNA _____ 2520i _____ 2530>

3010 3020 3030 3040 3050 3060
 CAACTTGCAC TGAAGTGGAT ATATATTCAA GAAGTTATC TCAAGAACT GGCTTGGAAA
 GTTTGAAGCTG ACTTGACCTA TATATAAGTT CTTCCAATAG AGTTCTTTGA CCGAAGCTTT
 A N L T E L D I Y S R R L S Q E T G L E>
 _____ CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON _____>
 _____ h _____ HYBRID ELA-CFTR-ELB MESSAGE _____>

_____2540i_____123 TO 4622 OF HUMAN CFTR CDNA_____2580i_____2590>

3070 3080 3090 3100 3110 3120

TAAGTGAAGA AATTAACGAA GAAGACTTAA AGGAGTGCCT TTTTGATGAT ATGGAGAGCA
ATTCACITCT TTAATTGCTT CTTCCTGAATT TCCTCACGGA AAAACTACTA TACCTCTCGT
I S E E I N E E D L K E C L F D D M E S>

____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____>
_____h_____HYBRID ELA-CFTR-ELB MESSAGE_____h_____>

_____2600i_____123 TO 4622 OF HUMAN CFTR CDNA_____2640i_____2650>

3130 3140 3150 3160 3170 3180

TACCAGCAGT GACTACATGG AACACATACC TTCGATATAT TACTGTCCAC AAGAGCTTAA
ATGGTCGTCA CTGATGTACC TTGTGTATGG AAGCTATATA ATGACAGGTG TTCTCGAATT
I P A V T T W N T Y L R Y I T V H K S L>

____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____>
_____h_____HYBRID ELA-CFTR-ELB MESSAGE_____h_____>

_____2660i_____123 TO 4622 OF HUMAN CFTR CDNA_____2700i_____2710>

3190 3200 3210 3220 3230 3240

TTTTTGTCCT AATTGGTGC TTAGTAATTT TTCTGGCAGA GGTGGCTGCT TCTTTGGTTG
AAAAACACGA TTAAACCACG AATCATTAAA AAGACCGTCT CCACCGACGA AGAAACCAAC
I F V L I W C L V I F L A E V A A S L V>

____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____>
_____h_____HYBRID ELA-CFTR-ELB MESSAGE_____h_____>

_____2720i_____123 TO 4622 OF HUMAN CFTR CDNA_____2760i_____2770>

3250 3260 3270 3280 3290 3300

TGCTGTGGCT CCTTGGAAAC ACTCCTCTTC AAGACAAAGG GAATAGTACT CATAGTAGAA
ACGACACCGA GGAACCTTTG TGAGGAGAAG TTCTGTTTCC CTTATCATGA GTATCATCTT
V L W L L G N T P L Q D K G N S T H S R>

____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____>
_____h_____HYBRID ELA-CFTR-ELB MESSAGE_____h_____>

_____2780i_____123 TO 4622 OF HUMAN CFTR CDNA_____2820i_____2830>

3310 3320 3330 3340 3350 3360

ATAACAGCTA TGCAGTGATT ATCACCAGCA CCAGTTCGTA TTATGTGTTT TACATTTACG
TATTGTGCGAT ACGTCACTAA TAGTGGTCGT GGTCAAGCAT AATACACAAA ATGTAAATGC
N N S Y A V I I T S T S S Y Y V F Y I Y>

____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____>
_____h_____HYBRID ELA-CFTR-ELB MESSAGE_____h_____>

_____2840i_____123 TO 4622 OF HUMAN CFTR CDNA_____2880i_____2890>

3370 3380 3390 3400 3410 3420

TGGGAGTAGC CGACACTTTG CTTGCTATGG GATTCTTCAG AGGTCTACCA CTGGTGCATA
ACCTTCATCG GCTGTGAAAC GAACGATACC CTAAGAAGTC TCCAGATGGT GACCACGTAT
V G V A D T L L A M G F F R G L P L V H>

____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____>
_____h_____HYBRID ELA-CFTR-ELB MESSAGE_____h_____>

_____2900i_____123 TO 4622 OF HUMAN CFTR CDNA_____2940i_____2950>

3430 3440 3450 3460 3470 3480

CTCTAATCAC AGTGTGCAAA ATTTTACACC ACAAATGTT ACATTCTGTT CTTCAAGCAC
GAGATTAGTG TCACAGCTTT TAAAATGTGG TGTITTACAA TGTAAGACAA GAAGTTCGTG
T L I T V S K I L H H K M L H S V L Q A>

____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____>

_____h_____HYBRID ELA-CFTR-E1B MESSAGE _____h_____
 _____2960i_____123 TO 4622 OF HUMAN CFTR CDNA _____3000i_____3010>
 3490 3500 3510 3520 3530 3540
 CTATGTCAAC CCTCAACACG TTGAAAGCAG GTGGGATTCT TAATAGATTG TCCAAGATA
 GATACAGTTG GGAGTTGTGC AACTTTCGTC CACCCTAAGA ATTATCTAAG AGGTTTCTAT
 P M S T L N T L K A G G I L N R F S K D>
 _____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON_____
 _____h_____HYBRID ELA-CFTR-E1B MESSAGE _____h_____
 _____3020i_____123 TO 4622 OF HUMAN CFTR CDNA _____3060i_____3070>
 3550 3560 3570 3580 3590 3600
 TAGCAATTTT GGATGACCTT CTGCCTCTTA CCATATTTGA CTTCATCCAG TTGTTATTAA
 ATCGTTAAAA CCTACTGGAA GACGGAGAAT GGTATAAACT GAAGTAGGTC AACAATAATT
 I A I L D D L L P L T I F D F I Q L L L>
 _____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON_____
 _____h_____HYBRID ELA-CFTR-E1B MESSAGE _____h_____
 _____3080i_____123 TO 4622 OF HUMAN CFTR CDNA _____3120i_____3130>
 3610 3620 3630 3640 3650 3660
 TTGTGATTGG AGCTATAGCA GTTGTGCGAG TTTTACAACC CTACATCTTT GTTGCAACAG
 AACACTAACC TCGATATCGT CAACAGCGTC AAAATGTTGG GATGTAGAAA CAACGTTGTC
 I V I G A I A V V A V L Q P Y I F V A T>
 _____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON_____
 _____h_____HYBRID ELA-CFTR-E1B MESSAGE _____h_____
 _____3140i_____123 TO 4622 OF HUMAN CFTR CDNA _____3180i_____3190>
 3670 3680 3690 3700 3710 3720
 TGCCAGTGAT AGTGGCTTTT ATTATGTTGA GAGCATATTT CCTCCAAACC TCACAGCAAC
 ACGGTCACCTA TCACCGAAAA TAATACAACCT CTCGTATAAA GGAGGTTTGG AGTGTCGTTG
 V P V I V A F I M L R A Y F L Q T S Q Q>
 _____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON_____
 _____h_____HYBRID ELA-CFTR-E1B MESSAGE _____h_____
 _____3200i_____123 TO 4622 OF HUMAN CFTR CDNA _____3240i_____3250>
 3730 3740 3750 3760 3770 3780
 TCAAACAACCT GGAATCTGAA GGCAGGAGTC CAATTTTCAC TCATCTTGTT ACAAGCTTAA
 AGTTTGTGTA CCTTAGACTT CCGTCCTCAG GTTAAAAGTG AGTAGAACAA TGTTCCGAATT
 L K Q L E S E G R S P I F T H L V T S L>
 _____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON_____
 _____h_____HYBRID ELA-CFTR-E1B MESSAGE _____h_____
 _____3260i_____123 TO 4622 OF HUMAN CFTR CDNA _____3300i_____3310>
 3790 3800 3810 3820 3830 3840
 AAGGACTATG GACACTTCGT GCCTTCGGAC GGCAGCCTTA CTTTGAAACT CTGTTCCACA
 TTCCTGATAC CTGTGAAGCA CGGAAGCCTG CCGTCGGAAT GAAACTTTGA GACAAGGTGT
 K G L W T L R A F G R Q P Y F E T L F H>
 _____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON_____
 _____h_____HYBRID ELA-CFTR-E1B MESSAGE _____h_____
 _____3320i_____123 TO 4622 OF HUMAN CFTR CDNA _____3360i_____3370>
 3850 3860 3870 3880 3890 3900
 AAGCTCTGAA TTTACATACT GCCAACTGGT TCTTGACCT GTCAACACTG CGCTGTTTCC
 TTCGAGACTT AAATGTATGA CGTTTGACCA AGAACATGGA CAGTTGTGAC GCGACCAAGG
 K A L N L H T A N W F L Y L S T L R W F>

____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____>
 _____h_____HYBRID ELA-CFTR-ELB MESSAGE _____h_____>
 _____3380i_____123 TO 4622 OF HUMAN CFTR CDNA _____3420i_____3430>

3910 3920 3930 3940 3950 3960

AAATGAGAAT AGAAATGATT TTGTGTCATCT TCTTCATTGC TGTTACCTTC ATTTCATTT
 TTTACTCTTA TCTTTACTAA AAACAGTAGA AGAAGTAACG ACAATGGAAG TAAAGGTAAA
 Q M R I E M I F V I F F I A V T F I S I>

____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____>
 _____h_____HYBRID ELA-CFTR-ELB MESSAGE _____h_____>
 _____3440i_____123 TO 4622 OF HUMAN CFTR CDNA _____3480i_____3490>

3970 3980 3990 4000 4010 4020

TAACAACAGG AGAAGGAGAA GGAAGAGTTG GTATTATCCT GACTTTAGCC ATGAATATCA
 ATTGTGTGCC TCTTCCTCTT CCTTCTCAAC CATAATAGGA CTGAAATCGG TACTTATAGT
 L T T G E G E G R V G I I L T L A M N I>

____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____>
 _____h_____HYBRID ELA-CFTR-ELB MESSAGE _____h_____>
 _____3500i_____123 TO 4622 OF HUMAN CFTR CDNA _____3540i_____3550>

4030 4040 4050 4060 4070 4080

TGAGTACATT GCAGTGGGCT GTAAACTCCA GCATAGATGT GGATAGCTTG ATGCGATCTG
 ACTCATGTAA CGTCACCCGA CATTGAGGT CGTATCTACA CCTATCGAAC TACGCTAGAC
 M S T L Q W A V N S S I D V D S L M R S>

____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____>
 _____h_____HYBRID ELA-CFTR-ELB MESSAGE _____h_____>
 _____3560i_____123 TO 4622 OF HUMAN CFTR CDNA _____3600i_____3610>

4090 4100 4110 4120 4130 4140

TGAGCCGAGT CTTTAAGTTC ATTGACATGC CAACAGAAGG TAAACCTACC AAGTCAACCA
 ACTCGGCTCA GAAATTCAAG TAACTGTACG GTTGCTCTCC ATTTGGATGG TTCAGTTGGT
 V S R V F K F I D M P T E G K P T K S T>

____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____>
 _____h_____HYBRID ELA-CFTR-ELB MESSAGE _____h_____>
 _____3620i_____123 TO 4622 OF HUMAN CFTR CDNA _____3660i_____3670>

4150 4160 4170 4180 4190 4200

AACCATACAA GAATGGCCAA CTCTCGAAAG TTATGATTAT TGAGAATTCA CACGTGAAGA
 TTGGTATGTT CTTACCGGTT GAGAGCTTTC AATACTAATA ACTCTTAAGT GTGCACTTCT
 K P Y K N G Q L S K V M I I E N S H V K>

____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____>
 _____h_____HYBRID ELA-CFTR-ELB MESSAGE _____h_____>
 _____3680i_____123 TO 4622 OF HUMAN CFTR CDNA _____3720i_____3730>

4210 4220 4230 4240 4250 4260

AAGATGACAT CTGCCCCCTCA GGGGGCCAAA TGACTGTCAA AGATCTCACA GCAAAATACA
 TTCTACTGTA GACCGGGAGT CCCCCGGTTT ACTGACAGTT TCTAGAGTGT CGTTTATGT
 K D D I W P S G G Q M T V K D L T A K Y>

____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____>
 _____h_____HYBRID ELA-CFTR-ELB MESSAGE _____h_____>
 _____3740i_____123 TO 4622 OF HUMAN CFTR CDNA _____3780i_____3790>

4270 4280 4290 4300 4310 4320

CAGAAGGTGG AAATGCCATA TTAGAGAAC TTTCTCTCTC AATAAGTCCT GGGCAGAGGC
 GTCTTCCACC TTTACGGTAT AATCTCTTCT AAAGGAAGAG TTATTCAGGA CCGGTCTCCC

-77-

T E G G N A I L E N I S F S I S P G Q R>
 ____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____>
 ____h____HYBRID ELA-CFTR-ELB MESSAGE____h____>
 ____3800i____123 TO 4622 OF HUMAN CFTR CDNA____3840i____3850>
 4330 4340 4350 4360 4370 4380

TGGGCCTCTT GGAAGAAGT GGATCAGGGA AGAGTACTTT GTTATCAGCT TTTTGTGAGAC
 ACCCGGAGAA CCTTCTTGA CCTAGTCCCT TCTCATGAAA CAATAGTCGA AAAAAGTCTG
 V G L L G R T G S G K S T L L S A F L R>
 ____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____>
 ____h____HYBRID ELA-CFTR-ELB MESSAGE____h____>
 ____3860i____123 TO 4622 OF HUMAN CFTR CDNA____3900i____3910>
 4390 4400 4410 4420 4430 4440

TACTGAACAC TGAAGGAGAA ATCCAGATCG ATGGTGTGTC TTGGGATTCA ATAAGTTTGC
 ATGACTTGTG ACTTCTCTT TAGGTCTAGC TACCACACAG AACCTAAGT TATTGAAACG
 L L N T E G E I Q I D G V S W D S I T L>
 ____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____>
 ____h____HYBRID ELA-CFTR-ELB MESSAGE____h____>
 ____3920i____123 TO 4622 OF HUMAN CFTR CDNA____3960i____3970>
 4450 4460 4470 4480 4490 4500

AACAGTGGAG GAAAGCCTTT GGAGTGATAC CACAGAAAGT ATTTATTTTT TCTGGAACAT
 TTGTCACTC CTTTCGGAAA CCTCACTATG GTGTCTTTCA TAAATAAAAA AGACCTTGTA
 Q Q W R K A F G V I P Q K V F I F S G T>
 ____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____>
 ____h____HYBRID ELA-CFTR-ELB MESSAGE____h____>
 ____3980i____123 TO 4622 OF HUMAN CFTR CDNA____4020i____4030>
 4510 4520 4530 4540 4550 4560

TTAGAAAAAA CTGGATCCC TATGAACAGT GGAGTGATCA AGAAATATGG AAAGTTGCAG
 AATCTTTTTT GAACCTAGGG ATACTTGTC CCTCACTAGT TCTTTATACC TTTCAACGTC
 F R K N L D P Y E Q W S D Q E I W K V A>
 ____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____>
 ____h____HYBRID ELA-CFTR-ELB MESSAGE____h____>
 ____4040i____123 TO 4622 OF HUMAN CFTR CDNA____4080i____4090>
 4570 4580 4590 4600 4610 4620

ATGAGGTTGG GCTCAGATCT GTGATAGAAC AGTTTCCTGG GAAGCTTGAC TTTGTCTTGG
 TACTCCAACC CGAGTCTAGA CACTATCTTG TCAAGGACC CTTCGAAGTC AACAGGAGC
 D E V G L R S V I E Q F P G K L D F V L>
 ____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____>
 ____h____HYBRID ELA-CFTR-ELB MESSAGE____h____>
 ____4100i____123 TO 4622 OF HUMAN CFTR CDNA____4140i____4150>
 4630 4640 4650 4660 4670 4680

TGGATGGGGG CTGTGCTCTA AGCCATGGCC ACPAGCAGTT GATGTGCTTG GCTAGATCTG
 ACCTACCCCC GACACAGGAT TCGGTACCCG TGTTCTGCAA CTACACGAAC CGATCTAGAC
 V D G G C V L S H G H K Q L M C L A R S>
 ____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____>
 ____h____HYBRID ELA-CFTR-ELB MESSAGE____h____>
 ____4160i____123 TO 4622 OF HUMAN CFTR CDNA____4200i____4210>
 4690 4700 4710 4720 4730 4740

TTCTCAGTAA GCGAAGATC TTGCTGCTG ATAAATCCAG TCTCTATTTC GATCCAGTAA

AAGAGTCATT CCGCTTCTAG AACGACGAAC TACTTGGGTC ACGAGTAAAC CTAGGTCATT
V L S K A K I L L L D E P S A H L D P V>
____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____>
____h____HYBRID ELA-CFTR-ELB MESSAGE____h____>
____4220i____123 TO 4622 OF HUMAN CFTR CDNA____4260i____4270>

4750 4760 4770 4780 4790 4800

CATACCAAAT AATTAGAAGA ACTCTAAAC AAGCATTTGC TGATTGCACA GTAATTCTCT
GTATGGTTTA TTAATCTTCT TGAGATTTTG TTCGTAAACG ACTAACGTGT CATTAGAGA
T Y Q I I R R T L K Q A F A D C T V I L>
____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____>
____h____HYBRID ELA-CFTR-ELB MESSAGE____h____>
____4280i____123 TO 4622 OF HUMAN CFTR CDNA____4320i____4330>

4810 4820 4830 4840 4850 4860

GTGAACACAG GATAGAAGCA ATGCTGGAAT GCCAACAATT TTTGGTCATA GAAGAGAACA
CACTTGTGTC CTATCTTCGT TACGACCTTA CGGTTGTAA AAACCAGTAT CTTCTCTTGT
C E H R I E A M L E C Q Q F L V I E E N>
____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____>
____h____HYBRID ELA-CFTR-ELB MESSAGE____h____>
____4340i____123 TO 4622 OF HUMAN CFTR CDNA____4380i____4390>

4870 4880 4890 4900 4910 4920

AAGTGGCGCA GTACGATTCC ATCCAGAAAC TGCTGAACGA GAGGAGCCTC TTCCGGCAAG
TTCACGCCGT CATGCTAAGG TAGGTCTTTG ACGACTTGCT CTCCTCGGAG AAGGCCGTTC
K V R Q Y D S I Q K L L N E R S L F R Q>
____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____>
____h____HYBRID ELA-CFTR-ELB MESSAGE____h____>
____4400i____123 TO 4622 OF HUMAN CFTR CDNA____4440i____4450>

4930 4940 4950 4960 4970 4980

CCATCAGCCC CTCCGACAGG GTGAAGCTCT TTCCCCACCG GAACTCAAGC AAGTGCAAGT
GGTAGTCGGG GAGGCTGTCC CACTTCGAGA AAGGGGTGGC CTTGAGTTTCG TTCACGTTCA
A I S P S D R V K L F P H R N S S K C K>
____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____>
____h____HYBRID ELA-CFTR-ELB MESSAGE____h____>
____4460i____123 TO 4622 OF HUMAN CFTR CDNA____4500i____4510>

4990 5000 5010 5020 5030 5040

CTAAGCCCCA GATTGCTGCT CTGAAGAGG AGACAGAAGA AGAGGTGCAA GATACAGGC
GATTGGGGST CTAACGACGA GACTTCTCC TCTGTCTTCT TCTCCACGTT CTATGTTCCG
S K P Q I A A L K E E T E E E V Q D T R>
____CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR; CODON____>
____h____HYBRID ELA-CFTR-ELB MESSAGE____h____>
____4520i____123 TO 4622 OF HUMAN CFTR CDNA____4560i____4570>

5050 5060 5070 5080 5090 5100

TTTAGAGAGC AGCATAAATG TTGACATGGG ACATTTGCTC ATGGAATTGG AGGTAGCGGA
AAATCTCTCG TCGTATTTAC AACTGTACCC TGTAACGAG TACCTTAACC TCCATCGCCT
L >

____h____HYBRID ELA-CFTR-ELB MESSAGE____h____>

____4580i____123 TO 4622 OF HUMAN CFTR CDNA____4620i____>

-79-

5110 5120 5130 5140 5150 5160
 TTGAGGTA CT GAAATGTGTG GCGGTGGCTT AAGGGTGGGA AAGAATATAT AAGGTGGGGG
 AACTCCATGA CTTTACACAC CCGCACCGAA TTCCACCCTT TCTTATATA TTCCACCCCC
 _____h_____ HYBRID ELA-CFTR-ELB MESSAGE _____h_____>
 _____10_g_____ ELB 3' UNTRANSLATED SEQUENCES _____50_g_____60_____>
 _____k_____10_____k_____ ELB 3' INTRON _____k_____40_____k_____50_____>

5170 5180 5190 5200 5210 5220
 TCTCATGTAG TTTTGTATCT GTTTTGCAGC AGCCGCCGCC ATGAGCGCCA ACTCGTTTGA
 AGAGTACATC AAAACATAGA CAAAACGTCG TCGGCGGCGG TACTCGCGGT TGAGCAAAC T
 M S A N S F D
 _____IX PROTEIN (HE)_____>
 _____h_____ HYBRID ELA-CFTR-ELB MESSAGE _____h_____>
 _____1_____1_____1_____ IX MRNA _____1_____1_____>
 _____70_g_____ ELB 3' UNTRANSLATED SEQUENCES _____110_g_____120_____>
 _____60_____ ELB 3' INTRON _____80_____>

5230 5240 5250 5260 5270 5280
 TGGAAGCATT GTGAGCTCAT ATTTGACAAC GCGCATGCCC CCATGGGCGG GGGTGGGTCA
 ACCTTCGTAA CACTCGAGTA TAAACTGTTG CGCGTACGGG GGTACCCGGC CCCACGCAGT
 G S I V S S Y L T T R M P P W A G V R Q>
 _____IX PROTEIN (HEXON-ASSOCIATED PROTEIN); CODON_START=1_____>
 _____h_____ HYBRID ELA-CFTR-ELB MESSAGE _____h_____>
 _____1_____1_____1_____ IX MRNA _____1_____1_____>
 _____130_g_____ ELB 3' UNTRANSLATED SEQUENCES _____170_g_____180_____>

5290 5300 5310 5320 5330 5340
 GAATGTGATG GGCTCCAGCA TTGATGGTCG CCCGTCCTG CCCGCAAAC TACTACCTT
 CTTACACTAC CCGAGGTCGT AACTACCAGC GGGCAGGAC GGGCGTTTGA GATGATGGAA
 N V M G S S I D G R P V L P A N S T T L>
 _____IX PROTEIN (HEXON-ASSOCIATED PROTEIN); CODON_START=1_____>
 _____h_____ HYBRID ELA-CFTR-ELB MESSAGE _____h_____>
 _____1_____1_____1_____ IX MRNA _____1_____1_____>
 _____190_g_____ ELB 3' UNTRANSLATED SEQUENCES _____230_g_____240_____>

5350 5360 5370 5380 5390 5400
 GACCTACGAG ACCGTGTCTG GAACGCCGTT GGAGACTGCA GCCTCCGCCG CCGCTTCAGC
 CTGGATGCTC TGGCAGAGAC CTTGCGGCAA CCTCTGACGT CGGAGCGCGC GGCGAAGTCG
 T Y E T V S G T P L E T A A S A A S A>
 _____IX PROTEIN (HEXON-ASSOCIATED PROTEIN); CODON_START=1_____>
 _____h_____ HYBRID ELA-CFTR-ELB MESSAGE _____h_____>
 _____1_____1_____1_____ IX MRNA _____1_____1_____>
 _____250_g_____ ELB 3' UNTRANSLATED SEQUENCES _____290_g_____300_____>

5410 5420 5430 5440 5450 5460
 CGCTGCAGCC ACCGCCCGCG GGATTGTGAC TGACTTTGCT TTCCTGAGCC CGCTTGCAAG
 GCGACGTCGG TGGCGGCGCG CCTAACACTG ACTGAAACGA AAGGACTCGG GCGAACGTTT
 A A A T A R G I V T D F A F L S P L A S>
 _____IX PROTEIN (HEXON-ASSOCIATED PROTEIN); CODON_START=1_____>
 _____h_____ HYBRID ELA-CFTR-ELB MESSAGE _____h_____>
 _____1_____1_____1_____ IX MRNA _____1_____1_____>
 _____310_g_____ ELB 3' UNTRANSLATED SEQUENCES _____350_g_____360_____>

5470 5480 5490 5500 5510 5520
 CAGTGCAGCT TCCCGTTTCT CCGCCCGCGA TGACAACTTG ACGGCTCTTT TGGCAACATT

GTCACGTCGA AGGGCAAGTA GCGGGGCGCT ACTGTTCAAC TGCCGAGAAA ACCGTGTTAA
 S A A S R S S A R D D K L T A L L A Q L>
 _____IX PROTEIN (HEXON-ASSOCIATED PROTEIN); CODON_START=1____>
 _____h_____HYBRID E1A-CFTR-E1B MESSAGE _____h____>
 _____1_____1_____IX MRNA _____1_____1____>
 _____370_g_____E1B 3' UNTRANSLATED SEQUENCES _____410_g_____420____>

5530 5540 5550 5560 5570 5580

GGATTCTTTG ACCCGGGAAC TTAATGTCGT TTCTCAGCAG CTGTTGGATC TGCGCCAGCA
 CCTAAGAAAC TGGGCCCTTG AATTACAGCA AAGAGTCGTC GACAACCTAG ACGCGGTCGT
 D S L T R E L N V V S Q Q L L D L R Q Q>
 _____IX PROTEIN (HEXON-ASSOCIATED PROTEIN); CODON_START=1____>
 _____h_____HYBRID E1A-CFTR-E1B MESSAGE _____h____>
 _____1_____1_____IX MRNA _____1_____1____>
 _____430_g_____E1B 3' UNTRANSLATED SEQUENCES _____470_g_____480____>

5590 5600 5610 5620 5630

GGTTTCTGCC CTGAAGGCTT CCTCCCCTCC CAATGCGGTT TAAACATAA ATAAA
 CCAAAGACGG GACTTCCGAA GGAGGGGAGG GTTACGCCAA ATTTTGTATT TATTT
 V S A L K A S S P P N A V *>
 _____IX PROTEIN (HEXON-ASSOCIATED PROTEIN); C____>
 _____h_____HYBRID E1A-CFTR-E1B MESSAGE _____h____>
 _____1_____1_____IX MRNA _____1_____1____>
 _____490_g_____E1B 3' UNTRANSLATED SEQUENCES _____530_g_____>

Table III

Nucleotide Sequence Analysis of Ad2-ORF6/PGK-CFTR

LOCUS	AD2-ORF6/P 36335 BP DS-DNA			
DEFINITION	-			
ACCESSION	-			
KEYWORDS	-			
SOURCE	-			
FEATURES	From	To/Span	Description	
frag	12915	36335	10676 to 34096 of Ad2-E4/ORF6	
frag	35069	35973	33178 to 34082 of Ad2 seq	
pre-msg	> 35973	< 35069	(C)	E4 mRNA [Nucleic Acids Res. 9, 1675-1689 (1981)], [J. Mol. Biol. 149, 189-221 (1981)], [Nucleic Acids Res. 12, 3503-3519 (1984)], [Unpublished (1984)] [Split]
IVS	35794	35084	(C)	E4 mRNA intron D7 [J. Virol. 50, 106-117 (1984)], [Nucleic Acids Res. 12, 3503-3519 (1984)], [Unpublished (1984)]
IVS	35794	35175	(C)	E4 mRNA intron D6 [Nucleic Acids Res. 12, 3503-3519 (1984)]
IVS	35794	35268	(C)	E4 mRNA intron D5 [J. Virol. 50, 106-117 (1984)]
IVS	35794	35295	(C)	E4 mRNA intron D4 [J. Virol. 50, 106-117 (1984)]
IVS	35794	35343	(C)	E4 mRNA intron D3 [J. Virol. 50, 106-117 (1984)]
IVS	35794	35501	(C)	E4 mRNA intron D2 [J. Virol. 50, 106-117 (1984)]
IVS	35794	35570	(C)	E4 mRNA intron D1 [J. Virol. 50, 106-117 (1984)]
IVS	35794	35766	(C)	E4 mRNA intron D [J. Virol. 50, 106-117 (1984)]
frag	35978	36335	35580 to 35937 of Ad2 seq	
pre-msg	36007	< 35978	(C)	E4 mRNA [Nucleic Acids Res. 9, 1675-1689 (1981)], [J. Mol. Biol. 149, 189-221 (1981)], [Nucleic Acids Res. 12, 3503-3519 (1984)], [Unpublished (1984)] [Split]
rpt	36234	36335	inverted terminal repetition; 99.54% [Biochem. Biophys. Res. Commun. 87, 671-678 (1979)], [J. Mol. Biol. 128, 577-594 (1979)]	
frag	< 12915	35054	1 to 32815 of Ad2 seq [Split]	
pept	< 28478	28790	3	33K protein (virion morphogenesis)
pept	28478	28790	1	33K protein (virion morphogenesis): codon_start=1
mRNA	29331	< 12915	(C)	E2b mRNA [J. Biol. Chem. 257, 13475-13491 (1982)] [Split]
pre-msg	< 12915	16352	major late mRNA L1 (alt.) [J. Mol. Biol. 149, 189-221 (1981)], [J. Virol. 48, 127-134 (1983)] [Split]	
pre-msg	< 12915	20208	major late mRNA L2 (alt.) [J. Mol. Biol. 149, 189-221 (1981)], [J. Virol. 38, 469-482 (1981)], [J. Virol. 48, 127-134 (1983)] [Split]	
pre-msg	< 12915	24682	major late mRNA L3 (alt.) [Nucleic Acids Res. 9, 1-17 (1981)], [J. Mol. Biol. 149, 189-221 (1981)], [J. Virol. 48, 127-134 (1983)] [Split]	
pre-msg	< 12915	30462	major late mRNA L4 (alt.) [J. Mol. Biol. 149, 189-221 (1981)], [J. Virol. 48, 127-134 (1983)] [Split]	
pre-msg	< 12915	35037	major late mRNA L5 (alt.) [J. Mol. Biol. 149, 189-221 (1981)], [J. Virol. 48, 127-134 (1983)] [Split]	

Nucleotide Sequence Analysis (cont.)

mRNA	< 12915	13278	major late mRNA intron (precedes 52,55K mRNA; 1st L1 mRNA) [Cell 16, 841-850 (1979)], [Cell 16, 851-861 (1979)], [J. Mol. Biol. 134, 143-158 (1979)], [J. Mol. Biol. 135, 413-433 (1979)], [Nature 292, 420-426 (1981)] [Split]
IVS	< 12915	16388	major late mRNA intron (precedes penton mRNA; 1st L2 mRNA) [J. Virol. 48, 127-134 (1983)] [Split]
IVS	< 12915	18754	major late mRNA intron (precedes pV mRNA; 2nd L2 mRNA) [J. Biol. Chem. 259, 13980-13985 (1984)] [Split]
IVS	< 12915	20238	major late mRNA intron (precedes pVI mRNA; 1st L3 mRNA) [J. Virol. 38, 469-482 (1981)] [Split]
IVS	< 12915	21040	major late mRNA intron (precedes hexon mRNA; 2nd L3 mRNA) [Proc. Natl. Acad. Sci. U.S.A. 75, 5822-5826 (1978)], [Cell 16, 841-850 (1979)] [Split]
IVS	< 12915	23888	major late mRNA intron (precedes 23K mRNA; 3rd L3 mRNA) [Nucleic Acids Res. 9, 1-17 (1981)] [Split]
IVS	< 12915	26333	major late mRNA intron (precedes 100K mRNA; 1st L4 mRNA) [Virology 128, 140-153 (1983)] [Split]
RNA	< 12915	13005	VA I RNA (alt.) [J. Biol. Chem. 252, 9043-9046 (1977)] [Split]
RNA	< 12915	13005	VA I RNA (alt.) [J. Biol. Chem. 246, 6991-7009 (1971)], [J. Biol. Chem. 252, 9047-9054 (1977)], [Proc. Natl. Acad. Sci. U.S.A. 77, 2424-2428 (1980)] [Split]
????	< 12915	13262	VA II RNA [Proc. Natl. Acad. Sci. U.S.A. 77, 3778-3782 (1980)], [Proc. Natl. Acad. Sci. U.S.A. 77, 2424-2428 (1980)] [Split]
pept	13279	14526	1 52,55K protein; codon_start=1
pept	14547	16304	1 IIIa protein (peripentonal hexon-associated protein; splice sites not sequenced); codon_start=1
signal	16331	16336	major late mRNA L1 poly-A signal (putative) 39.21%
pept	16390	18105	1 penton protein (virion component III); codon_start=1
pept	18112	18708	1 Pro-VII protein (precursor to major core protein); codon_start=1
pept	18778	19887	1 pV protein (minor core protein); codon_start=1
signal	20188	20193	major late mRNA L2 polyadenylation signal (putative) 49.94%
pept	20240	20992	1 pVI protein (hexon-associated precursor); codon_start=1
pept	21077	23983	1 hexon protein (virion component II); codon_start=1
????	< 12915	24631	23K protein (endopeptidase); codon_start=1 [Split]
signal	24657	24662	major late mRNA L3 polyadenylation signal (putative); 62.38%
pre-msg	28193	24659 (C)	E2a late mRNA (alt.) [J. Mol. Biol. 149, 189-221 (1981)]
pre-msg	28195	24659 (C)	E2a late mRNA (alt.) [Nucleic Acids Res. 12, 3503-3519 (1984)], [Unpublished (1984)]
pre-msg	29330	24659 (C)	E2a early mRNA (alt.) [J. Mol. Biol. 149,

Nucleotide Sequence Analysis (cont.)

			189-221 (1981)]
pre-msg	29331	24659 (C)	E2a early mRNA (alt.) [J. Mol. Biol. 149, 189-221 (1981)]
signal	24683	24678 (C)	E2a mRNA polyadenylation signal on comp strand (putative); 62.43%
pept	26318	24729 (C)	DBP protein (DNA binding or 72K protein); codon_start=1
IVS	26953	26328 (C)	E2a mRNA intron B [Nucleic Acids Res. 9, 4439-4457 (1981)]
pept	26347	28764 1	100K protein (hexon assembly); codon_start=1
IVS	29263	27031 (C)	E2a early mRNA intron A [Cell 18, 569-580 (1979)]
IVS	28124	27211 (C)	E2a late mRNA intron A [Virology 128, 140-153 (1983)]
IVS	28791	28992	33K-pept intron [J. Virol. 45, 251-263 (1983)]
pept	28993	> 29366 1	33K protein (virion morphogenesis)
pept	29454	30137 1	pVIII protein (hexon-associated precursor); codon_start=1
mRNA	29848	33103	E3-2 mRNA; 85.88% [Gene 22, 157-165 (1983)]
IVS	30220	30614	major late mRNA intron ('x' leader) [Gene 22, 157-165 (1983)], [J. Biol. Chem. 259, 13980-13985 (1984)]
signal	30444	30449	major late mRNA L4 polyadenylation signal; (putative) 78.48%
signal	< 12915	32676	major late mRNA intron ('y' leader) [J. Mol. Biol. 135, 413-433 (1979)], [J. Virol. 38, 469-482 (1981)], [EMBO J. 1, 249-254 (1982)], [Gene 22, 157-165 (1983)] [Split]
pept	31051	31530 1	E3 19K protein (glycosylated membrane protein); codon_start=1
pept	31707	32012 1	E3 11.6K protein; codon_start=1
signal	32008	32013	E3-1 mRNA polyadenylation signal (putative); 82.69%
IVS	32822	33268	major late mRNA intron ('z' leader) [Proc. Natl. Acad. Sci. U.S.A. 75, 5822-5826 (1978)], [Cell 16, 841-850 (1979)], [EMBO J. 1, 249-254 (1982)], [Gene 22, 157-165 (1983)]
signal	33081	33086	E3-2 mRNA polyadenylation signal; 85.82% (putative)
????	< 12915	35017	fiber protein (virion component IV); codon_start=1 [Split]
signal	35013	35018	major late mRNA L5 polyadenylation signal; (putative) 91.19%
pre-msg	35054	> 35041 (C)	E4 mRNA [Nucleic Acids Res. 9, 1675-1689 (1981)], [J. Mol. Biol. 149, 189-221 (1981)], [Nucleic Acids Res. 12, 3503-3519 (1984)], [Unpublished (1984)] [Split]
frag	1	12914	1 to 12914 of pAd2/PGK-CFTR
DNA	1	> 356	1 to 357 Ad2
rpt	1	> 103	inverted terminal repetition; 0.28% [Biochem. Biophys. Res. Commun. 87, 671-678 (1979)], [J. Mol. Biol. 128, 577-594 (1979)]
	< 10	103	inverted terminal repetition; 0.28% [Biochem. Biophys. Res. Commun. 87, 671-678 (1979)], [J. Mol. Biol. 128, 577-594 (1979)] [Split]
frag	357	379	linker segment
frag	915	> 923	polylinker cloning sites [Split]

-84-

Nucleotide Sequence Analysis (cont.)

DNA	< 924 > 954	polylinker cloning sites [Split]
signal	< 5567 > 12914	3328 to 10685 of Ad2 [Split]
frag	< 955 > 958	pgk promoter
	< 5501 5522	polylinker cloning sites [Split]
signal	5523 5555	polylinker cloning sites [Split]
frag	5555 > 5560	syn. BGH poly A
	< 5564 5567	linker [Split]
frag	959 5500	linker [Split]
revision	2868 2868	920 to 5461 of pCMV-CFTR-936C
modified	1814 1814	mistake in published sequence of Riordan et al. C not A is correct = N to H a.a. change
site	< 959 975	936 T to C mutation to inactivate cryptic bacterial promoter. Silent amino acid change
site	976 990	polylinker segment from pCMV-CFTR-936C (Rc/CMV-Invitrogen SpeI-BstXI) [Split]
site	991 1001	linker segment from pCMV-CFTR-936C. Originally Sall/BstXI adaptor oligo 1499DS
mRNA	1001 > 5500	linker segment from pCMV-CFTR-936C.
pept	1011 > 5453	Originally from pMT-CFTR construction oligo 1247 RG -Sal I to Aval sites.
		123 to 4622 of HUMCFTR
		1 cystic fibrosis transmembrane conductance regulator; codon_start=1
BASE COUNT	8597 A 10000 C	9786 G 7952 T 0 OTHER
ORIGIN	7	

Ad2-ORF6/P Length: 36335 Sep 16, 1993 - 08:13 PM Check: 1664 ..

```

1 CATCATCAAT AATATACTT ATTTTGGATT GAAGCCAATA TGATAATGAG GGGGTGGAGT
61 TTGTGACGTG GCGCGGGGCG TGGGAACGGG GCGGGTGAAG TAGTAGTGTG GCGGAAGTGT
121 GATGTTGCAA GTGTGGGGGA ACACATGTAA GCGCCGGATG TGGTAAAGT GACGTTTTTG
181 GTGTGCGCCG GTGTATACGG GAAGTGACAA TTTTGGGCG GTTTTAGGCG GATGTTGTAG
241 TAAATTTGGG CGTAACCAAG TAATGTTTGG CCATTTTGGC GGGAAACTG AATAACAGGA
301 AGTGAAATCT GAATAATTCT GTGTTACTCA TAGCGGTAA TATTTGTCTA GGGCGGCTCG
361 AGTTCGACGG TCTATCGATA AGCTTGATAT CGAATTCGGG GGTGGGGTTC GCGCTTTTTC
421 CAAGGCAGCC CTGGGTTTGC GCAGGGAGCG GGCTGCTCTG GCGGTGGTTC CCGGAAACGC
481 AGCGGCGCCG ACCCTGGGTC TCGCACATTC TTCACGTCCG TTCCGAGCGT CACCCGGATC
541 TTCGCGGCTA CCCTTGTGGG CCCCCCGGCG ACGCTTCCTC GTCCGCCCCCT AAGTCGGGAA
601 GGTTCCTTGC GGTTCGCGGC GTGCCGAGCG TGACAAACGG AAGCCGACG TCTACTAGT
661 ACCCTCGCAG ACGGACAGCG CCAGGAGACA ATGGCAGCG GCGGACCGG ATGGGCTGTG
721 GCCAATAGCG GCTGCTCAGC AGGCGCGGCC GAGAGCAGCG GCGGGAAGG GCGGTCGCG
781 GAGGOGGGGT GTGGGGGGGT CTGTTCTGCG CCGCGCGGTG TTCCGCATTC
841 TGCAAGCCTC CGGAGCGCAC GTGGCAGTC GGCTCCCTCG TTGACCGAAT CACCGACCTC
901 TCTCCCCAGG ATCCACTAGT ATTAAATCGT ACGCCTAGTA TTTAAATCGT ACGCCTAGTA
961 ACGGCGGCCA GTGTGCTGCA GATATCAAAG TCGACGGTAC CGGAGAGACC ATGCAGAGGT
1021 CGCCTCTGGA AAAGGCCAGC GTTGTCTCCA AACTTTTTTT CAGCTGGACC AGACCAATTT
1081 TGAGGAAAGG ATACAGACAG CGCCTGGAAT TGTCAGACAT ATACCAAATC CCTTCTGTTG
1141 ATTCTGCTGA CAATCTATCT TTTAAATTCG AAAGAGAATG GGATAGAGAG CTGGCTTCAA
1201 AGAAAAATCC TAAACTCATT AATGCCCTTC GGCGATGTTT TTTCTGGAGA TTTATGTTCT
1261 ATGGAATCTT TTTATATTTA GGGGAAGTCA CCAAAGCAGT ACAGCCTCTC TTACTGGGAA
1321 GAATCATAGC TTCTATGAC CCGATAACA AGGAGGAACG CTCTATCGCG ATTTATCTAG
1381 GCATAGGCTT ATGCCTTCTC TTTATTGTGA GGACACTGCT CCTACACCCA GCCATTTTTG
1441 GCCTTCATCA CATTGGAATG CAGATGAGAA TAGCTATGTT TAGTTTGATT TATAAGAAGA
1501 CTTTAAAGCT GTCAAGCCGT GTTCTAGATA AAATAAGTAT TGGACAACCT GTTAGTCTCC
1561 TTTCCAACAA CCTGAACAAA TTTGATGAAG GACTTGCAAT GGCACATTTT GTGTGGATCG
1621 CTCCTTTGCA AGTGGCACTC CTCATGGGGC TAATCTGGGA GTTGTTACAG GCGTCTGCCT
1681 TCTGTGGACT TGGTTTCTTG ATAGTCTTTC CCGTTTTTCA GGCTGGGCTA GGGAGAATGA
1741 TGATGAAGTA CAGAGATCAG AGAGCTGGGA AGATCAGTGA AAGACTTGTG ATTACCTCAG
1801 AAATGATTGA AAACATCCAA TCTGTTAAGG CATACTGCTG GGAAGAAGCA ATGGAAAAAA

```

Nucleotide Sequence Analysis (cont.)

1861	TGATTGAAAA	CTTAAGACAA	ACAGAACTGA	AAC TGACTCG	GAAGGCAGCC	TATGTGAGAT
1921	ACTTCAATAG	CTCAGCCTTC	TTCTTCTCAG	GGTTCCTTGT	GGTGTTTTTTA	TCTGTGCTTC
1981	CCTATGCACT	AATCAAAGGA	ATCATCTCC	GGAAAAATATT	CACCACCATC	TCATTCTGCA
2041	TTGTTCTGCG	CATGGCGGTC	ACTGGGCAAT	TTCCCTGGGC	TGTACAAACA	TGGTATGACT
2101	CTCTTGGAGC	AATAAACAAA	ATACAGGATT	TCTTACAAAA	GCAAGAATAT	AAGACATTGG
2161	AATATAACTT	AACGACTACA	GAAGTAGTGA	TGGAGAATGT	AACAGCCCTC	TGGGAGGAGG
2221	GATTTGCGGA	ATTATTTGAG	AAAGCAAAAC	AAAACAATAA	CAATAGAAAA	ACTTCTAATG
2281	GTGATGACAG	CCTCTTCTTC	AGTAATTTCT	CACCTCTTGG	TACTCCTGTC	CTGAAAGATA
2341	TTAATTTCAA	GATAGAAAGA	GGACAGTTGT	TGGCGGTTGC	TGGATCCACT	GGAGCAGGCA
2401	AGACTTCACT	TCTAATGATG	ATTATGGGAG	AAC TGAGGCC	TTCAGAGGGT	AAAATTAAGC
2461	ACAGTGGAG	AATTTTCATC	TGTTCTCAGT	TTTCTCTGGAT	TATGCTGGC	ACCATTAAAG
2521	AAAATATCAT	CTTTGGTGT	TOCTATGATG	AATATAGATA	CAGAAGCGTC	ATCAAAGCAT
2581	GCCAAC TAGA	AGAGGACATC	TCCAAGTTTG	CAGAGAAAGA	CAATATAGTT	CTTGGAGAAG
2641	GTGGAATCAC	ACTGAGTGGA	GGTCAACGAG	CAAGAATTTT	TTTAGCAAGA	GCAGTATACA
2701	AAGATGCTGA	TTTGATTTTA	TTAGACTCTC	CTTTTGGATA	CCTAGATGTT	TTAACAGAAA
2761	AAGAAATATT	TGAAAGCTGT	GTCTGTAAAC	TGATGGCTAA	CAAACTAGG	ATTTTGGTCA
2821	CTTCTAAAT	GGAACATTTA	AAGAAAGCTG	ACAAAATATT	AATTTTGCAT	GAAGGTAGCA
2881	GCTATTTTTA	TGGGACATTT	TCAGAACTCC	AAAATCTACA	GCCAGACTTT	AGCTCAAAAC
2941	TCATGGGATG	TGATTCTTTC	GACCAATTTA	GTGCAGAAAG	AAGAAATTC	ATCCTAACTG
3001	AGACCTTACA	COGTTTCTCA	TTAGAAGGAG	ATGCTCCTGT	CTCCTGGACA	GAAACAAAAA
3061	AACAATCTTT	TAAACAGACT	GGAGAGTTTG	GGGAAAAAAG	GAAGAATTCT	ATTCTCAATC
3121	CAATCAACTC	TATACGAAAA	TTTTCCATTG	TGCAAAAGAC	TCCCTTACAA	TCCATTTGGCA
3181	TCGAAGAGGA	TTCTGATGAG	CCTTTAGAGA	GAAGGCTGTC	CTTAGTACCA	GATTCGAGC
3241	AGGGAGAGGC	GATACTGCCT	CGCATCAGCG	TGATCAGCAC	TGGCCCCACG	CTTCAGGCAC
3301	GAAGGAGGCA	GTCTGTCTCG	AACCTGATGA	CACACTCAGT	TAACCAAGGT	CAGAACATTC
3361	ACOGAAAGAC	AACAGCATCC	ACACGAAAAG	TGTCACTGGC	CCCTCAGGCA	AAC TTGACTG
3421	AAC TGATAT	ATATTCAAGA	AGGTTATCTC	AAGAAACTGG	CTTGGAAATA	AGTGAAGAAA
3481	TTAOGAAGA	AGACTTAAAG	GAGTGCCCTT	TTGATGATAT	GGAGAGCATA	CCAGCAGTGA
3541	CTACATGGAA	CACATACCTT	CGATATATTA	CTGTCCACAA	GAGCTTAATT	TTTGTGCTAA
3601	TTTGGTGCCT	AGTAATTTTT	CTGGCAGAGG	TGGCTGCTTC	TTTGGTTGTG	CTGTGGCTCC
3661	TTGGAAACAC	TCCTCTTCAA	GACAAAGGGA	ATAGTACTCA	TAGTAGAAAT	CAGAGCTATG
3721	CAGTGATTAT	CACCAGCACC	AGTTCTGATT	ATGTGTTTTA	CATTTACGTG	GGAGTAGCCG
3781	ACACTTTGCT	TGCTATGGGA	TTCTTCAGAG	GTCTACCAC	GGTGCATACT	CTAATCACAG
3841	TGTCGAAAAT	TTTACACCAC	AAAATGTTAC	ATTCTGTTCT	TCAAGCACCT	ATGTCAACCC
3901	TCAACACGTT	GAAAGCAGGT	GGGATTCTTA	ATAGATTCTC	CAAAGATATA	GCAATTTTGG
3961	ATGACCTTCT	GCCTCTTACC	ATATTTGACT	TCATCCAGTT	GTTATTAAAT	GTGATTGGAG
4021	CTATAGCAGT	TGTCGCAGTT	TTACAACCC	ACATCTTTGT	TGCAACAGTG	CCAGTGATAG
4081	TGGCTTTTAT	TATGTTGAGA	GCATATTTCC	TCCAAACCTC	ACAGCAACTC	AAACAAC TGG
4141	AATCTGAAGG	CAGGAGTCCA	ATTTTCACTC	ATCTTGTTAC	AAGCTTAAAA	GGACTATGGA
4201	CACTTCTGTC	CTTCGGACGG	CAGCCTTACT	TTGAAACTCT	GTTCCACAAA	GCTCTGAATT
4261	TACATACTGC	CAACTGGTTC	TTGTACCTGT	CAACACTGCG	CTGGTTCCAA	ATGAGAATAG
4321	AAATGATTTT	TGTCATCTTC	TTTCATGCTG	TTACCTTCAT	TTCCATTTTA	ACAACAGGAG
4381	AAGGAGAAGG	AAGAGTTGGT	ATTATCTCTG	CTTTAGCCAT	GAATATCATG	AGTACATTGC
4441	AGTGGGCTGT	AAACTCCAGC	ATAGATGTGG	ATAGCTTGAT	GCGATCTGTG	AGCCGAGTCT
4501	TTAAGTTTAT	TGACATGCCA	ACAGAAGGTA	AACCTACCAA	GTCAACCAAA	CCATACAAGA
4561	ATGGCCAACT	CTCGAAAGTT	ATGATTATTG	ACAAATTCACA	CGTGAAGAAA	GATGACATCT
4621	GGCCCTCAGG	GGCCCAAATG	ACTGTCAAAG	ATCTCACAGC	AAAATACACA	GAAGGTGGAA
4681	ATGCCATATT	AGAGAACATT	TCCTTCTCAA	TAAGTCTCTG	CCAGAGGGTG	GGCCTCTTGG
4741	GAAGAACTGG	ATCAGOGAAG	AGTACTTTGT	TATCAGCTTT	TTTGAGACTA	CTGAACACTG
4801	AAGGAGAAAT	CCAGATCGAT	GGTGTGTCTT	GGGATTCAAT	AAGTTTGCAA	CAGTTGGAGG
4861	AAGCCTTTGG	AGTGATACCA	CAGAAAGTAT	TTATTTTTC	TGGAACATTT	AGAAAAAACT
4921	TGGATCCCTA	TGAACAGTGG	AGTGATCAAG	AAATATGGAA	AGTTGCAGAT	GAGGTTGGGC
4981	TCAGATCTGT	GATAGAACAG	TTTCTCTGGA	AGCTTGACTT	TGTCCTTGTG	GATGGGGGCT
5041	GTGTCCTAAG	CCATGGCCAC	AAGCAGTTGA	TGTGCTTGGC	TAGATCTGTT	CTCAGTAAGG
5101	CGAAGATCTT	GCTGCTTGAT	GAACCCAGTG	CTCATTTGGA	TCCAGTAACA	TACCAATAAA
5161	TTAGAAGAAC	TCTAAAACAA	GCATTTGCTG	ATTGCACAGT	AATTCTCTGT	GAACACAGGA
5221	TAGAAGCAAT	GCTGGAATGC	CAACAATTTT	TGCTCATAGA	AGAGAACAAA	GTGCGGCAGT

Nucleotide Sequence Analysis (cont.)

5281	ACGATTCCAT	CCAGAACTG	CTGAACGAGA	GGAGCCTCTT	CCGGCAAGCC	ATCAGCCCTT
5341	CCGACAGGGT	GAAGCTCTTT	CCOCACOGGA	ACTCAAGCAA	GTGCAAGTCT	AAGCCCCAGA
5401	TTGCTGCTCT	GAAAGAGGAG	ACAGAAGAAG	AGGTGCAAGA	TACAAGGCTT	TAGAGAGCAG
5461	CATAAATGTT	GACATGGGAC	ATTTGCTCAT	GGAAATTGGAG	AAATCGTACG	CTTAGGACGC
5521	GTAATAAAAT	GAGGAAATTG	CATCGCATTG	TCTGAOCOGT	TAOCGGGGAA	GGTGTCTAGG
5581	TACGATGAGA	CCCGCACCAG	GTGCAGACCC	TGOGAGTGTG	GCGGTAAACA	TATTAGGAAC
5641	CAGCCTGTGA	TGCTGGATGT	GACCGAGGAG	CTGAGGCCCG	ATCACTTGGT	GCTGGCCTGC
5701	ACCCGCGCTG	AGTTTGGCTC	TAGCGATGAA	GATACAGATT	GAGGTACTGA	AATGTGTGGG
5761	CGTGGCTTAA	GGGTGGGAAA	GAATATATAA	GGTGGGGGTC	TCATGTAGTT	TTGTATCTGT
5821	TTTGCAGCAG	CCGCGCCCAT	GAGOGCCAAC	TCGTTTGATG	GAAGCATTGT	GAGCTCATAT
5881	TTGACAACGC	GCATGCCCCC	ATOGGCOGGG	GTGCGTCAGA	ATGTGATGGG	CTCCAGCATT
5941	GATGCTCGCC	CCGTCCCTGCC	CGCAAACCTT	ACTACCTTGA	CCTACGAGAC	CGTGTCTGGA
6001	ACGCOGTGG	AGACTGCAGC	CTCOGCOGCC	GCTTCAGCCG	CTGCAGCCAC	CGCCCGCGGG
6061	ATTGTGACTG	ACTTTGCTTT	CCTGAGCCCG	CTTGCAAGCA	GTGCAGCTTC	CCGTTTCATCC
6121	GCCCGCGATG	ACAAGTTGAC	GGCTCTTTTG	GCACAATTGG	ATTCTTTGAC	CCGGGAACCT
6181	AATGTCGTTT	CTCAGCAGCT	GTTGGATCTG	CGCCAGCAGG	TTTCTGCCCT	GAAGGCTTCC
6241	TCCCTCCCA	ATGCGGTTTA	AAACATAAAT	AAAAACCAGA	CTCTGTTTGG	ATTTTGATCA
6301	AGCAAAGTGC	TTGCTGTCTT	TATTTAGGGG	TTTTGCGCGC	GCGGTAGGCC	CGCGACCAGC
6361	GGTCTCGGTC	GTGAGGGGTC	CTGTGTATTT	TTTCCAGGAC	GTGGTAAAGG	TGACTCTGGA
6421	TGTTTCAGATA	CATGGGCATA	AGCCCGTCTC	TGGGGTGGAG	GTAGCACCAC	TGCAGAGCTT
6481	CATGCTGCGG	GGTGGTGTG	TAGATGATCC	AGTCGTAGCA	GGAGOGCTGG	GCGTGGTGCC
6541	TAAAAATGTC	TTTCAGTAGC	AAGCTGATTG	CCAGGGGCAG	GCCCTTGGTG	TAGCTGTTTA
6601	CAAAGCGGTT	AAGCTGGGAT	GGGTGCATAC	GTGGGGATAT	GAGATGCATC	TTGGACTGTA
6661	TTTTTAGGTT	GGCTATGTTT	CCAGCCATAT	CCCTCCGGGG	ATTCATGTTG	TGCAGAACCA
6721	CCAGCACAGT	GTATCCGGTG	CACCTGGGAA	ATTTGTTCATG	TAGCTTAGAA	GGAAATGOGT
6781	GGAAGAACTT	GGAGACGCCC	TTGTGACCTC	CGAGATTTTC	CATGCATTCC	TCCATAATGA
6841	TGGCAATGGG	CCCACGGGCG	GCGGCCCTGG	OGAAGATATT	TCTGGGATCA	CTAACCTCAT
6901	AGTTGTGTTT	CAGGATGAGA	TCGTTCATAGG	CCATTTTTAC	AAAGCGCGGG	CGGAGGGTGC
6961	CAGACTGCGG	TATAATGGTT	CCATCCGGCC	CAGGGGCGTA	GTTACCCCTCA	CAGATTTGCA
7021	TTTCCACCGC	TTTGAGTTCA	GATGGGGGGA	TCATGTCTAC	CTCGCGGGCG	ATGAAGAAAA
7081	CCGTTTCCGG	GGTAGGGGAG	ATCAGCTGGG	AAGAAAGCAG	GTTCCTGAGC	GTTCCGCACT
7141	TACCGCAGCC	GGTGGGCCCC	TAAATCACAC	CTATTACCGG	CTGCAACTGG	TAGTTAAGAG
7201	AGCTGCAGCT	GCCGTCATCC	CTGAGCAGGG	GGGCCACTTC	GTTAAGCATG	TCCCTGACTT
7261	GCATGCTTTC	CCTGACCATA	TGCGCCAGAA	GGCGCTCGCC	GCCCAGCGAT	ACCAGTTCTT
7321	GCAAGGAAGC	AAAGTTTTTC	AACGGTTTGA	GGCCGTCCGC	CGTAGGCATG	CTTTTGAGCG
7381	TTTGACCAAG	CAGTTCCAGG	CGGTCCCAAC	GCTCGGTCAC	GTGCTCTACG	GCATCTCGAT
7441	CCAGCATATC	TCCTCGTTTT	GCGGGTTGGG	GCGGCTTTCC	CTGTACGGCA	GTAGTCCGGT
7501	CTCGTCCAGA	CGGGCCAGGG	TCATGTCTTT	CCACGGGCGC	AGGGTCCCTG	TCAGCGTAGT
7561	CTGGGTCACG	GTGAAGGGGT	GCGCTCCGGG	CTGCGCGCTG	GCCAGGGTGC	GCTTGAGGCT
7621	GGTCTCTGCT	GTGCTGAAGC	GCTGCCGGTC	TTCCGCCCTG	GCGTCCGGCA	GGTAGCATTT
7681	GACCATGGTG	TCATAGTCCA	GCCCCCTCCG	GGCGTGGCCC	TTGGCGGCGA	GCTTGCCCTT
7741	GGAGGAGGCG	CCGCACGAGG	GGCAGTGCAG	ACTTTTAAAG	GCGTAGAGCT	TGGGCGCGAG
7801	AAATACCGAT	TCCGGGGAGT	AGGCATCCGC	CCCGCAGGCC	CCGCAGACGG	TCTCGCATTC
7861	CACGAGCCAG	GTGAGCTCTG	GCGGTTCCGG	GTCAAAAACC	AGGTTTCCCC	CATGCTTTTT
7921	GATGCGTTTC	TTACCTCTGG	TTTCCATGAG	CCGGTGTCCA	CGCTCGGTGA	CGAAAAGGCT
7981	GTCCGTGTCC	CCGTATACAG	ACTTGAGAGG	CCTGTCTCTG	AGCGGTGTTT	CGCGGTCTCT
8041	CTCGTATAGA	AACTCGGACC	ACTCTGAGAC	GAAGGCTCGC	GTCCAGGCCA	GCACGAAGGA
8101	GGCTAAGTGG	GAGGGGTAGC	GGTGGTTGTC	CACTAGGGGG	TCCACTCGCT	CCAGGGTGTG
8161	AAGACACATG	TGCCCCCTCT	CGGCATCAAG	GAAGGTGATT	GGTTTATAGG	TGTAGGCCAC
8221	GTGACCGGGT	GTTCCTGAAG	GGGGGCTATA	AAAGGGGGTG	GGGGCGGGTT	CGTCTCTACT
8281	CTCTTCCGCA	TGCTGTCTGT	CGAGGGCCAG	CTGTGGGGT	GAGTACTCCC	TCTCAAAAGC
8341	GGGCATGACT	TCTGCGCTAA	GATTGTTCAGT	TTCCAAAACC	GAGGAGGATT	TGATATTAC
8401	CTGGCCCCGG	GTGATGCCTT	TGAGGTGGGC	CGCGTCCATC	TGGTCAGAAA	AGACAATCTT
8461	TTTGTGTGTA	AGCTTGGTGG	CAAACGACCC	GTAGAGGGCG	TTGGACAGCA	ACTTGGCGAT
8521	GGAGCGCAGG	GTTTGGTTTT	TGTCGCGATC	GGCGCGCTCC	TTGGCCGCGA	TGTTTAGCTG
8581	CACGTATTCC	CGCGCAACGC	ACCGCCATT	GGGAAAGACG	GTGGTGCGCT	CGTCCGGGCAC
8641	CAGGTGCACG	CGCCAACCGC	GGTTGTGCAG	GGTGACAAGG	TCAACGCTGG	TGGCTACCTC

Nucleotide Sequence Analysis (cont.)

8701	TCCGCGTAGG	CGCTCGTTGG	TCCAGCAGAG	GCGGCGGCCC	TTGCGCGAAC	AGAATGGCGG
8761	TAGTGGGTCT	AGCTGCGTCT	CGTCCGGGGG	GTCTGCGTCC	ACGGTAAAGA	CCCCGGGCGAG
8821	CAGGCGGCGG	TOGAAGTAGT	CTATCTTGCA	TCCTTGCAAG	TCTAGOGCCT	GCTGCCATGC
8881	GCGGCGGCGA	AGCGGCGGCT	CGTATGGGTT	GAGTGGGGGA	CCCCATGGCA	TGGGGTGGGT
8941	GAGCGCGGAG	GCGTACATGC	CGCAAATGTC	GTAAACGTAG	AGGGGCTCTC	TGAGTATTCC
9001	AAGATATGTA	GGGTAGCATC	TTCCACGCGG	GATGCTGGCG	CGCACGTAAT	CGTATAGTTC
9061	GTGCGAGGGA	GCGAGGAGGT	CGGGACCGAG	GTGCTAOGG	GCGGGCTGCT	CTGCTCGGAA
9121	GACTATCTGC	CTGAAGATGG	CATGTGAGTT	GGATGATATG	GTGAGCGCT	GGAAGACGTT
9181	GAAGCTGGCG	TCTGTGAGAC	CTACCGCGTC	ACGCAOGAAG	GAGGCGTAGG	AGTOGCGCAG
9241	CTGTGTGACC	AGCTCGGCGG	TGACCTGCAC	GTCTAGGGCG	CAGTAGTCCA	GGGTTTCCCT
9301	GATGATGTCA	TACTTATCCT	GTCCCTTTTT	TTTCCACAGC	TCGCGGTTGA	GGACAAACTC
9361	TTCCGGGTCT	TTCCAGTACT	CTTGGATCGG	AAACCCGTCG	GCCTCCGAAC	GGTAAGAGCC
9421	TAGCATGTAG	AACTGGTTGA	CGGCCTGGTA	GCGCGAGCAT	CCCTTTTCTA	CGGGTAGCGC
9481	GTATGCGTGC	GCGGCCTTCC	GGAGCGAGGT	GTGGGTGAGC	GCAAAGGTGT	CCCTAACCAT
9541	GACTTTGAGG	TACTGGTATT	TGAAGTCAGT	GTOGTGCGAT	CCGCCCTGCT	CCCAGAGCAA
9601	AAAGTCCGTC	CGCTTTTTGG	AACGCGGGTT	TGGCAGGGCG	AAGGTGACAT	CGTTGAAAAG
9661	TATCTTTCCC	GCGCGAGGCA	TAAAGTTGCG	TGTGATGOGG	AAGGGTCCCG	GCACCTCGGA
9721	ACGGTGTGTA	ATTACCTGGG	CGGCGAGCAC	GATCTCGTCG	AAGCCGTTGA	TGTTGTGGCC
9781	CAGGATGTAA	AGTTCCAAGA	AGCGCGGGGT	GCCCTTGATG	GAGGGCAATT	TTTTAAGTTC
9841	CTGTAGGTG	AGCTCCTCAG	GGGAGCTGAG	CCCGTGTCT	GACAGGGCCC	AGTCTGCAAG
9901	ATGAGCGTTG	GAAGCGAOGA	ATGAGCTCCA	CAGGTCAOGG	GCCATTAGCA	TTTGCAGGTG
9961	GTCGCGAAAG	GTCCATAACT	GGCGACCTAT	GGCCATTTTT	TCTGGGGTGA	TGCAGTAGAA
10021	GGTAAGCGGG	TCTTGTTCCT	AGCGGTCCCA	TCCAAGGTCC	ACGGCTAGGT	CTCGCGCGGC
10081	GGTCACCAGA	GGCTCATCTC	CGCCGAACCT	CATAACCAGC	ATGAAGGGCA	CGAGCTGCTT
10141	CCCAAAGGCC	CCCATCCAAG	TATAGGTCTC	TACATCGTAG	GTGACAAAGA	GACGCTGGGT
10201	GCGAGGATGC	GAGCCGATCG	GGAAGAAGTC	GATCTCCCGC	CACCACTTGG	AGGAGTGGCT
10261	GTTGATGTGG	TGAAAGTAGA	AGTCCCTGCG	ACGGGCCGAA	CACTCGTGCT	GGCTTTTGTA
10321	AAAACGTGCG	CAGTACTGGC	ACCGGTGCAC	GGGCTGTACA	TCCTGCAOGA	GGTTGACCTG
10381	ACGACCGGCG	ACAAGGAAGC	AGAGTGGGAA	TTTGAGCCCC	TCGCCTGGCG	GGTTTGGCTG
10441	GTGGTCTTCT	ACTTCGGCTG	CTTGTCTCTG	ACCGTCTGCG	TGCTCGAGGG	GAGTTATGGT
10501	GGATCGGACC	ACCACGCCGC	GCGAGCCCCA	AGTCCAGATG	TCCGCGCGCG	GCGGTGCGAG
10561	CTTGATGACA	ACATCGCGCA	GATGGGAGCT	GTCCATGGTC	TGGAGCTCCC	GCGGCGACAG
10621	GTCAGGCGGG	AGCTCCTGCA	GGTTTACCTC	GCATAGCCGG	GTGAGGGGCG	GGGCTAGGTC
10681	CAGGTGATAC	CTGATTTCCA	GGGGCTGGTT	GGTGGCGGCG	TCGATGACTT	GCAAGAGCCC
10741	GCATCCCCGC	GGCGCGACTA	CGGTACCGCG	CGGCGGGGCG	TGGGCGCGCG	TGGTGTCTCT
10801	GGATGATGCA	TCTAAAAGCG	GTGACGCGGG	CGGGCCCCCG	GAGGTAGGGG	GGGCTCGGGA
10861	CCCGCCGGGA	GAGGGGGCAG	GGGCACGTCG	GCGCGCGCGG	CGGGCAGGAG	CTGGTGCTGC
10921	GCGCGGAGGT	TGCTGGCGAA	CGCGACGACG	CGGCGGTTGA	TCTCCTGAAT	CTGGCGCCTC
10981	TGCGTGAAGA	CGACGGGGCC	GGTGAGCTTG	AACCTGAAAG	AGAGTTGAC	AGAATCAATT
11041	TCGGTGTGCT	TGACGGCGGC	CTGGCGCAA	ATCTCCTGCA	CGTCTCCTGA	GTTGTCTTGA
11101	TAGGCGATTT	CGGCCATGAA	CTGCTGATC	TCTTCTCTCT	GGAGATCTCC	GCGTCCGGCT
11161	CGCTCCACGG	TGGCGGCGAG	GTOGTTGGAG	ATGCGGGCCA	TGAGCTGCGA	GAAGGCGTTG
11221	AGGCCTCCCT	CGTTCCAGAC	GCGGCTGTAG	ACCACGCCCC	CTTCGGCATC	GCGGGCGCGC
11281	ATGACCACCT	GCGCGAGATT	GAGCTCCACG	TGCGGGGCGA	AGACGGCGTA	GTTTCGCAGG
11341	CGCTGAAAGA	GGTAGTTGAG	GGTGGTGGCG	GTGTGTTCTG	CCACGAAGAA	GTACATAACC
11401	CAGCGTCGCA	ACGTGGATTTC	GTTGATATCC	CCCAAGGCCT	CAAGGCGCTC	CATGGCCTCG
11461	TAGAAGTCCA	CGGCGAAGTT	GAAAACTGG	GAGTTGCGCG	CCGACACGGT	TAACCTCTCC
11521	TCCAGAAGAC	GGATGAGCTC	GGCGACAGTG	TCGCGCACCT	CGCGCTCAAA	GGCTACAGGG
11581	GCCTCTTCTT	CTTCAATCTC	CTCTTCCATA	AGGGCCTCCC	CTTCTTCTTC	TTCTTCTGGC
11641	GCGCGTGGGG	GAGGGGGGAC	ACGGCGGCGA	CGACGGGCGA	CCGGGAGGCG	GTCGACAAAG
11701	CGCTCGATCA	TCTCCCCGCG	GCGACGGCGC	ATGGTCTCGG	TGACGGGCGG	GCCGTCTCTG
11761	CGGGGGCGCA	GTTGGAAGAC	GCGGCCCCGC	ATGTCCCGGT	TATGGGTTGG	CGGGGGGCTG
11821	CCGTGCGGCA	GGGATACGGC	GCTAACCATG	CATCTCAACA	ATTGTTGTGT	AGGTACTCCG
11881	CCACCGAGGG	ACCTGAGCGA	GTCCGCATCG	ACCGGATCGG	AAAACCTCTC	GAGAAAGGCG
11941	TCTAACCAGT	CACAGTCGCA	AGGTAGGCTG	AGCACCGTGG	CGGGCGGCGG	GCGGTGGCGG
12001	TGCGGGTTGT	TTCTGGCGGA	GGTGCTGCTG	ATGATGTAAT	TAAAGTAGGC	GGTCTTGAGA
12061	CGGCGGATGG	TCGACAGAAG	CACCATGTCC	TTGGGTCCGG	CCTGCTGAAT	GCGCAGGCGG

Nucleotide Sequence Analysis (cont.)

12121	TGGCCATGC	CCCAGGCTTC	GTTTGTGACAT	CGGCGCAGGT	CTTTGTAGTA	GTCTTGCAATG
12181	AGCCTTTCTA	CCGGCACTTC	TTCTTCTCCT	TCCTCTTGTC	CTGCATCTCT	TGCATCTATC
12241	GCTACGGCGG	CGGCGGAGTT	TGGCGGTAGG	TGGCGCCCTC	TTCTTCCCAT	GCGTGTGACC
12301	CGGAAGCCCC	TCATCGGCTG	AAGCAGGGCC	AGGTCGGGGA	CAACGGGCTC	GGCTAATATG
12361	GCCTGCTGCA	CCTGCGTGAG	GGTAGACTGG	AAGTCATCCA	TGTCCACAAA	GCGGTGGTAT
12421	GCGCCCGTGT	TGATGGTGTA	AGTGCACTTG	GCCATAACGG	ACCACTTAAC	GGTCTGGTGA
12481	CCCGGCTGCG	AGAGCTCGGT	GTACCTGAGA	CGCGAGTAAG	CCCTTGAGTC	AAAGACGTAG
12541	TGGTTGCAAG	TCCGCACCAAG	GTACTGATAT	CCCACCAAAA	AGTGCGGCGG	CGGCTGGGCG
12601	TAGAGGGGCC	ACCGTAGGGT	GGCGGGGCTT	CGCGGGGCGA	GGTCTTCCAA	CATAAGGCGA
12661	TGATATCCGT	AGATGTACCT	GGACATCCAG	GTGATGCCGG	CGGCGGTGGT	GGAGGCGGCG
12721	GGAAAGTCGC	GGACGCGGTT	CCAGATGTTG	CGCAGCGGCA	AAAAGTGCTC	CATGGTGGGG
12781	ACGCTCTGGC	CGGTGAGGCG	TGCGCAGTCG	TTGACGCTCT	AGACCGTGCA	AAAAGAGAGC
12841	CTGTAAGCGG	GCACTCTTCC	GTGGTCTGGT	GGATAAATTC	GCAAGGGTAT	CATGGCGGAC
12901	GACCGGGGTT	CGAACCCCGG	ATCCGGCCGT	CGCGGTGAT	CCATGCGGTT	ACCGCCCGCG
12961	TGTCGAACCC	AGGTGTGCGA	CGTCAGACAA	CGGGGGAGCG	CTCCTTTTGG	CTTCCCTTCCA
13021	GGCGCGGCGG	CTGCTGCGCT	AGCTTTTTTG	GCCACTGGCC	GCGCGCGGGG	GCGCGCGTTA
13081	GGCTGGAAG	CGAAAGCATT	AAGTGCGCTCG	CTCCCTGTAG	CCGGAGGGTT	ATTTTCCAAAG
13141	GGTTGAGTCG	CAGGACCCCC	GGTTGAGTTC	TGGGGCCGGC	CGGACTGCGG	CGAACGGGGG
13201	TTTGCCCTCCC	CGTCATGCAA	GACCCCGCTT	GCAAAATTCCT	CCGGAACACAG	GGACGAGCCC
13261	CTTTTTTGCT	TTTCCAGAT	GCATCCGCTG	CTGCGGCAGA	TGCGCCCCCC	TCCTCAGCAG
13321	CGGCAAGAGC	AAGAGCAGCG	GCAGACATGC	AGGGCACCCCT	CCCTTCTCTC	TACCGCGTCA
13381	GGAGGGGCAA	CATCCGCGGC	TGACCGCGCG	GCAGATGGTG	ATTACGAACC	CCCGCGGCGC
13441	GGGGCCCGGC	ACTACCTGGA	CTTGAGGAG	GGCGAGGGCC	TGGGCGGGCT	AGGAGCGCCC
13501	TCTCCTGAGC	GACACCCAAG	GGTGCACTCG	AAGCGTGACA	CGCGGAGGGC	GTACGTGCCG
13561	CGGCAGAACC	TGTTTTCGCG	CGCGGAGGGA	GAGGAGCCCG	AGGAGATGCG	GGATCGAAG
13621	TTCCACGCAG	GGCGCGAGTT	GCGGCATGGC	CTGAACCGCG	AGCGGTGCT	GCGGAGGAG
13681	GACTTTGAGC	CCGACGCGCG	GACCGGGATT	AGTCCCGGCG	GCGCACAGCT	GCGGCGCGCC
13741	GACCTGGTAA	CCGCGTACGA	GCAGACGGTG	AACCAAGGAG	TTAACTTTCA	AAAAAGCTTT
13801	AACAACCAAG	TGCGCACGCT	TGTGGCGCGC	GACGAGGTGG	CTATAGGACT	GATGCATCTG
13861	TGGGCAAGT	TAAGCGCGCT	GGAGCAAAAC	CCAAATAGCA	AGCGGCTCAT	GGCGCAGCTG
13921	TTCCCTTATG	TGCAGCACAG	CAGGACAAAC	GAGGCATTCA	GGGATGCGCT	GCTAAACATA
13981	GTAGAGCCCG	AGGGCGGCTG	GCTGCTCGAT	TTGATAAACA	TTCTGCAGAG	CATAGTGGTG
14041	CAGGAGCGCA	GCTTGAGCCT	GGCTGACAAG	GTGGCCGCCA	TTAACTATTTC	CATGCTCAGT
14101	CTGGGCAAGT	TTTACGCCCC	CAAGATATAC	CATACCCCTT	ACGTTCCCAT	AGACAAGGAG
14161	GTAAAGATCG	AGGGGTTCCTA	CATGCGCATG	GCGTTGAAGG	TGCTTACCTT	GAGCGACGAC
14221	CTGGGCGTTT	ATCGCAACGA	GCGCATCCAC	AAGGCCGTGA	GCGTGAGCCG	GCGGCGCGAG
14281	CTCAGCGACC	GCGAGCTGAT	GCACAGCCTG	CAAAGGGGCC	TGGCTGGCAC	GGGCGACGGC
14341	GATAGAGAGG	CCGAGTCCTA	CTTTGACGCG	GGCGCTGACC	TGCGCTGGGC	CCCAAGCCGA
14401	CGCGCCCTGG	AGGCAGCTGG	GGCGGACCTT	GGGCTGGCGG	TGGCACCCGC	GCGCGCTGGC
14461	AACGTGCGCG	GCGTGGAAGA	ATATCACGAG	GACGATGAGT	ACGAGCCAGA	GGACGCGGAG
14521	TACTAAGCGG	TGATGTTTCT	GATCAGATGA	TGCAAGACGC	AACGGACCCG	GCGGTGCGGG
14581	CGGCGCTGCA	GAGCCAGCCG	TCCGGCCCTTA	ACTCCACGGA	CGACTGGCGC	CAGGTCATCG
14641	ACCGCATCAT	GTCGCTGACT	GCGCGTAACC	CTGACGCGTT	CCGGCAGCAG	CCGCAGGCCA
14701	ACCGGCTCTC	CGCAATTCTG	GAAGCGGTGG	TCCCGGCGCG	CGCAAACCCC	ACGCACGAGA
14761	AGGTGCTGGC	GATCGTAAAC	GCGCTGGCCG	AAAACAGGGC	CATCCGGCCC	GATGAGGCCG
14821	GCTTGTCTA	CGACGCGCTG	CTTCAGCGCG	TGGCTCGTTA	CAACAGCGGC	AACGTGCAGA
14881	CCAACCTGGA	CCGGCTGGTG	GGGGATGTGC	GCGAGGCCGT	GGCGCAGCGC	GCGGCGCGCG
14941	AGCAGCAGGG	CAACCTGGGC	TCCATGGTTG	CACATAAACGC	CTTCCTGAGT	ACACAGCCCG
15001	CCAACGTGCC	GCGGGGACAG	GAGGACTACA	CCAACTTTGT	GAGCGCACTG	CGGCTAATGG
15061	TGACTGAGAG	ACCGCAAAGT	GAGGTGTACC	AGTCGGGGCC	AGACTATTTT	TTCCAGACCA
15121	GTAGCAAGAG	CCTGCAGACC	GTAACCTGA	GCCAGGGCTT	CAAGAACTTG	CAGGGGCTGT
15181	GGGGGGTGCG	GGCTCCCAACA	GGCGACCGCG	CGACCGTGTC	TAGCTTGCTG	ACGCCCCAAT
15241	CGCGCCTGTT	GCTGCTGCTA	ATAGCGCCCT	TCACGGACAG	TGGCAGCGTG	TCCCGGGACA
15301	CATACCTAGG	TCACTTGCTG	ACACTGTACC	GCGAGGCCAT	AGGTGAGGCG	CATGTGGACG
15361	AGCATACTTT	CCAGGAGATT	ACAAGTGTC	GCCGCGCGCT	GGGGCAGGAG	GACACGGGCA
15421	GCCTGGAGGC	AACCTGAAC	TACCTGCTGA	CCAACCGGCG	GCAGAAGATC	CCCTCGTTGC
15481	ACAGTTTAAA	CAGCGAGGAG	GAGCGCATCT	TGCGCTATGT	GCAGCAGAGC	GTGAGCCTTA

Nucleotide Sequence Analysis (cont.)

15541	ACCTGATGCG	CGACGGGGTA	ACGCCCAGCG	TGGCGCTGGA	CATGACCGCG	CGCAACATGG
15601	AACCGGGCAT	GTATGCCTCA	AACCGGCCGT	TTATCAATCG	CCTAATGGAC	TACTTGCATC
15661	GCGCGGCCGC	CGTGAACCCC	GAGTATTTCA	CCAATGCCAT	CTTGAACCCG	CACTGGCTAC
15721	CGCCCCCTGG	TTTCTACACC	GGGGGATTTG	AGGTGCCCGA	GGGTAACGAT	GGATTCCCTCT
15781	GGGACGACAT	AGACGACAGC	GTGTTTTCCT	CGCAACCGCA	GACCCCTGCTA	GAGTTGCAAC
15841	AGCGGAGCA	GGCAGAGGGC	GCGCTGCGAA	AGGAAAGCTT	CCGCAGGCECA	AGCAGCTTGT
15901	CGATCTAGG	CGCTGCGGCC	CGCGGCTCAG	ATGCGAGTAG	CCCATTTCCTA	AGCTTGATAG
15961	GGTCTTTTAC	CAGCACTCGC	ACCACCCGCC	CGCGCTGCT	GGGCGAGGAG	GAGTACCTAA
16021	ACAACCTCGT	GCTGCAGCG	CAGCGCGAAA	AGAACCTGCC	TCCGGCATTT	CCCAACAAACG
16081	GGATAGAGAG	CCTAGTGGAC	AAGATGAGTA	GATGGAAGAC	GTATGCGCAG	GAGCACAGGG
16141	ATGTGCCCGG	CCCGCGCCCG	CCCACCCGTC	GTCAAAGGCA	CGACCGTCAG	CGGGGTCTGG
16201	TGTGGGAGGA	CGATGACTCG	GCAGACGACA	GCAGCGTCCT	GGATTTCGGA	GGGAGTGGCA
16261	ACCCGTTTGC	GCACCTTCGC	CCCAGGCTGG	GGAGAATGTT	TTAAAAAAA	AAAAAAAAG
16321	CATGATGCAA	AATAAAAAAC	TCACCAAGGC	CATGGCACCG	AGGTTGGTT	TTCTTGATT
16381	CCCCTTAGTA	TGCAGCGCGC	GGCGATGTAT	GAGGAAGGTC	CTCCTCCCTC	CTACGAGAGC
16441	GTGGTGAGCG	CGGCGCCAGT	GGCGGCGGCG	CTGGGTTCCT	CCTTCGATGC	TCCCTGGAC
16501	CGCGGTTTG	TGCCTCCGCG	GTACCTCCGG	CCTACCGGGG	GGAGAAACAG	CATCOGTTAC
16561	TCTGAGTTGG	CACCCCTATT	CGACGCCACC	CGTGTGTACC	TTGTGGACAA	CAAGTCAACG
16621	GATGTGGCAT	CCCTGAACCTA	CCAGAACGAC	CACAGCAACT	TTCTAACCCAC	GGTCATTCAA
16681	AACAATGACT	ACAGCCCGGG	GGAGGCAAGC	ACACAGACCA	TCAATCTTGA	CGACCGTTCCG
16741	CACTGGGGCG	GCGACCTGAA	AACCATCCTG	CATACCAACA	TGCCAAATGT	GAAACGAGTTC
16801	ATGTTTACCA	ATAAGTTTAA	GGCGGGGGTG	ATGGTGTGCG	GCTCGCTTAC	TAAGGACAAA
16861	CAGGTGGAGC	TGAAATATGA	GTGGGTGGAG	TTACGCTGCG	CGAGGGGCAA	CTACTCCGAG
16921	ACCATGACCA	TAGACCTTAT	GAACAACGCG	ATCGTGGAGC	ACTACTTGAA	AGTGGGCAGG
16981	CAGAACGGGG	TTCTGGAAAG	CGACATCGGG	GTAAAGTTTG	ACACCCGCAA	CTTCAGACTG
17041	GGGTTTGACC	CAGTCACTGG	TCTTGTTCATG	CCTGGGGTAT	ATACAAACGA	AGCCTTCCAT
17101	CCAGACATCA	TTTTGCTGCC	AGGATGCGGG	GTGGACTTCA	CCCACAGCCG	CCTGAGCAAC
17161	TTGTTGGGCA	TCCGCAAGCG	GCAACCTTTC	CAGGAGGGCT	TTAGGATCAC	CTACGATGAC
17221	CTGGAGGGTG	GTAACATTCC	CGCACTGTTG	GATGTGGACG	CCTACCAGGC	AAGCTTAAAA
17281	GATGACACCG	AACAGGGCGG	GGATGGCGCA	GGCGGCGGCA	ACAACAGTGG	CAGCGGCGCG
17341	GAAGAGAACT	CCAACGCGGC	AGCCGCGGCA	ATGCAGCOGG	TGGAGGACAT	GAACGATCAT
17401	GCCATTGCGG	GCGACACCTT	TGCCACACGG	GCGGAGGAGA	AGCGCGCTGA	GGCCGAGGCA
17461	GCGGCAGAAG	CTGCCGCCCC	CGCTGCGCAA	CCCGAGGTGCG	AGAAGCCTCA	GAAGAAACCG
17521	GTGATCAAAC	CCCTGCACAGA	GGACAGCAAG	AAACGCAGTT	ACAACCTAAT	AAGCAATGAC
17581	AGCACCTTCA	CCCAGTACCG	CAGCTGGTAC	CTTGCAACA	ACTACGGCGA	CCCTCAGACC
17641	GGGATCCGCT	CATGGACCCCT	CCTTGCACCT	CCTGACGTAA	CCTGCGGCTC	GGAGCAGGTC
17701	TACTGGTGGT	TGCCAGACAT	GATGCAAGAC	CCCGTGACCT	TCCGCTCCAC	GAGCCAGATC
17761	AGCAACTTTC	CGGTGGTGCG	CGCCGAGCTG	TTGCCCGTGC	ACTCCAAGAG	CTTCTACAAC
17821	GACCAGGCCG	TCTACTCCCA	GCTCATCCGC	CAGTTTACCT	CTCTGACCCA	CGTGTTCAT
17881	CGCTTTCCCG	AGAACCAGAT	TTTGGCGCGC	CCGCCAGCCC	CCACCATCAC	CACCGTCAGT
17941	GAAGACGTTT	CTGCTCTCAC	AGATCACGGG	ACGCTACOGC	TGCGCAACAG	CATCGGAGGA
18001	GTCCAGCGAG	TGACCATTAC	TGACGCCAGA	CGCCGCACCT	GCCCCCTACGT	TTACAAGGCC
18061	CTGGGCATAG	TCTCGCCGCG	CGTCTATCG	AGCCGCACTT	TTTGAGCAAA	CATGTCCATC
18121	CTTATATCGC	CCAGCAATAA	CACAGGCTGG	GGCCTGCGCT	TCCCAAGCAA	GATGTTTGCG
18181	GGGGCAAAGA	AGCGCTCCGA	CCAACACCCA	GTGCGCGTGC	GCGGGCACTA	CCGCGCGCCC
18241	TGGGGCGCGC	ACAAACGCGG	CGCACTGGG	CGCACACCG	TCGATGACCG	CATTGACGCG
18301	TGGGTGGAGG	AGGCGCGCAA	CTACACGCC	ACGCGGCCAC	CAGTGTCCAC	AGTGGACGCG
18361	GCCATTGAGA	CCGTGGTGCG	CGGAGCCCGG	CGTTATGCTA	AAATGAAGAG	ACGGCGGAGG
18421	CGCGTAGCAC	GTCGCCACCG	CCGCCGACCC	GGCACTGCGG	CCCAACGCGC	GGCGGCGGCC
18481	CTGCTTAACC	GCGCAOGTCG	CACCGGCCGA	CGGGCGGCCA	TGCGGGCCGC	TCGAAGGCTG
18541	GCCGCGGGTA	TTGTCACTGT	GCCCCCAGG	TCCAGGCGAC	GAGCGGCCGC	CGCAGCAGCC
18601	GCGGCCATTA	GTGCTATGAC	TCAGGGTTCG	AGGGGCAACG	TGTACTGGGT	GCGCGACTCG
18661	GTTAGCGGCC	TGCGCGTGCC	CGTCCGCACC	CGCCCCCGC	GCAACTAGAT	TGCAAGAAAA
18721	AACTACTTAG	ACTCGTACTG	TTGTATGTAT	CCAGCGCGCG	CGGCGCGCAA	GGAAGCTATG
18781	TCCAAGCGCA	AAATCAAAGA	AGAGATGCTC	CAGGTACATG	CGCCGGAGAT	CTATGGCCCC
18841	CCGAAGAAGG	AAGAGCAGGA	TTACAAGCCC	CQAAAGCTAA	AGCGGGTCAA	AAAGAAAAAG
18901	AAAGATGATG	ATGATGATGA	ACTTGACGAC	GAGGTGGAAC	TGCTGCACCG	AACCGCGCCC

Nucleotide Sequence Analysis (cont.)

18961	AGGCGGCGGG	TACAGTGGAA	AGGTGACGCG	GTAAGACGTG	TTTTGCGACC	CGGCAACCACC
19021	GTAAGTTTAA	CGCCCGGTGA	GCGCTCCACC	CGCACCTACA	AGCGCGTGTA	TGATGAGGTG
19081	TACGGCGAAG	AGGACCTGCT	TGAGCAGGCC	AACGAGCGCC	TGGGGAGTT	TGCGTACGGA
19141	AAGCGGCATA	AGGACATGTT	GGOGTTGCCG	CTGGACGAGG	GCAACCCAAC	ACTTAGCCTA
19201	AAGCCCGTGA	CACTGCAGCA	GGTGTGCCC	ACGCTTGAC	CGTOOGAAGA	AAAGCGGGC
19261	CTAAAGCGCG	AGTCTGGTGA	CTTGGCACCC	ACCGTGCAGC	TGATGGTACC	CAAGCGCCAG
19321	CGACTGGAAG	ATGTCTTGGG	AAAAATGACC	GTGGAGCCTG	GGCTGGAGCC	CGAGGTCCGC
19381	GTGCGGCCAA	TCAAGCAGGT	GGCACCGGGA	CTGGGCGTGC	AGACCGTGGG	CGTTCAGATA
19441	CCCAACCA	GTAAGCACTAG	TATTGCACT	GCCACAGAGG	GCATGGAGAC	ACAAACGTCC
19501	CGGTTGCTCT	CGGCGGTGGC	AGATGCCGCG	GTGCAGGCGG	CGGCTGCGGC	CGCGTCCAAA
19561	ACCTCTACGG	AGGTGCAAA	GGACCGGTGG	ATGTTTGGCG	TTTCAGCCCC	CCGGCGCCCCG
19621	CGCGGTCCA	GGAAGTACGG	CACCGCCAGC	GCACTACTGC	CCGAATATGC	CCGACATCCT
19681	TCCATCGCGC	CTACCCCGCG	CTATCGTGGC	TACACCTACC	GCCCCAGAAG	ACGAGCGACT
19741	ACCCGACGCC	GAACCACCAC	TGGAACCGCG	CGCGCGGTC	GCCGTCGCCA	GCCCGTGCTG
19801	GCCCGGATT	CGGTGCGCAG	GGTGGCTGCG	GAAGGAGGCA	GGACCGTGGT	GCTGCCAACA
19861	GCGCGCTACC	ACCCAGCAT	CGTTTAAAG	CCGGTCTTTG	TGGTTCTTGC	AGATATGGCC
19921	CTCACCTGCC	GCCTCGGTTT	CCCGGTGCCG	GGATTGCGAG	GAAGAATGCA	CCGTAGGAGG
19981	GGCATGCGCG	GCCACGGCCT	GACGGGCGGC	ATGCGTGGTG	CGCACCCACG	GCGGCGGCGC
20041	GCGTCGCAAC	GTCGATGCG	CGGCGGTATC	CTGCCCCCTC	TTATTCCACT	GATCGCGCGG
20101	GCGATTGCGG	CCGTGCCCGG	AATTGCATCC	GTGGCCTTGC	AGGCGCAGAG	ACACTGATTA
20161	AAAACAAGTT	GCATGTGGAA	AAATCAAAAT	AAAAAGTCTG	GAGTCTCACG	CTCGCTTGGT
20221	CTGTAACTA	TTTTGTAGAA	TGGAAGACAT	CAACTTTGCG	TCTCTGGCCC	CGCGACACGG
20281	CTCGCGCCCG	TTCATGGGAA	ACTGGCAAGA	TATCGGCACC	AGCAATATGA	GCGGTGGGCG
20341	CTTCAGCTGG	GGCTCGCTGT	GGAGCGGCAT	TAAAAATTTT	GGTTCCACCA	TTAAGAACTA
20401	TGGCAGCAAG	GCCTGGAACA	GCAGCACAGG	CCAGATGCTG	AGGGACAAGT	TGAAGAGCA
20461	AAATTTCCAA	CAAAAGGTGG	TAGATGGCCT	GGCCTCTGGC	ATTAGCGGGG	TGGTGGACCT
20521	GGCCAACCAG	GCAGTGCAAA	ATAAGATTAA	CAGTAAGCTT	GATCCCCGCC	CTCCCGTAGA
20581	GGAGCCTCCA	CGGCGGTGG	AGACAGTGTG	TCCAGAGGGG	CGTGGGGA	AGCGTCCGCG
20641	GCCCGACAGG	GAAGAACTC	TGGTGAACGA	AATAGATGAG	CCTCCCTCGT	ACGAGGAGGC
20701	ACTAAAGCAA	GGCCTGCCCA	CCACCGTCC	CATCGCGCCC	ATGGCTACCG	GAGTGCTGGG
20761	CCAGGACACA	CCTGTAAAGC	TGGACCTGCC	TCCCCCGCT	GACACCCAGC	AGAAACCTGT
20821	GCTGCCAGGG	CCGTCCGCGG	TTGTGTAAAC	CCGCCCTAGC	CGCGGTCC	TGCGCGTGC
20881	CGCCAGCGGT	CCGCGATCGA	TGCGGCCCGT	AGCCAGTGGC	AACTGGCAAA	GCACACTGAA
20941	CAGCATCGTG	GGTCTGGGGG	TGCAATCCCT	GAAGCGCCGA	CGATGCTTCT	AAATAGCTAA
21001	CGTGTGCTAT	GTGTCACTGA	TGCGTCCATG	TGCGCGCCAG	AGGAGCTGCT	GAGCGCGCGT
21061	GCGCCCGCTT	TCCAAGATGG	CTACCCCTTC	GATGATGCGG	CAGTGGTCTT	ACATGCACAT
21121	CTCGGGCCAG	GACGCCCTCG	AGTACCTGAG	CCCCGGGCTG	GTGCAGTTTG	CCCGCGCCAC
21181	CGAGACGTAC	TTCAGCCTGA	ATAACAAGTT	TAGAAACCCC	ACGGTGGCAC	CTACGCACGA
21241	CGTAACCACA	GACCGGTCCC	AGCGTTTGAC	GCTGCGGTTT	ATCCCTGTGG	ACCGCGAGGA
21301	TACCGCGTAC	TGCTACAAAG	CGCGGTTTAC	CCTGGCTGTG	GGTGACAACC	GTGTGCTTGA
21361	TATGGCTTCC	ACGTACTTTG	ACATCCGCGG	CGTGTGAGAC	AGGGGGCCTA	CTTTTAAGCC
21421	CTACTCCGGC	ACTGCCTACA	ACGCTCTAGC	TCCCAAGGGC	GCTCCTAACT	CCTGTGAGTG
21481	GGAACAAACC	GAAGATAGCG	GCCGGGCACT	TGCCGAGGAT	GAAGAAGAGG	AAGATGAAGA
21541	TGAAGAAGAG	GAAGAAGAAG	AGCAAAACCG	TGAGATCAG	CCTACTAAGA	AAACACATGT
21601	CTATGCCCCAG	GCTCCTTTGT	CTGGAGAAAC	AATTACAAA	AGCGGGCTAC	AAATAGGATC
21661	AGACAATGCA	GAAACACAG	CTAAACCTGT	ATAOGCAGAT	CCTTCCTATC	AACCAGAAC
21721	TCAAATTGGC	GAATCTCAGT	GGAACGAAGC	TGATGCTAAT	GCGGCAGGAG	GGAGAGTGCT
21781	TAAAAAACA	ACTCCCATGA	AACCATGCTA	TGGATCTTAT	GCCAGGCCTA	CAAATCCTTT
21841	TGGTGGTCAA	TCCGTTCTGG	TCCGGATGA	AAAAGGGGTG	CCTCTTCCAA	AGGTTGACTT
21901	GCAATTCTTC	TCAAATACTA	CCTCTTTGAA	CGACCGGCA	GGCAATGCTA	CTAAACCAAA
21961	AGTGGTTTTG	TACAGTGAAG	ATGTAAATAT	GGAAACCCCA	GACACACATC	TGCTTTACAA
22021	ACCTGGAAAA	GGTGATGAAA	ATTCTAAAGC	TATGTTGGGT	CAACAATCTA	TGCCAAACAG
22081	ACCCAATTAC	ATTGCTTTCA	GGGACAATTT	TATTGGCCCTA	ATGTATTATA	ACAGCACTGG
22141	CAACATGGGT	GTTCTTGCTG	GTCAGGCATC	GCAGCTAAAT	GCCGTGGTAG	ATTTGCAAGA
22201	CAGAAACACA	GAGCTGTCCT	ATCAACTCTT	GCTTGATTCC	ATAGGTGATA	GAACCAGATA
22261	TTTTTCTATG	TGGAATCAGG	CTGTAGACAG	CTATGATCCA	GATGTTAGAA	TCATTGAAAA
22321	CCATGGAAC	GAGGATGAAT	TGCCAAATTA	TTGTTTTCTT	CTTGGGGGTA	TTGGCGTAAC

Nucleotide Sequence Analysis (cont.)

22381	TGACACCTAT	CAAGCTATTA	AGGCTAATGG	CAATGGCTCA	GGCGATAATG	GAGATACTAC
22441	ATGGACAAAA	GATGAAACTT	TTGCAACACG	TAATGAAATA	GGAGTGGGTA	ACAACCTTGC
22501	CATGGAAATT	AACCTAAATG	CCAACCTATG	GAGAAATTTT	CTTTACTCCA	ATATTGOGCT
22561	GTACCTGCCA	GACAAGCTAA	AATACAACCC	CACCAATGTG	GAAATATCTG	ACAACCCCAA
22621	CACCTACGAC	TACATGAACA	AGCGAGTGGT	GGCTCCCGGG	CTTGTAAGCT	GCTACATTAA
22681	CCTTGGGGCG	CGCTGGTCTC	TGGACTACAT	GGACAAOGTT	AATCCCTTTA	ACCACCACCG
22741	CAATGCGGGC	CTCCGTTATC	GCTCCATGTT	GTTGGGAAAC	GGCCGCTACG	TGCCCTTTCA
22801	CATTCAAGTG	CCCCAAAAGT	TTTTTGCCAT	TAAAAACCTC	CTCCTCCTGC	CAGGCTCATA
22861	TACATATGAA	TGGAACCTCA	GGAAOGATGT	TAACATGGTT	CTGCAGAGCT	CTCTGGGAAA
22921	CGATCTTAGA	GTTGACGGCG	CTAGCATTTAA	GTTTGACAGC	ATTTGTCTTT	ACGCCACCTT
22981	CTTCCCCATG	GCCCACAACA	CGGCTCCAC	GCTGGAAGCC	ATGCTCAGAA	ATGACACCAA
23041	CGACCAGTCC	TTPAATGACT	ACCTTTCCGC	CGCCAACATG	CTATACCCCA	TACCGCCAA
23101	CGCEACCAAC	GTGCCCATCT	CCATCCCATC	GCGCAACTGG	GCAGCATTTT	GCGGTTGGGG
23161	CTTCACACGC	TTGAAGACAA	AGGAAACCCC	TTCCCTGGGA	TCAGGCTACG	ACCTTTACTA
23221	CACCTACTCT	GGCTCCATAC	CATACCTTGA	CGGAACCTTC	TATCTTAATC	ACACCTTTAA
23281	GAAGGTGGCC	ATTACCTTTG	ACTCTTCTGT	TAGCTGGCCG	GGCAACGACC	GCTGCTTAC
23341	TCCCAATGAG	TTTGAGATTA	AACGCTCAGT	TGACGGGGAG	GGCTACAACG	TAGCTCAGTG
23401	CAACATGACC	AAGGACTGGT	TCCTGGTGCA	GATGTTGGCC	AACCTACAATA	TTGGCTACCA
23461	GGGCTTCTAC	ATTCCAGAAA	GCTACAAGGA	CCGCATGTAC	TGGTTCTTCA	GAAACTTTCA
23521	GCCCATGAGC	CGGCAAGTGG	TTGACGATAC	TAAATACAAG	GAGTATCAGC	AGGTTGGAAT
23581	TCCTTACCAG	CATAACAAC	CAGGATTGGT	AGGCTACCTC	GCTCCACCCA	TGCGCGAGGG
23641	ACAGGCTTAC	CCCGCCAACG	TGCCCTACCC	ACTAATAGGC	AAAACCGCGG	TTGACAGTAT
23701	TACCCAGAAA	AAGTTTCTTT	GCGATCGCAC	CCTTTGGCGC	ATCCCATTTT	CCAGTAACTT
23761	TATGTCCATG	GGCGCACTCA	CAGACCTGGG	CCAAAACCTT	CTCTAOGCCA	ACTCGCCCCA
23821	CGCGCTAGAC	ATGACTTTTG	AGGTGGATCC	CATGGAOGAG	CCCACCCCTT	TTTATGTTT
23881	GTTTGAGTTC	TTTGACGTGG	TCCGTGTGCA	CCAGCOGCAC	OGGGOGTCA	TCGAGACCGT
23941	GTACCTGGCG	ACGCCCTTCT	CGGCCGGCAA	CGCCACAACA	TAAAAGAAGC	AAGCAACATC
24001	AACRAACAGT	GCCGCCATGG	GCTCCAGTGA	CGAGGAAC	AAAGCCATTG	TCAAAGATCT
24061	TGGTTGTGGG	CCATATTTTT	TGGGCACCTA	TGACAAGCGC	TTTCCAGGCT	TTGTTTCTCC
24121	ACACAAGCTC	GCCTGGGCCA	TAGTCAATAC	GGCCGGTGGC	GAGACTGGGG	GCGTACACTG
24181	GATGGCCCTT	GCCTGGAAAC	CGCGCTCAAA	AACATGCTAC	CTCTTTGAGC	CCTTTGGCTT
24241	TTCTGACCAA	CGACTCAAGC	AGGTTTACCA	GTTTGAGTAC	GAGTCACTCC	TGCGCCGTAG
24301	CGCCATTGCT	TCTTCCCCCG	ACCGCTGTAT	AACGCTGGAA	AAGTCCACCC	AAAGCGTGCA
24361	GGGGCCCAAC	TCGGCCGCTT	GTGGACTATT	CTGCTGCATG	TTTCTCCACG	CCTTTGCCAA
24421	CTGGCCCCAA	ACTCCCATGG	ATCACAACCC	CACCATGAAC	CTTATTACCG	GGGTACCCAA
24481	CTCCATGCTT	AACAGTCCCC	AGGTACAGCC	CACCCCTGGT	CGCAACCAGG	AACAGCTCTA
24541	CAGCTTCTCT	GAGCGCCACT	CGCCCTACTT	CCGCAGCCAC	AGTGCGCAGA	TTAGGAGCGC
24601	CACCTCTTTT	TGTCACCTGA	AAAACATGTA	AAAATAATGT	ACTAGGAGAC	ACTTTCAATA
24661	AAGGCAATG	TTTTTATTTG	TACACTCTCG	GGTGATTATT	TACCCCCCAC	CCTTGCCGTC
24721	TGCGCGGTTT	AAAAATCAAA	GGGGTTCTGC	CGCGCATCGC	TATGCGCCAC	TGGCAGGGAC
24781	ACGTTGCGAT	ACTGGTGTTT	AGTGCTCCAC	TAAAACTCAG	GCACAACCAT	CCGCGGCAGC
24841	TCGGTGAAGT	TTTCACTCCA	CAGGCTGCGC	ACCATCACCA	ACCGGTTTAG	CAGGTGGGGC
24901	GCCGATATCT	TGAAGTCGCA	GTTGGGGCCT	CCGCCCTGCG	CGCGCGAGTT	GCGATACACA
24961	GGGTTGCAGC	ACTGGAACAC	TATCAGCGCC	GGGTGGTGCA	CGCTGGCCAG	CACGCTCTTG
25021	TCGGAGATCA	GATCCGCGTC	CAGGTCCCTC	GCGTTGCTCA	GGGCGAACGG	AGTCAACTTT
25081	GGTAGCTGCT	TTCCCAAAAA	GGGTGCATGC	CCAGGCTTTG	AGTTGCACTC	GCACOGTAGT
25141	GGCATCAGAA	GGTGACCGTG	CCCGGCTCTG	GCGTTAGGAT	ACAGCGCCTG	CATGAAAGCC
25201	TTGATCTGCT	TAAAAGCCAC	CTGAGCCTTT	GCGCCTTCAG	AGAAGAACAT	GCCGCAAGAC
25261	TTGCCGGA	ACTGATTGGC	CGGACAGGCC	GCGTCATGCA	CGCAGCACCT	TGCGTGGGTG
25321	TTGGAGATCT	GCACCACATT	TCGGCCCCAC	CGGTTCTTCA	CGATCTTGGC	CTTGCTAGAC
25381	TTCTCCTTCA	GCGCGCGCTG	CCCGTTTTCG	CTGTTCACAT	CCATTTCAAT	CACGTGCTCC
25441	TGATTTATCA	TAATGCTCCC	GTGTAGACAC	TAAAGCTCCG	CTTCGATCTC	AGCGCAGCGG
25501	TGCAGCCACA	ACGCGCAGCC	CGTGGGCTCG	TGGTGCTTGT	AGGTTACCTC	TGCAAAACGAC
25561	TGCAGGTACG	CCTGCAGGAA	TCGCCCCATC	ATCGTCACAA	AGGCTCTGTT	GCTGGTGAAG
25621	GTCAGCTGCA	ACCCGCGGTG	CTCCTCGTTT	AGCCAGGTCT	TGCATACGGC	CGCCAGAGCT
25681	TCCACTTGGT	CAGGCAGTAG	CTTGAAGTTT	GCCTTTAGAT	CGTTATCCAC	GTGGTACTTG
25741	TCCATCAACG	CGCGCGCAGC	CTCCATGCCC	TTCTCCACG	CAGACACGAT	CGGCAGGCTC

Nucleotide Sequence Analysis (cont.)

25801	AGCGGGTTTA	TCACCGTGCT	TTCACTTTCC	GCTTCACTGG	ACTCTTCCTT	TTCTCTTGCC
25861	GTCGCGATAC	CCCGCGCCAC	TGGGTGCTCT	TCATTTCAGCC	GCGGCACCGT	GCGCTTACCT
25921	CCCTTGCGGT	GCTTGATTAG	CACCGGTGGG	TTGCTGAAAC	CCACCATTTG	TAGCGCCACA
25981	TCTTCTCTTT	CTTCTCTCGT	GTCCAGGATC	ACCTCTGGCG	ATGGCGGGCG	CTGGGGCTTG
26041	GGAGAGGGGC	GCTTCTTTTT	CTTTTGGAC	GCAATGGCCA	AATCCGCGGT	CGAGGTCCAT
26101	GGCGCGGGC	TGGGTGTGCG	CGGCACCAGC	GCATCTTGTT	ACGAGTCTTC	TTCTCTCTCG
26161	GACTCGAGAC	GCCGCCTCAG	COGCTTTTTT	GGGGGCGCGC	GGGGAGGCGG	CGGOGAOGGC
26221	GACGGGGACG	ACACGTCCTC	CATGGTTGGT	GGACGTGCGG	COGCACCGCG	TCCGCGCTOG
26281	GGGGTGGTTT	CGCGCTGCTC	CTCTTCCCGA	CTGGCCATTT	CCTTCTCTTA	TAGGCAGAAA
26341	AAGATCATGG	AGTCAGTCGA	GAAGGAGGAC	AGCCTAACCG	CCCCCTTTGA	GTTGCGCCACC
26401	ACGCGCTCCA	CCGATGCGCG	CAACGCGCCT	ACCACCTTCC	CCGTGAGGGC	ACCCCGGCTT
26461	GAGGAGGAGG	AAGTGATTAT	CGAGCAGGAC	CCAGGTTTTG	TAAGCGAAGA	CGACGAGGAT
26521	CGCTCAGTAC	CAACAGAGGA	TAAAAAGCAA	GACCAGGACG	ACGCAGAGGC	AAACGAGGAA
26581	CAAGTCGGGC	GGGGGGACCA	AAGGCATGGC	GACTACCTAG	ATGTGGGAGA	CGACGTGCTG
26641	TTGAAGCATC	TGCAGCGCCA	GTGCGCCATT	ATCTGCGACG	CGTTGCAAGA	GCGCAGCGAT
26701	GTGCCCCCTG	CCATAGCGGA	TGTCAGCCCT	GCCTACGAAC	GCCACCTGTT	CTCACC GGCG
26761	GTACCCCCCA	AAAGCCCAAGA	AAACGGCACA	TGCGAGCCCA	ACCCGCGCCT	CAACTTCTAC
26821	CCGTTATTTG	CCGTGCCAGA	GGTGCTTGCC	ACCTATCACA	TCTTTTTCCA	AAACTGCAAG
26881	ATACCCCTAT	CCTGCGGTGC	CAACCGCAGC	CGAGCGGACA	AGCAGCTGGC	CTTGGCGCAG
26941	GGCGCTGTCA	TACCTGATAT	CGCCTCGCTC	GACGAAGTGC	CAAAAATCTT	TGAGGGTCTT
27001	GGAGCGGACG	AGAAAAGCGC	GGCAAACGCT	CTGCAACAAG	AAAACAGCGA	AAATGAAAGT
27061	CACGTGTGGG	TGCTGGTGGG	ACTTGAGGGT	GACAACGCGC	GCCTAGCCGT	GCTGAAACGC
27121	AGCATCGAGG	TCACCCACTT	TGCCTACCCG	GCACCTAACC	TACCCGCCAA	GGTTATGAGC
27181	ACAGTCATGA	GCGAGCTGAT	CGTGCGCCGT	GCACGACCCC	TGGAGAGGGA	TGCAAACTTG
27241	CAAGAACAAA	CCGAGGAGGG	CCTACCCGCA	GTTGGCGATG	AGCAGCTGGC	GCGCTGGCTT
27301	GAGAGCGCGG	AGCCTGCGGA	CTTGAGGAGG	CGAAGCAAGC	TAATGATGGC	CGCAGTGCTT
27361	GTTACCGTGG	AGCTTGAGTG	CATGCAGCGG	TTCTTTGCTG	ACCCGGAGAT	GCAGCGCAAG
27421	CTAGAGGAAA	CGTTGCACTA	CACCTTTTCG	CAGGGCTAAG	TGCGCCAGGC	CTGCAAAATT
27481	TCCAACGTGG	AGCTCTGCAA	CCTGGTCTCC	TACCTTGGA	TTTTGCAAGA	AAACCGCCTC
27541	GGGCAAAACG	TGCTTCATTC	CACGCTCAAG	GGCGAGGCGC	GCGCGACTA	GCGCGCGAC
27601	TGCGTTTACT	TATTTCTGTG	CTACACCTGG	CAAACGGCCA	TGGGCGTGTG	GCAGCAATGC
27661	CTGGAGGAGC	GCAACCTAAA	GGAGCTGCAG	AAGCTGCTAA	AGCAAAACTT	GAAGGACCTA
27721	TGGACGGCCT	TCAACGAGCG	CTCCGTGGCC	GCGCACCTGG	CGGACATTAT	CTTCCCGGAA
27781	CGCCTGCTTA	AAACCCCTGCA	ACAGGGTCTG	CCAGACTTCA	CCAGTCAAAG	CATGTTGCAA
27841	AACTTTAGGA	ACTTTATCCT	AGAGCGTTCA	GGAAATTCTG	CCGCCACCTG	CTGTGCGCTT
27901	CCTAGCGACT	TTGTGCCCCAT	TAAGTACCGT	GAATGCCCTC	CGCCGCTTTG	GGGTCACTGC
27961	TACCTTCTGC	AGCTAGCCAA	CTACCTTGCC	TACCACTCCG	ACATCATGGA	AGACGTGAGC
28021	GGTGACGGCC	TACTGGAGTG	TCACTGTGCG	TGCAACCTAT	GCACCCCGCA	CCGCTCCCTG
28081	GTCTGCAATT	CGCAACTGCT	TAGCGAAAGT	CAAATTATOG	GTACCTTTGA	GCTGCAGGGT
28141	CCCTCGCCTG	ACGAAAAGTC	CGCGGCTCCG	GGGTTGAAAC	TCACCTCGGG	GCTGTGGACG
28201	TGCGCTTACC	TTCGCAAATT	TGTACCTGAG	GACTACCACG	CCCACGAGAT	TAGGTTCTAC
28261	GAAGACCAAT	CCCGCCCGCC	AAATGCGGAG	CTTACCGCCT	GCGTCATTAC	CCAGGGCCAC
28321	ATCCTTGGCC	AATTGCAAGC	CATCAACAAA	GCCCGCCAAG	AGTTTCTGCT	ACGAAAGGGA
28381	CGGGGGGTTT	ACCTGGACCC	CCAGTCCGGC	GAGGAGCTCA	ACCCAATCCC	CCCGCGCGCG
28441	CAGCCCTATC	AGCAGCCGCG	GGCCCTTGCT	TCCAGGATG	GCACCCAAAA	AGAAGCTGCA
28501	GCTGCCGCCG	CCGTCACCCA	CGGACGAGGA	GGAATACTGG	GACAGTCAGG	CAGAGGAGGT
28561	TTTGGAACGAG	GAGGAGGAGA	TGATGGAAGA	CTGGGACAGC	CTAGACGAAG	CTTCCGAGGC
28621	CGAAGAGGTG	TCAGACGAAA	CACCGTCAAC	CTCGGTGCGA	TTCCCTCTGC	CGGCGCCCCA
28681	GAAATTGGCA	ACCGTTCCCA	GCATOGCTAC	AACCTCGCT	CCTCAGGCGC	CGCCGGCACT
28741	GCCTGTTCCG	CGACCCAAAC	GTAAGTGGGA	CACCACTGGA	ACCAGGGCCG	GTAAGTCTAA
28801	GCAGCCGCCG	CCGTTAGCCC	AAGAGCAACA	ACAGCGCCAA	GGCTACCGCT	CGTGGCGCGG
28861	GCACAAGAAC	GCCATAGTTG	CTTGCTTGCA	AGACTGTGGG	GGCAACATCT	CCTTCGCCCC
28921	CCGCTTTCTT	CTCTACCATC	ACGGCGTGGC	CTTCCCCCGT	AACATCCTGC	ATTACTACCG
28981	TCATCTCTAC	AGCCCCTACT	GCACCGGCGG	CAGCGGCAGC	GGCAGCAACA	GCAGCGGTCA
29041	CACAGAAGCA	AAGGCGACCG	GATAGCAAGA	CTCTGACAAA	GCCCAAGAAA	TCCACAGCGG
29101	CGGCAGCAGC	AGGAGGAGGA	GCGCTGCGTC	TGGCGCCCAA	CGAACCCGTA	TGACCCCGCG
29161	AGCTTAGAAA	TAGGATTTTT	CCCACTCTGT	ATGCTATATT	TCAACAAAGC	AGGGGCCAAG

Nucleotide Sequence Analysis (cont.)

29221	AACAAGAGCT	GAAAATAAAA	AACAGGTCTC	TGCGCTCCCT	CACCGGCAGC	TGCCTGTATC
29281	ACAAAAGCGA	AGATCAGCTT	CGGCGCAOCC	TGGAAGACGC	GGAGGCTCTC	TTCAGCAAAT
29341	ACTGCGCGCT	GACTCTTAAG	GACTAGTTTC	GCGCCCTTTC	TCAAATTTAA	GCGCGAAAAC
29401	TACGTCACTCT	CCAGCGGCCA	CACCGGGGCG	CAGCACCTGT	CGTCAGCGCC	ATTATGAGCA
29461	AGGAAATTCC	CACGCCCTAC	ATGTGGAGTT	ACCAGCCACA	AATGGGACTT	GCGGCTGGAG
29521	CTGCCCAAGA	CTACTCAACC	CGAATAAACT	ACATGAGCGC	GGGACCCAC	ATGATATCCC
29581	GGGTCAACGG	AATCCGCGCC	CACOGAAACC	GAATTCTCCT	CGAACAGGCG	GCTATTACCA
29641	CCACACCTCG	TAATAACCTT	AATCCCCGTA	GTTGGCCCCG	TGCCCTGGTG	TACCAGGAAA
29701	GTCCCGCTCC	CACCACTGTG	GTACTTCCCA	GAGACGCCCA	GGCCGAAGTT	CAGATGACTA
29761	ACTCAGGGGC	GCAGCTTGCG	GGCGGCTTTC	GTCACAGGGT	GCGGTGCGCC	GCGCAGGGTA
29821	TAACTCACCT	GAAAATCAGA	GGGCGAGGTA	PTCAGCTCAA	CGACGAGTCG	GTGAGCTCCT
29881	CTCTTGGTCT	CCGTCCGGAC	GGGACATTTT	AGATOGGCGG	CGCTGGCGCG	TCTTCATTTA
29941	CGCCCCGTCA	GGCGATCCTA	ACTCTGCAGA	CCTCGTCTCT	CGAGCGCGCG	TCCGGAGGCA
30001	TGGGAACCTCT	ACAATTTATT	GAGGAGTTGG	TGCCCTTCGGT	TTACTTCAAC	CCCTTTTCTG
30061	GACCTCCCGG	CCACTACCCG	GACCACTTTA	TTCCCAACTT	TGACGCGGTG	AAAGACTCGG
30121	CGGACGGCTA	CGACTGAATG	ACCACTGGAG	AGGCAGAGCG	ACTGCGCCTG	ACACACCTCG
30181	ACCACTGCCG	CCGCCACAAG	TGCTTTTGCCC	GCGGCTCCCG	TGAGTTTTGT	TACTTTGAAT
30241	TGCCCGAAGA	GCATATCGAG	GGCCCGGCGC	ACGCGCTCCG	GCTCACCACC	CAGGTAGAGC
30301	TTACACGTAG	CCTGATTCCG	GAGTTTACCA	AGCGCCCCCT	GCTAGTGGAG	CGGGAGCGGG
30361	GTCCCTGTGT	TCTGACCGTG	GTTTGCAACT	GTCTTAACCC	TGGATTACAT	CAAGATCTTT
30421	GTTGTCACTCT	CTGTGCTGAG	TATAATAAAT	ACAGAAATTA	GAATCTACTG	GGGCTCCTGT
30481	CGCCATCCTG	TGAACGCCAC	CGTTTTTTACC	CACCCAAAGC	AGACCAAAGC	AAACCTCACC
30541	TCGGTTTTGC	ACAAGCGGGC	CAATAAGTAC	CTTACCTGGT	ACTTTAACGG	CTCTTCATTT
30601	GTAATTTTACA	ACAGTTTCCA	GCGAGACGAA	GTAAGTTTGC	CACACAACCT	TCTCGGCTTC
30661	AACCTACCCG	TCAAGAAAAA	CACCACCACC	ACCACCTCC	TCACCTGCGG	GGAAAGTACG
30721	AGTGCGTCAC	CGGTGCTGCG	GCCACACCT	ACAGCCTGAG	CGTAACCAGA	CATTACTCCC
30781	ATTTTTTCCAA	AACAGGAGGT	GAGCTCAACT	CCCGGAACTC	AGGTCAAAAA	AGCATTTTGC
30841	GGGGTGCTGG	GATTTTTTAA	TTAAGTATAT	GAGCAATTCA	AGTAACCTTA	CAAGCTTGTC
30901	TAATTTTTTCT	GGAATTGCGG	TCCGGGTTAT	CCTTACTCTT	GTAATTCTGT	TTATTCTTAT
30961	ACTAGCACTT	CTGTGCCCTTA	GGGTGCGCGC	CTGCTGCACG	CACGTTTGTA	CCTATTGTCA
31021	GCTTTTTTAAA	CGCTGGGGGC	AACATCCAAG	ATGAGGTACA	TGATTTTAGG	CTTGCTCGCC
31081	CTTGCGGCAG	TCTGCAGCGC	TGCCAAAAAG	GTTGAGTTTA	AGGAACCAGC	TTGCAATGTT
31141	ACATTTTAAAT	CAGAAGCTAA	TGAATGCACT	ACTCTTATAA	AATGCACCAC	AGAACATGAA
31201	AAGTTTATTA	TTCGCCACAA	AGACAAAATT	GGCAAGTATG	CTGTATATGC	TATTTGGCAG
31261	CCAGGTGACA	CTAACGACTA	TAATGTCACA	GTCTTCCAAG	GTGAAAATCG	TAAAACCTTTT
31321	ATGTATAAAT	TTCCATTTTA	TGAAATGTGC	GATATTACCA	TGTACATGAG	CAAACAGTAC
31381	AAGTTGTGGC	CCCCACAAAA	GTGTTTAGAG	AACACTGGCA	CCTTTTGTTC	CACGCTCTG
31441	CTTATTACAG	CGCTTGCTTT	GGTATGTACC	TTACTTTATC	TCAAATACAA	AAGCAGACGC
31501	AGTTTTATTG	ATGAAAAGAA	AATGCCTTGA	TTTTCCGCTT	GCTTGATTTC	CCCTGGACAA
31561	TTTACTCTAT	GTGGGATATG	CTCCAGGCGG	GCAAGATTAT	ACCCACAACC	TTCAAAATCAA
31621	ACTTTCCTGG	ACGTTAGCGC	CTGATTTCTG	CEAGCGCCTG	CACCTGCAAT	TTGATCAAAC
31681	CCAGCTTCAG	CTTGCCCTGCT	CCAGAGATGA	COGGCTCAAC	CATCGCGCCC	ACAACGGACT
31741	ATCGCAACAC	CACCTGCTACC	GGACTAACAT	CTGCCCTAAA	TTTACCCCAA	GTTCATGCCT
31801	TTGTCAATGA	CTGGGCGAGC	TTGGACATGT	GGTGGTTTTT	CATAGCGCTT	ATGTTTGTFT
31861	GCCTTATTAT	TATGTGGCTT	ATTTGTTGCC	TAAAGCGCAG	ACGCGCCAGA	CCCCCATCT
31921	ATAGGCCCTAT	CATTGTGCTC	AACCCACACA	ATGAAAAAAT	TCATAGATTG	GACGGTCTGA
31981	AACCATGTTT	TCTTCTTTTA	CAGTATGATT	AAATGAGACA	TGATTCTCTG	AGTTCTTATA
32041	TTATTGACCC	TTGTTGCGCT	TTTCTGTGCG	TGCTCTACAT	TGGCCGCGGT	CGCTCACATC
32101	GAAGTAGATT	GCATCCCACC	TTTCACAGTT	TACCTGCTTT	ACGGATTGTG	CACCCTTATC
32161	CTCATCTGCA	GCCTCGTCAC	TGTAGTCATC	GCCTTCATTG	AGTTTCATTG	CTGGGTTTGT
32221	GTGCGCATTG	CGTACCTCAG	GCACCATCOG	CAATACAGAG	ACAGGACTAT	AGCTGATCTT
32281	CTCAGAAATC	TTTAAATTATG	AAACGGAGTG	TCATTTTTGT	TTTGCTGATT	TTTTGCGCCC
32341	TACCTGTGCT	TTGCTCCCAA	ACCTCAGCGC	CTCCCAAAAG	ACATATTTC	TGCAGATTCA
32401	CTCAAAATAG	GAACATTCCC	AGCTGCTACA	ACAAACAGAG	CGATTGTGTA	GAAGCCTGGT
32461	TATACGCCAT	CATCTCTGTC	ATGGTTTTTT	GCAGTACCAT	TTTTGCCCTA	GCCATATATC
32521	CATACCTTGA	CATTGGCTGG	AATGCCATAG	ATGCCATGAA	CCACCCTACT	TTCCAGTGTC
32581	CCGCTGTCAT	ACCACTGCAA	CAGGTTATTG	CCCCAATCAA	TCAGCCTCGC	CCCCCTTCTC

Nucleotide Sequence Analysis (cont.)

```

32641 CCACCCCCAC TGAGATTAGC TACTTTAATT TGACAGGTGG AGATGACTGA ATCTCTAGAT
32701 CTAGAATTGG ATGGAATTAA CACCGAACAG CGCCTACTAG AAAGGCGCAA GGCGGCGTCC
32761 GAGCGAGAAC GCCTAAAACA AGAAGTTGAA GACATGGTTA ACCTACACCA GTGTAAAAGA
32821 GGTATCTTTT GTGTGGTCAA GCAGGCCAAA CTTACCTACG AAAAAACCAC TACCGGCAAC
32881 CGCCTCAGCT ACAAGCTACC CACCCAGOGC CAAAACTGG TGCTTATGGT GGGAGAAAAA
32941 CCTATCACCG TCACCCAGCA CTGGGCAGAA ACAGAGGGCT GCCTGCACCTT CCCCTATCAG
33001 GGTCCAGAGG ACCTCTGCAC TCTTATTAAA ACCATGTGTG GTATTAGAGA TCTTATTCCA
33061 TTCAACTAAC ATAAACACAC AATAAATTAC TTACTTAAAA TCAGTCAGCA AATCTTTGTC
33121 CAGCTTATTC AGCATCACCT CCTTTCCTTC CTCCCAACTC TGGTATCTCA GCCCGCTTTT
33181 AGCTGCAAAAC TTTCTCCAAA GTTTAAATGG GATGTCAAAT TCCTCATGTT CTGTGCCCTC
33241 CGCACCCACT ATCTTCATAT TGTGTGAGAT GAAACGGCGC AGACCGTCTG AAGACACCTT
33301 CAACCCCGTG TATCCATATG ACACAGAAAC CGGGCCTCCA ACTGTGCCCT TTCTTACCCC
33361 TCCATTTGTT TCACCCAATG GTTTCCAAGA AAGTCCCCCT GGAGTTCTCT CTCTAOGGOT
33421 CTCGGAACCT TTGGACACCT CCCACGGCAT GCTTGOGCTT AAAATGGGCA GOGGTCTTAC
33481 CCTAGACAAG GCCGGAACCC TCACCTOCCA AATGTAAACC ACTGTTACTC AGCCACTTAA
33541 AAAAACAAAG TCAAAACATA GTTTGACAC CTCCGCACCA CTTACAATTA CCTCAGGGCG
33601 CCTAACAGTG GCAACCACCG CTCCTCTGAT AGTTACTAGC GCGCTCTTA GCGTACAGTC
33661 ACAAGCCCCA CTGACCGTGC AAGACTCCAA ACTAAGCATT GCTACTAAAG GGCCATTAC
33721 AGTGTGAGAT GGAAGCTAG CCCTGCAAC ATCAGCCCCC CTCTCTGGCA GTGACAGOGA
33781 CACCCTTACT GTAAGTGCAT CACCCCGCT AACTACTGCC ACGGGTAGCT TGGGCATTAA
33841 CATGGAAGAT CCTATTTATG TAAATAATGG AAAAATAGGA ATTAAAATAA GCGGTCTTTT
33901 GCAAGTAGCA CAAAACCTCG ATACACTAAC AGTAGTTACT GGACCAGGTG TCACCGTTGA
33961 ACAAACTCC CTTAGAACCA AAGTTGCAGG AGCTATTGGT TATGATTCAT CAAACAACAT
34021 GGAAATTAAA ACGGGCGGTG GCATGOGTAT AAATAACAAC TTGTTAATTC TAGATGTGGA
34081 TTACCCATTT GATGCTCAA CAAAACCTAC TCTTAACTG GGCAGGGAC CCCTGTATAT
34141 TAATGCATCT CATAACTTGG ACATAAATA TAACAGAGGC CTATACCTTT TTAATGCATC
34201 AAACAATACT AAAAACTGG AAGTTAGCAT AAAAAATCC AGTGGACTAA ACTTTGATAA
34261 TACTGCCATA GCTATAAATG CAGGAAAGGG TCTGGAGTTT GATACAAACA CATCTGAGTC
34321 TCCAGATATC AACCCAATAA AAATAAAAT TGGCTCTGGC ATTGATTACA ATGAAAAOGG
34381 TGCCATGATT ACTAAACTTG GAGOGGGTTT AAGCTTTGAC AACTCAGGGG CCATTACAAT
34441 AGGAAACAAA AATGATGACA AACTTACCCT GTGGACAACC CCAGACCCAT CTCCTAACTG
34501 CAGAATTCAT TCAGATAATG ACTGCAAAT TACTTTGGTT CTTACAAAAT GTGGGAGTCA
34561 AGTACTAGCT ACTGTAGCTG CTTTGGCTGT ATCTGGAGAT CTTTCATCCA TGACAGGCAC
34621 CGTTGCAAGT GTTAGTATAT TCCCTAGATT TGACCAAAAC GGTGTTCTAA TGGAGAACTC
34681 CTCACTTAAA AAACATTACT GGAACCTTAG AAATGGGAAC TCAACTAATG CAAATCCATA
34741 CACAAATGCA GTTGGATTTA TGCCTAACCT TCTAGCCTAT CCAAAAACCC AAAGTCAAAC
34801 TGCTAAAAAT AACATTGTCA GTCAAGTTTA CTGTCATGGT GATAAACTA AACCTATGAT
34861 ACTTACCATT ACACCTAATG GCACTAGTGA ATCCACAGAA ACTAGCGAGG TAAGCACTTA
34921 CTCTATGTCT TTTACATGGT CCTGGGAAAG TGGAAAATAC ACCACTGAAA CTTTTGCTAC
34981 CAACTCTTAC ACCTTCTCCT ACATTGCCCA GGAATAAAGA ATCGTGAACC TGTGTCATGT
35041 TATGTTTCAA CGTGGGATCC TTTATTATAG GGGAAGTCCA CGCCTACATG GGGGTAGAGT
35101 CATAATCGTG CATCAGGATA GGGCGGTGGT GCTGCAGCAG CGCGGAATA AACTGTCTCC
35161 GCCGCCGCTC CGTCTGCGAG GAATACAACA TGGCAGTGGT CTCCTCAGCG ATGATTGCGA
35221 CCGCCCGCAG CATGAGACGC CTTGTCTCTC GGCACAGCA GCGCACCTTG ATCTCACTTA
35281 AATCAGCACA GTAAGTGCAG CACAGCACA CAATATTGTT CAAAATCCCA CAGTGCAAGG
35341 CGCTGTATCC AAAGCTCATG GCGGGGACCA CAGAACCAC GTGGCCATCA TACCACAAGC
35401 GCAGGTAGAT TAAGTGGCGA CCCCTCATAA ACACGCTGGA CATAAACATT ACCTCTTTTG
35461 GCATGTTGTA ATTCACCACC TCCCGGTACC ATATAAACCT CTGATTAAAC ATGGCGCCAT
35521 CCACCACCAT CCTAAACCAG CTGGCCAAAA CCTGCCCGCC GGCTATGCAC TGCAGGGAAC
35581 CGGGACTGGA ACAATGACAG TGGAGAGCCC AGGACTCGTA ACCATGGATC ATCATGCTCG
35641 TCATGATATC AATGTTGGCA CAACACAGGC ACACGTGCAT ACACTTCTCT AGGATTACAA
35701 GCTCCTCCCG CGTCAGAACC ATATCCCAGG GAACAACCCA TTCCTGAATC AGCGTAAATC
35761 CCACACTGCA GGAAGACCT CGCAGGTAAC TCACGTTGTG CATGTGCAAA GTGTTACATT
35821 CGGGCAGCAG CGGATGATCC TCCAGTATGG TGGCGCGGT CTCTGTCTCA AAAGGAGGTA
35881 GGCGATCCCT ACTGTACGGA GTGCGCCGAG ACAACCGAGA TCGTGTGGT CGTAGTGTCA
35941 TGCACAAATG AACGCCGAG GTAGTCATAT TTCATCGACA CGGCACCAGC TCAATCACTC
36001 ACAGTGTAAG AAGGCCAAG TACAGAGCGA GTATATATAG GACTAAAAAA TGACGTAACG

```

Nucleotide Sequence Analysis (cont.)

36061 GTTAAAGTCC ACAAAAACA CCCAGAAAAC CGCACGOGAA CCTACGCCCA GAAACGAAAG
36121 CCAAAAACC CACAACCTCC TCAAATCTTC ACTTCOGTTT TCCCACGATA CGTCACTTCC
36181 CATTTTAAAA AACTACAAT TCCCAATACA TGCAAGTTAC TCGCCCTAA AACCTACGTC
36241 ACCCGCCCCG TTCCCACGCC CCGGCCACG TCACAAACTC CACCCCTCA TTATCATATT
36301 GGCTTCAATC CAAAATAAGG TATATTATGA TGATG

//

SEQUENCE LISTING

(1) GENERAL INFORMATION:

5

(i) APPLICANTS: Gregory, R.J., Armentano, D., Couture, L.A., Smith,
A.E.

10

(ii) TITLE OF INVENTION: GENE THERAPY FOR CYSTIC FIBROSIS

(iii) NUMBER OF SEQUENCES: 9

15

(iv) CORRESPONDENCE ADDRESS:

- (A) ADDRESSEE: LAHIVE & COCKFIELD
- (B) STREET: 60 STATE STREET, SUITE 510
- (C) CITY: BOSTON
- (D) STATE: MASSACHUSETTS
- (E) COUNTRY: USA
- (F) ZIP: 02109

20

(v) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: ASCII

25

(vi) CURRENT APPLICATION DATA:

- (A) APPLICATION NUMBER:
- (B) FILING DATE: 02-DEC-1993
- (C) CLASSIFICATION:

30

(vii) PRIOR APPLICATION DATA:

- (A) APPLICATION NUMBER: US 07/985,478
- (B) FILING DATE: 02-DEC-1992
- (C) CLASSIFICATION:

35

(viii) ATTORNEY/AGENT INFORMATION:

- (A) NAME: Hanley, Elizabeth A.
- (B) REGISTRATION NUMBER: 33,505
- (C) REFERENCE/DOCKET NUMBER: NZI-014CP2PC

40

(ix) TELECOMMUNICATION INFORMATION:

- (A) TELEPHONE: (617) 227-7400
- (B) TELEFAX: (617) 227-5941

45

(2) INFORMATION FOR SEQ ID NO:1:

50

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 6129 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

55

(ii) MOLECULE TYPE: cDNA

- 97 -

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 133..4572

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

10	AATTGGAAGC AAATGACATC ACAGCAGGTC AGAGAAAAAG GGTTGAGCGG CAGGCACCCA	60
	GAGTAGTAGG TCTTTGGCAT TAGGAGCTTG AGCCCAGACG GCCCTAGCAG GGACCCCAGC	120
	GCCCCGAGAGA CC ATG CAG AGG TCG CCT CTG GAA AAG GCC AGC GTT GTC	168
15	Met Gln Arg Ser Pro Leu Glu Lys Ala Ser Val Val	
	1 5 10	
	TCC AAA CTT TTT TTC AGC TGG ACC AGA CCA ATT TTG AGG AAA GGA TAC	216
	Ser Lys Leu Phe Phe Ser Trp Thr Arg Pro Ile Leu Arg Lys Gly Tyr	
	15 20 25	
20	AGA CAG CGC CTG GAA TTG TCA GAC ATA TAC CAA ATC CCT TCT GTT GAT	264
	Arg Gln Arg Leu Glu Leu Ser Asp Ile Tyr Gln Ile Pro Ser Val Asp	
	30 35 40	
25	TCT GCT GAC AAT CTA TCT GAA AAA TTG GAA AGA GAA TGG GAT AGA GAG	312
	Ser Ala Asp Asn Leu Ser Glu Lys Leu Glu Arg Glu Trp Asp Arg Glu	
	45 50 55 60	
30	CTG GCT TCA AAG AAA AAT CCT AAA CTC ATT AAT GCC CTT CGG CGA TGT	360
	Leu Ala Ser Lys Lys Asn Pro Lys Leu Ile Asn Ala Leu Arg Arg Cys	
	65 70 75	
35	TTT TTC TGG AGA TTT ATG TTC TAT GGA ATC TTT TTA TAT TTA GGG GAA	408
	Phe Phe Trp Arg Phe Met Phe Tyr Gly Ile Phe Leu Tyr Leu Gly Glu	
	80 85 90	
	GTC ACC AAA GCA GTA CAG CCT CTC TTA CTG GGA AGA ATC ATA GCT TCC	456
	Val Thr Lys Ala Val Gln Pro Leu Leu Leu Gly Arg Ile Ile Ala Ser	
	95 100 105	
40	TAT GAC CCG GAT AAC AAG GAG GAA CGC TCT ATC GCG ATT TAT CTA GGC	504
	Tyr Asp Pro Asp Asn Lys Glu Glu Arg Ser Ile Ala Ile Tyr Leu Gly	
	110 115 120	
45	ATA GGC TTA TGC CTT CTC TTT ATT GTG AGG ACA CTG CTC CTA CAC CCA	552
	Ile Gly Leu Cys Leu Leu Phe Ile Val Arg Thr Leu Leu Leu His Pro	
	125 130 135 140	
50	GCC ATT TTT GGC CTT CAT CAC ATT GGA ATG CAG ATG AGA ATA GCT ATG	600
	Ala Ile Phe Gly Leu His His Ile Gly Met Gln Met Arg Ile Ala Met	
	145 150 155	
55	TTT AGT TTG ATT TAT AAG AAG ACT TTA AAG CTG TCA AGC CGT GTT CTA	648
	Phe Ser Leu Ile Tyr Lys Lys Thr Leu Lys Leu Ser Ser Arg Val Leu	
	160 165 170	

	GAT AAA ATA AGT ATT GGA CAA CTT GTT AGT CTC CTT TCC AAC AAC CTG	696
	Asp Lys Ile Ser Ile Gly Gln Leu Val Ser Leu Leu Ser Asn Asn Leu	
	175 180 185	
5	AAC AAA TTT GAT GAA GGA CTT GCA TTG GCA CAT TTC GTG TGG ATC GCT	744
	Asn Lys Phe Asp Glu Gly Leu Ala Leu Ala His Phe Val Trp Ile Ala	
	190 195 200	
10	CCT TTG CAA GTG GCA CTC CTC ATG GGG CTA ATC TGG GAG TTG TTA CAG	792
	Pro Leu Gln Val Ala Leu Leu Met Gly Leu Ile Trp Glu Leu Leu Gln	
	205 210 215 220	
15	GCG TCT GCC TTC TGT GGA CTT GGT TTC CTG ATA GTC CTT GCC CTT TTT	840
	Ala Ser Ala Phe Cys Gly Leu Gly Phe Leu Ile Val Leu Ala Leu Phe	
	225 230 235	
20	CAG GCT GGG CTA GGG AGA ATG ATG ATG AAG TAC AGA GAT CAG AGA GCT	888
	Gln Ala Gly Leu Gly Arg Met Met Met Lys Tyr Arg Asp Gln Arg Ala	
	240 245 250	
25	GGG AAG ATC AGT GAA AGA CTT GTG ATT ACC TCA GAA ATG ATT GAA AAT	936
	Gly Lys Ile Ser Glu Arg Leu Val Ile Thr Ser Glu Met Ile Glu Asn	
	255 260 265	
30	ATC CAA TCT GTT AAG GCA TAC TGC TGG GAA GAA GCA ATG GAA AAA ATG	984
	Ile Gln Ser Val Lys Ala Tyr Cys Trp Glu Glu Ala Met Glu Lys Met	
	270 275 280	
35	ATT GAA AAC TTA AGA CAA ACA GAA CTG AAA CTG ACT CGG AAG GCA GCC	1032
	Ile Glu Asn Leu Arg Gln Thr Glu Leu Lys Leu Thr Arg Lys Ala Ala	
	285 290 295 300	
40	TAT GTG AGA TAC TTC AAT AGC TCA GCC TTC TTC TTC TCA GGG TTC TTT	1080
	Tyr Val Arg Tyr Phe Asn Ser Ser Ala Phe Phe Phe Ser Gly Phe Phe	
	305 310 315	
45	GTG GTG TTT TTA TCT GTG CTT CCC TAT GCA CTA ATC AAA GGA ATC ATC	1128
	Val Val Phe Leu Ser Val Leu Pro Tyr Ala Leu Ile Lys Gly Ile Ile	
	320 325 330	
50	CTC CGG AAA ATA TTC ACC ACC ATC TCA TTC TGC ATT GTT CTG CGC ATG	1176
	Leu Arg Lys Ile Phe Thr Thr Ile Ser Phe Cys Ile Val Leu Arg Met	
	335 340 345	
55	GCG GTC ACT CGG CAA TTT CCC TGG GCT GTA CAA ACA TGG TAT GAC TCT	1224
	Ala Val Thr Arg Gln Phe Pro Trp Ala Val Gln Thr Trp Tyr Asp Ser	
	350 355 360	
60	CTT GGA GCA ATA AAC AAA ATA CAG GAT TTC TTA CAA AAG CAA GAA TAT	1272
	Leu Gly Ala Ile Asn Lys Ile Gln Asp Phe Leu Gln Lys Gln Glu Tyr	
	365 370 375 380	
65	AAG ACA TTG GAA TAT AAC TTA ACG ACT ACA GAA GTA GTG ATG GAG AAT	1320
	Lys Thr Leu Glu Tyr Asn Leu Thr Thr Thr Glu Val Val Met Glu Asn	
	385 390 395	

	GTA	ACA	GCC	TTC	TGG	GAG	GAG	GGA	TTT	GGG	GAA	TTA	TTT	GAG	AAA	GCA	1368
	Val	Thr	Ala	Phe	Trp	Glu	Glu	Gly	Phe	Gly	Glu	Leu	Phe	Glu	Lys	Ala	
			400					405						410			
5	AAA	CAA	AAC	AAT	AAC	AAT	AGA	AAA	ACT	TCT	AAT	GGT	GAT	GAC	AGC	CTC	1416
	Lys	Gln	Asn	Asn	Asn	Asn	Arg	Lys	Thr	Ser	Asn	Gly	Asp	Asp	Ser	Leu	
			415					420					425				
10	TTC	TTC	AGT	AAT	TTC	TCA	CTT	CTT	GGT	ACT	CCT	GTC	CTG	AAA	GAT	ATT	1464
	Phe	Phe	Ser	Asn	Phe	Ser	Leu	Leu	Gly	Thr	Pro	Val	Leu	Lys	Asp	Ile	
			430				435					440					
15	AAT	TTC	AAG	ATA	GAA	AGA	GGA	CAG	TTG	TTG	GCG	GTT	GCT	GGA	TCC	ACT	1512
	Asn	Phe	Lys	Ile	Glu	Arg	Gly	Gln	Leu	Leu	Ala	Val	Ala	Gly	Ser	Thr	
	445					450					455					460	
20	GGA	GCA	GGC	AAG	ACT	TCA	CTT	CTA	ATG	ATG	ATT	ATG	GGA	GAA	CTG	GAG	1560
	Gly	Ala	Gly	Lys	Thr	Ser	Leu	Leu	Met	Met	Ile	Met	Gly	Glu	Leu	Glu	
				465					470						475		
25	CCT	TCA	GAG	GGT	AAA	ATT	AAG	CAC	AGT	GGA	AGA	ATT	TCA	TTC	TGT	TCT	1608
	Pro	Ser	Glu	Gly	Lys	Ile	Lys	His	Ser	Gly	Arg	Ile	Ser	Phe	Cys	Ser	
			480					485						490			
30	CAG	TTT	TCC	TGG	ATT	ATG	CCT	GGC	ACC	ATT	AAA	GAA	AAT	ATC	ATC	TTT	1656
	Gln	Phe	Ser	Trp	Ile	Met	Pro	Gly	Thr	Ile	Lys	Glu	Asn	Ile	Ile	Phe	
			495				500						505				
35	GGT	GTT	TCC	TAT	GAT	GAA	TAT	AGA	TAC	AGA	AGC	GTC	ATC	AAA	GCA	TGC	1704
	Gly	Val	Ser	Tyr	Asp	Glu	Tyr	Arg	Tyr	Arg	Ser	Val	Ile	Lys	Ala	Cys	
		510					515					520					
40	CAA	CTA	GAA	GAG	GAC	ATC	TCC	AAG	TTT	GCA	GAG	AAA	GAC	AAT	ATA	GTT	1752
	Gln	Leu	Glu	Glu	Asp	Ile	Ser	Lys	Phe	Ala	Glu	Lys	Asp	Asn	Ile	Val	
	525					530					535					540	
45	CTT	GGA	GAA	GGT	GGA	ATC	ACA	CTG	AGT	GGA	GGT	CAA	CGA	GCA	AGA	ATT	1800
	Leu	Gly	Glu	Gly	Gly	Ile	Thr	Leu	Ser	Gly	Gly	Gln	Arg	Ala	Arg	Ile	
				545					550						555		
50	TCT	TTA	GCA	AGA	GCA	GTA	TAC	AAA	GAT	GCT	GAT	TTG	TAT	TTA	TTA	GAC	1848
	Ser	Leu	Ala	Arg	Ala	Val	Tyr	Lys	Asp	Ala	Asp	Leu	Tyr	Leu	Leu	Asp	
			560					565						570			
55	TCT	CCT	TTT	GGA	TAC	CTA	GAT	GTT	TTA	ACA	GAA	AAA	GAA	ATA	TTT	GAA	1896
	Ser	Pro	Phe	Gly	Tyr	Leu	Asp	Val	Leu	Thr	Glu	Lys	Glu	Ile	Phe	Glu	
			575				580						585				
60	AGC	TGT	GTC	TGT	AAA	CTG	ATG	GCT	AAC	AAA	ACT	AGG	ATT	TTG	GTC	ACT	1944
	Ser	Cys	Val	Cys	Lys	Leu	Met	Ala	Asn	Lys	Thr	Arg	Ile	Leu	Val	Thr	
		590					595					600					
65	TCT	AAA	ATG	GAA	CAT	TTA	AAG	AAA	GCT	GAC	AAA	ATA	TTA	ATT	TTG	CAT	1992
	Ser	Lys	Met	Glu	His	Leu	Lys	Lys	Ala	Asp	Lys	Ile	Leu	Ile	Leu	His	
	605					610					615					620	

- 100 -

	GAA GGT AGC AGC TAT TTT TAT GGG ACA TTT TCA GAA CTC CAA AAT CTA	2040
	Glu Gly Ser Ser Tyr Phe Tyr Gly Thr Phe Ser Glu Leu Gln Asn Leu	
	625 630 635	
5	CAG CCA GAC TTT AGC TCA AAA CTC ATG GGA TGT GAT TCT TTC GAC CAA	2088
	Gln Pro Asp Phe Ser Ser Lys Leu Met Gly Cys Asp Ser Phe Asp Gln	
	640 645 650	
10	TTT AGT GCA GAA AGA AGA AAT TCA ATC CTA ACT GAG ACC TTA CAC CGT	2136
	Phe Ser Ala Glu Arg Arg Asn Ser Ile Leu Thr Glu Thr Leu His Arg	
	655 660 665	
15	TTC TCA TTA GAA GGA GAT GCT CCT GTC TCC TGG ACA GAA ACA AAA AAA	2184
	Phe Ser Leu Glu Gly Asp Ala Pro Val Ser Trp Thr Glu Thr Lys Lys	
	670 675 680	
20	CAA TCT TTT AAA CAG ACT GGA GAG TTT GGG GAA AAA AGG AAG AAT TCT	2232
	Gln Ser Phe Lys Gln Thr Gly Glu Phe Gly Glu Lys Arg Lys Asn Ser	
	685 690 695 700	
25	ATT CTC AAT CCA ATC AAC TCT ATA CGA AAA TTT TCC ATT GTG CAA AAG	2280
	Ile Leu Asn Pro Ile Asn Ser Ile Arg Lys Phe Ser Ile Val Gln Lys	
	705 710 715	
30	ACT CCC TTA CAA ATG AAT GGC ATC GAA GAG GAT TCT GAT GAG CCT TTA	2328
	Thr Pro Leu Gln Met Asn Gly Ile Glu Glu Asp Ser Asp Glu Pro Leu	
	720 725 730	
35	GAG AGA AGG CTG TCC TTA GTA CCA GAT TCT GAG CAG GGA GAG GCG ATA	2376
	Glu Arg Arg Leu Ser Leu Val Pro Asp Ser Glu Gln Gly Glu Ala Ile	
	735 740 745	
40	CTG CCT CGC ATC AGC GTG ATC AGC ACT GGC CCC ACG CTT CAG GCA CGA	2424
	Leu Pro Arg Ile Ser Val Ile Ser Thr Gly Pro Thr Leu Gln Ala Arg	
	750 755 760	
45	AGG AGG CAG TCT GTC CTG AAC CTG ATG ACA CAC TCA GTT AAC CAA GGT	2472
	Arg Arg Gln Ser Val Leu Asn Leu Met Thr His Ser Val Asn Gln Gly	
	765 770 775 780	
50	CAG AAC ATT CAC CGA AAG ACA ACA GCA TCC ACA CGA AAA GTG TCA CTG	2520
	Gln Asn Ile His Arg Lys Thr Thr Ala Ser Thr Arg Lys Val Ser Leu	
	785 790 795	
55	GCC CCT CAG GCA AAC TTG ACT GAA CTG GAT ATA TAT TCA AGA AGG TTA	2568
	Ala Pro Gln Ala Asn Leu Thr Glu Leu Asp Ile Tyr Ser Arg Arg Leu	
	800 805 810	
60	TCT CAA GAA ACT GGC TTG GAA ATA AGT GAA GAA ATT AAC GAA GAA GAC	2616
	Ser Gln Glu Thr Gly Leu Glu Ile Ser Glu Glu Ile Asn Glu Glu Asp	
	815 820 825	
65	TTA AAG GAG TGC CTT TTT GAT GAT ATG GAG AGC ATA CCA GCA GTG ACT	2664
	Leu Lys Glu Cys Leu Phe Asp Asp Met Glu Ser Ile Pro Ala Val Thr	
	830 835 840	

- 101 -

5	ACA TGG AAC ACA TAC CTT CGA TAT ATT ACT GTC CAC AAG AGC TTA ATT	2712
	Thr Trp Asn Thr Tyr Leu Arg Tyr Ile Thr Val His Lys Ser Leu Ile	
	845 850 855 860	
10	TTT GTG CTA ATT TGG TGC TTA GTA ATT TTT CTG GCA GAG GTG GCT GCT	2760
	Phe Val Leu Ile Trp Cys Leu Val Ile Phe Leu Ala Glu Val Ala Ala	
	865 870 875	
15	TCT TTG GTT GTG CTG TGG CTC CTT GGA AAC ACT CCT CTT CAA GAC AAA	2808
	Ser Leu Val Val Leu Trp Leu Leu Gly Asn Thr Pro Leu Gln Asp Lys	
	880 885 890	
20	GGG AAT AGT ACT CAT AGT AGA AAT AAC AGC TAT GCA GTG ATT ATC ACC	2856
	Gly Asn Ser Thr His Ser Arg Asn Asn Ser Tyr Ala Val Ile Ile Thr	
	895 900 905	
25	AGC ACC AGT TCG TAT TAT GTG TTT TAC ATT TAC GTG GGA GTA GCC GAC	2904
	Ser Thr Ser Ser Tyr Tyr Val Phe Tyr Ile Tyr Val Gly Val Ala Asp	
	910 915 920	
30	ACT TTG CTT GCT ATG GGA TTC TTC AGA GGT CTA CCA CTG GTG CAT ACT	2952
	Thr Leu Leu Ala Met Gly Phe Phe Arg Gly Leu Pro Leu Val His Thr	
	925 930 935 940	
35	CTA ATC ACA GTG TCG AAA ATT TTA CAC CAC AAA ATG TTA CAT TCT GTT	3000
	Leu Ile Thr Val Ser Lys Ile Leu His His Lys Met Leu His Ser Val	
	945 950 955	
40	CTT CAA GCA CCT ATG TCA ACC CTC AAC ACG TTG AAA GCA GGT GGG ATT	3048
	Leu Gln Ala Pro Met Ser Thr Leu Asn Thr Leu Lys Ala Gly Gly Ile	
	960 965 970	
45	CTT AAT AGA TTC TCC AAA GAT ATA GCA ATT TTG GAT GAC CTT CTG CCT	3096
	Leu Asn Arg Phe Ser Lys Asp Ile Ala Ile Leu Asp Asp Leu Leu Pro	
	975 980 985	
50	CTT ACC ATA TTT GAC TTC ATC CAG TTG TTA TTA ATT GTG ATT GGA GCT	3144
	Leu Thr Ile Phe Asp Phe Ile Gln Leu Leu Leu Ile Val Ile Gly Ala	
	990 995 1000	
55	ATA GCA GTT GTC GCA GTT TTA CAA CCC TAC ATC TTT GTT GCA ACA GTG	3192
	Ile Ala Val Val Ala Val Leu Gln Pro Tyr Ile Phe Val Ala Thr Val	
	1005 1010 1015 1020	
60	CCA GTG ATA GTG GCT TTT ATT ATG TTG AGA GCA TAT TTC CTC CAA ACC	3240
	Pro Val Ile Val Ala Phe Ile Met Leu Arg Ala Tyr Phe Leu Gln Thr	
	1025 1030 1035	
65	TCA CAG CAA CTC AAA CAA CTG GAA TCT GAA GGC AGG AGT CCA ATT TTC	3288
	Ser Gln Gln Leu Lys Gln Leu Glu Ser Glu Gly Arg Ser Pro Ile Phe	
	1040 1045 1050	
70	ACT CAT CTT GTT ACA AGC TTA AAA GGA CTA TGG ACA CTT CGT GCC TTC	3336
	Thr His Leu Val Thr Ser Leu Lys Gly Leu Trp Thr Leu Arg Ala Phe	
	1055 1060 1065	

- 102 -

	GGA CGG CAG CCT TAC TTT GAA ACT CTG TTC CAC AAA GCT CTG AAT TTA	3384
	Gly Arg Gln Pro Tyr Phe Glu Thr Leu Phe His Lys Ala Leu Asn Leu	
	1070 1075 1080	
5	CAT ACT GCC AAC TGG TTC TTG TAC CTG TCA ACA CTG CGC TGG TTC CAA	3432
	His Thr Ala Asn Trp Phe Leu Tyr Leu Ser Thr Leu Arg Trp Phe Gln	
	1085 1090 1095 1100	
10	ATG AGA ATA GAA ATG ATT TTT GTC ATC TTC TTC ATT GCT GTT ACC TTC	3480
	Met Arg Ile Glu Met Ile Phe Val Ile Phe Phe Ile Ala Val Thr Phe	
	1105 1110 1115	
15	ATT TCC ATT TTA ACA ACA GGA GAA GGA GAA GGA AGA GTT GGT ATT ATC	3528
	Ile Ser Ile Leu Thr Thr Gly Glu Gly Glu Gly Arg Val Gly Ile Ile	
	1120 1125 1130	
20	CTG ACT TTA GCC ATG AAT ATC ATG AGT ACA TTG CAG TGG GCT GTA AAC	3576
	Leu Thr Leu Ala Met Asn Ile Met Ser Thr Leu Gln Trp Ala Val Asn	
	1135 1140 1145	
25	TCC AGC ATA GAT GTG GAT AGC TTG ATG CGA TCT GTG AGC CGA GTC TTT	3624
	Ser Ser Ile Asp Val Asp Ser Leu Met Arg Ser Val Ser Arg Val Phe	
	1150 1155 1160	
30	AAG TTC ATT GAC ATG CCA ACA GAA GGT AAA CCT ACC AAG TCA ACC AAA	3672
	Lys Phe Ile Asp Met Pro Thr Glu Gly Lys Pro Thr Lys Ser Thr Lys	
	1165 1170 1175 1180	
35	CCA TAC AAG AAT GGC CAA CTC TCG AAA GTT ATG ATT ATT GAG AAT TCA	3720
	Pro Tyr Lys Asn Gly Gln Leu Ser Lys Val Met Ile Ile Glu Asn Ser	
	1185 1190 1195	
40	CAC GTG AAG AAA GAT GAC ATC TGG CCC TCA GGG GGC CAA ATG ACT GTC	3768
	His Val Lys Lys Asp Asp Ile Trp Pro Ser Gly Gly Gln Met Thr Val	
	1200 1205 1210	
45	AAA GAT CTC ACA GCA AAA TAC ACA GAA GGT GGA AAT GCC ATA TTA GAG	3816
	Lys Asp Leu Thr Ala Lys Tyr Thr Glu Gly Gly Asn Ala Ile Leu Glu	
	1215 1220 1225	
50	AAC ATT TCC TTC TCA ATA AGT CCT GGC CAG AGG GTG GGC CTC TTG GGA	3864
	Asn Ile Ser Phe Ser Ile Ser Pro Gly Gln Arg Val Gly Leu Leu Gly	
	1230 1235 1240	
55	AGA ACT GGA TCA GGG AAG AGT ACT TTG TTA TCA GCT TTT TTG AGA CTA	3912
	Arg Thr Gly Ser Gly Lys Ser Thr Leu Leu Ser Ala Phe Leu Arg Leu	
	1245 1250 1255 1260	
50	CTG AAC ACT GAA GGA GAA ATC CAG ATC GAT GGT GTG TCT TGG GAT TCA	3960
	Leu Asn Thr Glu Gly Glu Ile Gln Ile Asp Gly Val Ser Trp Asp Ser	
	1265 1270 1275	
55	ATA ACT TTG CAA CAG TGG AGG AAA GCC TTT GGA GTG ATA CCA CAG AAA	4008
	Ile Thr Leu Gln Gln Trp Arg Lys Ala Phe Gly Val Ile Pro Gln Lys	
	1280 1285 1290	

- 103 -

5	GTA TTT ATT TTT TCT GGA ACA TTT AGA AAA AAC TTG GAT CCC TAT GAA	4056
	Val Phe Ile Phe Ser Gly Thr Phe Arg Lys Asn Leu Asp Pro Tyr Glu	
	1295 1300 1305	
	CAG TGG AGT GAT CAA GAA ATA TGG AAA GTT GCA GAT GAG GTT GGG CTC	4104
	Gln Trp Ser Asp Gln Glu Ile Trp Lys Val Ala Asp Glu Val Gly Leu	
10	1310 1315 1320	
	AGA TCT GTG ATA GAA CAG TTT CCT GGG AAG CTT GAC TTT GTC CTT GTG	4152
	Arg Ser Val Ile Glu Gln Phe Pro Gly Lys Leu Asp Phe Val Leu Val	
	1325 1330 1335 1340	
	GAT GGG GGC TGT GTC CTA AGC CAT GGC CAC AAG CAG TTG ATG TGC TTG	4200
15	Asp Gly Gly Cys Val Leu Ser His Gly His Lys Gln Leu Met Cys Leu	
	1345 1350 1355	
	GCT AGA TCT GTT CTC AGT AAG GCG AAG ATC TTG CTG CTT GAT GAA CCC	4248
	Ala Arg Ser Val Leu Ser Lys Ala Lys Ile Leu Leu Leu Asp Glu Pro	
	1360 1365 1370	
20	AGT GCT CAT TTG GAT CCA GTA ACA TAC CAA ATA ATT AGA AGA ACT CTA	4296
	Ser Ala His Leu Asp Pro Val Thr Tyr Gln Ile Ile Arg Arg Thr Leu	
	1375 1380 1385	
	AAA CAA GCA TTT GCT GAT TGC ACA GTA ATT CTC TGT GAA CAC AGG ATA	4344
	Lys Gln Ala Phe Ala Asp Cys Thr Val Ile Leu Cys Glu His Arg Ile	
25	1390 1395 1400	
	GAA GCA ATG CTG GAA TGC CAA CAA TTT TTG GTC ATA GAA GAG AAC AAA	4392
	Glu Ala Met Leu Glu Cys Gln Gln Phe Leu Val Ile Glu Glu Asn Lys	
	1405 1410 1415 1420	
	GTG CGG CAG TAC GAT TCC ATC CAG AAA CTG CTG AAC GAG AGG AGC CTC	4440
30	Val Arg Gln Tyr Asp Ser Ile Gln Lys Leu Leu Asn Glu Arg Ser Leu	
	1425 1430 1435	
	TTC CGG CAA GCC ATC AGC CCC TCC GAC AGG GTG AAG CTC TTT CCC CAC	4488
	Phe Arg Gln Ala Ile Ser Pro Ser Asp Arg Val Lys Leu Phe Pro His	
	1440 1445 1450	
35	CGG AAC TCA AGC AAG TGC AAG TCT AAG CCC CAG ATT GCT GCT CTG AAA	4536
	Arg Asn Ser Ser Lys Cys Lys Ser Lys Pro Gln Ile Ala Ala Leu Lys	
	1455 1460 1465	
	GAG GAG ACA GAA GAA GAG GTG CAA GAT ACA AGG CTT TAGAGAGCAG	4582
	Glu Glu Thr Glu Glu Glu Val Gln Asp Thr Arg Leu	
40	1470 1475 1480	
	CATAAATGTT GACATGGGAC ATTTGCTCAT GGAATTGGAG CTCGTGGGAC AGTCACCTCA	4642
	TGGAATTGGA GCTCGTGGAA CAGTTACCTC TGCCTCAGAA AACAAGGATG AATTAAGTTT	4702
	TTTTTTAAAA AAGAAACATT TGGTAAGGGG AATTGAGGAC ACTGATATGG GTCTTGATAA	4762
	ATGGCTTCCT GGCAATAGTC AAATTGTGTG AAAGGTACTT CAAATCCTTG AAGATTTACC	4822
45	ACTTGTGTTT TGCAAGCCAG ATTTTCCTGA AAACCCTTGC CATGTGCTAG TAATTGGAAA	4882
50		
55		

- 104 -

GGCAGCTCTA AATGTCAATC AGCCTAGTTG ATCAGCTTAT TGTCTAGTGA AACTCGTTAA 4942
 TTTGTAGTGT TGGAGAAGAA CTGAAATCAT ACTTCTTAGG GTTATGATTA AGTAATGATA 5002
 5 ACTGGAAACT TCAGCGGTTT ATATAAGCTT GTATTCCCTT TTCTCTCCTC TCCCCATGAT 5062
 GTTTAGAAAC ACAACTATAT TGTTTGCTAA GCATTCCAAC TATCTCATTT CCAAGCAAGT 5122
 10 ATTAGAATAC CACAGGAACC ACAAGACTGC ACATCAAAAT ATGCCCCATT CAACATCTAG 5182
 TGAGCAGTCA GGAAAGAGAA CTTCCAGATC CTGGAAATCA GGGTTAGTAT TGTCCAGGTC 5242
 TACCAAAAAT CTCAATATTT CAGATAATCA CAATACATCC CTTACCTGGG AAAGGGCTGT 5302
 15 TATAATCTTT CACAGGGGAC AGGATGGTTC CCTTGATGAA GAAGTTGATA TGCCTTTTCC 5362
 CAACTCCAGA AAGTGACAAG CTCACAGACC TTTGAACTAG AGTTTAGCTG GAAAAGTATG 5422
 TTAGTGCAAA TTGTCACAGG ACAGCCCTTC TTTCCACAGA AGCTCCAGGT AGAGGGGTGTG 5482
 20 TAAGTAGATA GGCCATGGGC ACTGTGGGTA GACACACATG AAGTCCAAGC ATTTAGATGT 5542
 ATAGGTTGAT GGTGGTATGT TTTCAGGCTA GATGTATGTA CTTCATGCTG TCTACACTAA 5602
 25 GAGAGAATGA GAGACACACT GAAGAAGCAC CAATCATGAA TTAGTTTTAT ATGCTTCTGT 5662
 TTTATAATTT TGTGAAGCAA AATTTTTTCT CTAGGAAATA TTTATTTTAA TAATGTTTCA 5722
 AACATATATT ACAATGCTGT ATTTTAAAAG AATGATTATG AATTACATTT GTATAAAATA 5782
 30 ATTTTATAT TTGAAATATT GACTTTTTAT GGCACATGTA TTTTATGAA ATATTATGTT 5842
 AAAACTGGGA CAGGGGAGAA CCTAGGGTGA TATTAACCAG GGGCCATGAA TCACCTTTTG 5902
 35 GTCTGGAGGG AAGCCTTGGG GCTGATCGAG TTGTTGCCCA CAGCTGTATG ATTCCCAGCC 5962
 AGACACAGCC TCTTAGATGC AGTTCTGAAG AAGATGGTAC CACCAGTCTG ACTGTTTCCA 6022
 TCAAGGGTAC ACTGCCTTCT CAACTCCAAA CTGACTCTTA AGAAGACTGC ATTATATTTA 6082
 40 TTACTGTAAG AAAATATCAC TTGTCAATAA AATCCATACA TTTGTGT 6129

(2) INFORMATION FOR SEQ ID NO:2:

45

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1480 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

50

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

55

Met Gln Arg Ser Pro Leu Glu Lys Ala Ser Val Val Ser Lys Leu Phe

1

5

10

15

- 105 -

	Phe Ser Trp Thr Arg Pro Ile Leu Arg Lys Gly Tyr Arg Gln Arg Leu	
	20	25 30
5	Glu Leu Ser Asp Ile Tyr Gln Ile Pro Ser Val Asp Ser Ala Asp Asn	
	35	40 45
	Leu Ser Glu Lys Leu Glu Arg Glu Trp Asp Arg Glu Leu Ala Ser Lys	
	50	55 60
10	Lys Asn Pro Lys Leu Ile Asn Ala Leu Arg Arg Cys Phe Phe Trp Arg	
	65	70 75 80
	Phe Met Phe Tyr Gly Ile Phe Leu Tyr Leu Gly Glu Val Thr Lys Ala	
15	85	90 95
	Val Gln Pro Leu Leu Leu Gly Arg Ile Ile Ala Ser Tyr Asp Pro Asp	
	100	105 110
20	Asn Lys Glu Glu Arg Ser Ile Ala Ile Tyr Leu Gly Ile Gly Leu Cys	
	115	120 125
	Leu Leu Phe Ile Val Arg Thr Leu Leu Leu His Pro Ala Ile Phe Gly	
	130	135 140
25	Leu His His Ile Gly Met Gln Met Arg Ile Ala Met Phe Ser Leu Ile	
	145	150 155 160
	Tyr Lys Lys Thr Leu Lys Leu Ser Ser Arg Val Leu Asp Lys Ile Ser	
30	165	170 175
	Ile Gly Gln Leu Val Ser Leu Leu Ser Asn Asn Leu Asn Lys Phe Asp	
	180	185 190
35	Glu Gly Leu Ala Leu Ala His Phe Val Trp Ile Ala Pro Leu Gln Val	
	195	200 205
	Ala Leu Leu Met Gly Leu Ile Trp Glu Leu Leu Gln Ala Ser Ala Phe	
	210	215 220
40	Cys Gly Leu Gly Phe Leu Ile Val Leu Ala Leu Phe Gln Ala Gly Leu	
	225	230 235 240
	Gly Arg Met Met Met Lys Tyr Arg Asp Gln Arg Ala Gly Lys Ile Ser	
45	245	250 255
	Glu Arg Leu Val Ile Thr Ser Glu Met Ile Glu Asn Ile Gln Ser Val	
	260	265 270
50	Lys Ala Tyr Cys Trp Glu Glu Ala Met Glu Lys Met Ile Glu Asn Leu	
	275	280 285
	Arg Gln Thr Glu Leu Lys Leu Thr Arg Lys Ala Ala Tyr Val Arg Tyr	
	290	295 300
55	Phe Asn Ser Ser Ala Phe Phe Phe Ser Gly Phe Phe Val Val Phe Leu	
	305	310 315 320

- 106 -

	Ser Val Leu Pro Tyr Ala Leu Ile Lys Gly Ile Ile Leu Arg Lys Ile	325	330	335	
5	Phe Thr Thr Ile Ser Phe Cys Ile Val Leu Arg Met Ala Val Thr Arg	340	345	350	
	Gln Phe Pro Trp Ala Val Gln Thr Trp Tyr Asp Ser Leu Gly Ala Ile	355	360	365	
10	Asn Lys Ile Gln Asp Phe Leu Gln Lys Gln Glu Tyr Lys Thr Leu Glu	370	375	380	
	Tyr Asn Leu Thr Thr Thr Glu Val Val Met Glu Asn Val Thr Ala Phe	385	390	395	400
15	Trp Glu Glu Gly Phe Gly Glu Leu Phe Glu Lys Ala Lys Gln Asn Asn	405	410	415	
	Asn Asn Arg Lys Thr Ser Asn Gly Asp Asp Ser Leu Phe Phe Ser Asn	420	425	430	
	Phe Ser Leu Leu Gly Thr Pro Val Leu Lys Asp Ile Asn Phe Lys Ile	435	440	445	
25	Glu Arg Gly Gln Leu Leu Ala Val Ala Gly Ser Thr Gly Ala Gly Lys	450	455	460	
	Thr Ser Leu Leu Met Met Ile Met Gly Glu Leu Glu Pro Ser Glu Gly	465	470	475	480
30	Lys Ile Lys His Ser Gly Arg Ile Ser Phe Cys Ser Gln Phe Ser Trp	485	490	495	
	Ile Met Pro Gly Thr Ile Lys Glu Asn Ile Ile Phe Gly Val Ser Tyr	500	505	510	
	Asp Glu Tyr Arg Tyr Arg Ser Val Ile Lys Ala Cys Gln Leu Glu Glu	515	520	525	
40	Asp Ile Ser Lys Phe Ala Glu Lys Asp Asn Ile Val Leu Gly Glu Gly	530	535	540	
	Gly Ile Thr Leu Ser Gly Gly Gln Arg Ala Arg Ile Ser Leu Ala Arg	545	550	555	560
45	Ala Val Tyr Lys Asp Ala Asp Leu Tyr Leu Leu Asp Ser Pro Phe Gly	565	570	575	
	Tyr Leu Asp Val Leu Thr Glu Lys Glu Ile Phe Glu Ser Cys Val Cys	580	585	590	
50	Lys Leu Met Ala Asn Lys Thr Arg Ile Leu Val Thr Ser Lys Met Glu	595	600	605	
55	His Leu Lys Lys Ala Asp Lys Ile Leu Ile Leu His Glu Gly Ser Ser	610	615	620	

- 107 -

	Tyr	Phe	Tyr	Gly	Thr	Phe	Ser	Glu	Leu	Gln	Asn	Leu	Gln	Pro	Asp	Phe	625	630	635	640
5	Ser	Ser	Lys	Leu	Met	Gly	Cys	Asp	Ser	Phe	Asp	Gln	Phe	Ser	Ala	Glu	645	650	655	
	Arg	Arg	Asn	Ser	Ile	Leu	Thr	Glu	Thr	Leu	His	Arg	Phe	Ser	Leu	Glu	660	665	670	
10	Gly	Asp	Ala	Pro	Val	Ser	Trp	Thr	Glu	Thr	Lys	Lys	Gln	Ser	Phe	Lys	675	680	685	
15	Gln	Thr	Gly	Glu	Phe	Gly	Glu	Lys	Arg	Lys	Asn	Ser	Ile	Leu	Asn	Pro	690	695	700	
	Ile	Asn	Ser	Ile	Arg	Lys	Phe	Ser	Ile	Val	Gln	Lys	Thr	Pro	Leu	Gln	705	710	715	720
20	Met	Asn	Gly	Ile	Glu	Glu	Asp	Ser	Asp	Glu	Pro	Leu	Glu	Arg	Arg	Leu	725	730	735	
	Ser	Leu	Val	Pro	Asp	Ser	Glu	Gln	Gly	Glu	Ala	Ile	Leu	Pro	Arg	Ile	740	745	750	
25	Ser	Val	Ile	Ser	Thr	Gly	Pro	Thr	Leu	Gln	Ala	Arg	Arg	Arg	Gln	Ser	755	760	765	
30	Val	Leu	Asn	Leu	Met	Thr	His	Ser	Val	Asn	Gln	Gly	Gln	Asn	Ile	His	770	775	780	
	Arg	Lys	Thr	Thr	Ala	Ser	Thr	Arg	Lys	Val	Ser	Leu	Ala	Pro	Gln	Ala	785	790	795	800
35	Asn	Leu	Thr	Glu	Leu	Asp	Ile	Tyr	Ser	Arg	Arg	Leu	Ser	Gln	Glu	Thr	805	810	815	
	Gly	Leu	Glu	Ile	Ser	Glu	Glu	Ile	Asn	Glu	Glu	Asp	Leu	Lys	Glu	Cys	820	825	830	
40	Leu	Phe	Asp	Asp	Met	Glu	Ser	Ile	Pro	Ala	Val	Thr	Thr	Trp	Asn	Thr	835	840	845	
45	Tyr	Leu	Arg	Tyr	Ile	Thr	Val	His	Lys	Ser	Leu	Ile	Phe	Val	Leu	Ile	850	855	860	
	Trp	Cys	Leu	Val	Ile	Phe	Leu	Ala	Glu	Val	Ala	Ala	Ser	Leu	Val	Val	865	870	875	880
50	Leu	Trp	Leu	Leu	Gly	Asn	Thr	Pro	Leu	Gln	Asp	Lys	Gly	Asn	Ser	Thr	885	890	895	
	His	Ser	Arg	Asn	Asn	Ser	Tyr	Ala	Val	Ile	Ile	Thr	Ser	Thr	Ser	Ser	900	905	910	
55	Tyr	Tyr	Val	Phe	Tyr	Ile	Tyr	Val	Gly	Val	Ala	Asp	Thr	Leu	Leu	Ala	915	920	925	

- 108 -

Met Gly Phe Phe Arg Gly Leu Pro Leu Val His Thr Leu Ile Thr Val
930 935 940

5 Ser Lys Ile Leu His His Lys Met Leu His Ser Val Leu Gln Ala Pro
945 950 955 960

Met Ser Thr Leu Asn Thr Leu Lys Ala Gly Gly Ile Leu Asn Arg Phe
965 970 975

10 Ser Lys Asp Ile Ala Ile Leu Asp Asp Leu Leu Pro Leu Thr Ile Phe
980 985 990

15 Asp Phe Ile Gln Leu Leu Leu Ile Val Ile Gly Ala Ile Ala Val Val
995 1000 1005

Ala Val Leu Gln Pro Tyr Ile Phe Val Ala Thr Val Pro Val Ile Val
1010 1015 1020

20 Ala Phe Ile Met Leu Arg Ala Tyr Phe Leu Gln Thr Ser Gln Gln Leu
1025 1030 1035 1040

Lys Gln Leu Glu Ser Glu Gly Arg Ser Pro Ile Phe Thr His Leu Val
1045 1050 1055

25 Thr Ser Leu Lys Gly Leu Trp Thr Leu Arg Ala Phe Gly Arg Gln Pro
1060 1065 1070

30 Tyr Phe Glu Thr Leu Phe His Lys Ala Leu Asn Leu His Thr Ala Asn
1075 1080 1085

Trp Phe Leu Tyr Leu Ser Thr Leu Arg Trp Phe Gln Met Arg Ile Glu
1090 1095 1100

35 Met Ile Phe Val Ile Phe Phe Ile Ala Val Thr Phe Ile Ser Ile Leu
1105 1110 1115 1120

Thr Thr Gly Glu Gly Glu Gly Arg Val Gly Ile Ile Leu Thr Leu Ala
1125 1130 1135

40 Met Asn Ile Met Ser Thr Leu Gln Trp Ala Val Asn Ser Ser Ile Asp
1140 1145 1150

Val Asp Ser Leu Met Arg Ser Val Ser Arg Val Phe Lys Phe Ile Asp
1155 1160 1165

45 Met Pro Thr Glu Gly Lys Pro Thr Lys Ser Thr Lys Pro Tyr Lys Asn
1170 1175 1180

50 Gly Gln Leu Ser Lys Val Met Ile Ile Glu Asn Ser His Val Lys Lys
1185 1190 1195 1200

Asp Asp Ile Trp Pro Ser Gly Gly Gln Met Thr Val Lys Asp Leu Thr
1205 1210 1215

55 Ala Lys Tyr Thr Glu Gly Gly Asn Ala Ile Leu Glu Asn Ile Ser Phe
1220 1225 1230

- 109 -

	Ser Ile Ser Pro Gly Gln Arg Val Gly Leu Leu Gly Arg Thr Gly Ser	
	1235	1240 1245
5	Gly Lys Ser Thr Leu Leu Ser Ala Phe Leu Arg Leu Leu Asn Thr Glu	
	1250	1255 1260
	Gly Glu Ile Gln Ile Asp Gly Val Ser Trp Asp Ser Ile Thr Leu Gln	
	1265	1270 1275 1280
10	Gln Trp Arg Lys Ala Phe Gly Val Ile Pro Gln Lys Val Phe Ile Phe	
		1285 1290 1295
	Ser Gly Thr Phe Arg Lys Asn Leu Asp Pro Tyr Glu Gln Trp Ser Asp	
15		1300 1305 1310
	Gln Glu Ile Trp Lys Val Ala Asp Glu Val Gly Leu Arg Ser Val Ile	
	1315	1320 1325
20	Glu Gln Phe Pro Gly Lys Leu Asp Phe Val Leu Val Asp Gly Gly Cys	
	1330	1335 1340
	Val Leu Ser His Gly His Lys Gln Leu Met Cys Leu Ala Arg Ser Val	
	1345	1350 1355 1360
25	Leu Ser Lys Ala Lys Ile Leu Leu Leu Asp Glu Pro Ser Ala His Leu	
		1365 1370 1375
	Asp Pro Val Thr Tyr Gln Ile Ile Arg Arg Thr Leu Lys Gln Ala Phe	
30		1380 1385 1390
	Ala Asp Cys Thr Val Ile Leu Cys Glu His Arg Ile Glu Ala Met Leu	
	1395	1400 1405
35	Glu Cys Gln Gln Phe Leu Val Ile Glu Glu Asn Lys Val Arg Gln Tyr	
	1410	1415 1420
	Asp Ser Ile Gln Lys Leu Leu Asn Glu Arg Ser Leu Phe Arg Gln Ala	
	1425	1430 1435 1440
40	Ile Ser Pro Ser Asp Arg Val Lys Leu Phe Pro His Arg Asn Ser Ser	
		1445 1450 1455
	Lys Cys Lys Ser Lys Pro Gln Ile Ala Ala Leu Lys Glu Glu Thr Glu	
45		1460 1465 1470
	Glu Glu Val Gln Asp Thr Arg Leu	
	1475	1480

50 (2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 5635 base pairs

(B) TYPE: nucleic acid

55 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

5	CATCATCAAT AATATACCTT ATTTTGGATT GAAGCCAATA TGATAATGAG GGGGTGGAGT	60
	TTGTGACGTG GCGCGGGGCG TGGGAACGGG GCGGGTGACG TAGTAGTGTG GCGGAAGTGT	120
	GATGTTGCAA GTGTGGCGGA ACACATGTAA GCGCCGGATG TGGTAAAAGT GACGTTTTTG	180
10	GTGTGCGCCG GTGTATACGG GAAGTGACAA TTTTCGCGCG GTTTTAGGCG GATGTTGTAG	240
	TAAATTTGGG CGTAACCAAG TAATGTTTGG CCATTTTCGC GGGAAACTG AATAAGAGGA	300
15	AGTGAAATCT GAATAATTCT GTGTTACTCA TAGCGCGTAA TATTTGTCTA GGGCCGCGGG	360
	GACTTTGACC GTTTACGTGG AGACTCGCCC AGGTGTTTTT CTCAGGTGTT TTCCGCGTTC	420
	CGGGTCAAAG TTGGCGTTTT ATTATTATAG TCAGCTGACG CGCAGTGTAT TTATACCCGG	480
20	TGAGTTCCTC AAGAGGCCAC TCTTGAGTGC CAGCGAGTAG AGTTTTCTCC TCCGAGCCGC	540
	TCCGAGCTAG TAACGGCCGC CAGTGTGCTG CAGATATCAA AGTCGACGGT ACCCGAGAGA	600
25	CCATGCAGAG GTCGCCTCTG GAAAAGGCCA GCGTTGTCTC CAAACTTTTT TTCAGCTGGA	660
	CCAGACCAAT TTTGAGGAAA GGATACAGAC AGCGCCTGGA ATTGTCAGAC ATATACCAAA	720
	TCCCTTCTGT TGATTCTGCT GACAATCTAT CTGAAAAATT GGAAAGAGAA TGGGATAGAG	780
30	AGCTGGCTTC AAAGAAAAAT CCTAAACTCA TTAATGCCCT TCGGCGATGT TTTTCTGGA	840
	GATTTATGTT CTATGGAATC TTTTATATT TAGGGGAAGT CACCAAAGCA GTACAGCCTC	900
35	TCTTACTGGG AAGAATCATA GCTTCCTATG ACCCGGATAA CAAGGAGGAA CGCTCTATCG	960
	CGATTTATCT AGGCATAGGC TTATGCCTTC TCTTTATTGT GAGGACACTG CTCCTACACC	1020
	CAGCCATTTT TGGCCTTCAT CACATTGGAA TGCAGATGAG AATAGCTATG TTTAGTTTGA	1080
40	TTTATAAGAA GACTTTAAAG CTGTCAAGCC GTGTTCTAGA TAAAATAAGT ATTGGACAAC	1140
	TTGTTAGTCT CCTTTCCAAC AACCTGAACA AATTTGATGA AGGACTTGCA TTGGCACATT	1200
45	TCGTGTGGAT CGCTCCTTTG CAAGTGGCAC TCCTCATGGG GCTAATCTGG GAGTTGTTAC	1260
	AGGCGTCTGC CTTCTGTGGA CTTGGTTTCC TGATAGTCCT TGCCCTTTTT CAGGCTGGGC	1320
	TAGGGAGAAT GATGATGAAG TACAGAGATC AGAGAGCTGG GAAGATCAGT GAAAGACTTG	1380
50	TGATTACCTC AGAAATGATT GAAAACATCC AATCTGTTAA GGCATACTGC TGGGAAGAAG	1440
	CAATGGAAAA AATGATTGAA AACTTAAGAC AAACAGAACT GAAACTGACT CGGAAGGCAG	1500
55	CCTATGTGAG ATACTTCAAT AGCTCAGCCT TCTTCTTCTC AGGGTTCTTT GTGGTGTTTT	1560
	TATCTGTGCT TCCCTATGCA CTAATCAAAG GAATCATCCT CCGGAAAATA TTCACCACCA	1620
	TCTCATTCTG CATTGTTCTG CGCATGGCGG TCACTCGGCA ATTTCCCTGG GCTGTACAAA	1680

- 111 -

	CATGGTATGA	CTCTCTTGGA	GCAATAAACA	AAATACAGGA	TTTCTTACAA	AAGCAAGAAT	1740
	ATAAGACATT	GGAATATAAC	TTAACGACTA	CAGAAGTAGT	GATGGAGAAT	GTAACAGCCT	1800
5	TCTGGGAGGA	GGGATTTGGG	GAATTATTTG	AGAAAGCAAA	ACAAAACAAT	AACAATAGAA	1860
	AAACTTCTAA	TGGTGATGAC	AGCCTCTTCT	TCAGTAATTT	CTCACTTCTT	GGTACTCCTG	1920
10	TCCTGAAAGA	TATTAATTTT	AAGATAGAAA	GAGGACAGTT	GTTGGCGGTT	GCTGGATCCA	1980
	CTGGAGCAGG	CAAGACTTCA	CTTCTAATGA	TGATTATGGG	AGAACTGGAG	CCTTCAGAGG	2040
	GTAAAAATTAA	GCACAGTGGA	AGAATTTTCAT	TCTGTTCTCA	GTTTTCTCTG	ATTATGCCTG	2100
15	GCACCATTAA	AGAAAATATC	ATCTTTGGTG	TTTCTATGA	TGAATATAGA	TACAGAAGCG	2160
	TCATCAAAGC	ATGCCAACTA	GAAGAGGACA	TCTCCAAGTT	TGCAGAGAAA	GACAATATAG	2220
20	TTCTTGGAGA	AGGTGGAATC	ACACTGAGTG	GAGGTCAACG	AGCAAGAATT	TCTTTAGCAA	2280
	GAGCAGTATA	CAAAGATGCT	GATTTGTATT	TATTAGACTC	TCCTTTTGGA	TACCTAGATG	2340
	TTTTAACAGA	AAAAGAAATA	TTTGAAAGCT	GTGTCTGTAA	ACTGATGGCT	AACAAAACATA	2400
25	GGATTTTGGT	CACTTCTAAA	ATGGAACATT	TAAAGAAAGC	TGACAAAATA	TTAATTTTGC	2460
	ATGAAGGTAG	CAGCTATTTT	TATGGGACAT	TTTCAGAACT	CCAAAATCTA	CAGCCAGACT	2520
30	TTAGCTCAAA	ACTCATGGGA	TGTGATTCTT	TCGACCAATT	TAGTGCAGAA	AGAAGAAATT	2580
	CAATCCTAAC	TGAGACCTTA	CACCGTTTCT	CATTAGAAGG	AGATGCTCCT	GTCTCCTGGA	2640
	CAGAAACAAA	AAAACAATCT	TTTAAACAGA	CTGGAGAGTT	TGGGGAAAAA	AGGAAGAATT	2700
35	CTATTCTCAA	TCCAATCAAC	TCTATACGAA	AATTTTCCAT	TGTGCAAAAG	ACTCCCTTAC	2760
	AAATGAATGG	CATCGAAGAG	GATTCTGATG	AGCCTTTAGA	GAGAAGGCTG	TCCTTAGTAC	2820
40	CAGATTCTGA	GCAGGGAGAG	GCGATACTGC	CTCGCATCAG	CGTGATCAGC	ACTGGCCCCA	2880
	CGCTTCAGGC	ACGAAGGAGG	CAGTCTGTCC	TGAACCTGAT	GACACACTCA	GTTAACCAAG	2940
	GTCAGAACAT	TCACCGAAAG	ACAACAGCAT	CCACACGAAA	AGTGTCCTG	GCCCCTCAGG	3000
45	CAAACCTGAC	TGAACTGGAT	ATATATTCAA	GAAGGTTATC	TCAAGAAACT	GGCTTGGAAG	3060
	TAAGTGAAGA	AATTAACGAA	GAAGACTTAA	AGGAGTGCCT	TTTGATGAT	ATGGAGAGCA	3120
50	TACCAGCAGT	GACTACATGG	AACACATACC	TTCGATATAT	TACTGTCCAC	AAGAGCTTAA	3180
	TTTTTGTGCT	AATTTGGTGC	TTAGTAATTT	TTCTGGCAGA	GGTGGCTGCT	TCTTTGGTTG	3240
	TGCTGTGGCT	CCTTGGAAC	ACTCCTCTTC	AAGACAAAGG	GAATAGTACT	CATAGTAGAA	3300
55	ATAACAGCTA	TGCAGTGATT	ATCACCAGCA	CCAGTTCGTA	TTATGTGTTT	TACATTTACG	3360
	TGGGAGTAGC	CGACACTTTG	CTTGCTATGG	GATTCTTCAG	AGGTCTACCA	CTGGTGCATA	3420
	CTCTAATCAC	AGTGTCGAAA	ATTTTACACC	ACAAAATGTT	ACATTCTGTT	CTTCAAGCAC	3480

	CTATGTCAAC	CCTCAACACG	TTGAAAGCAG	GTGGGATTCT	TAATAGATTG	TCCAAAGATA	3540
5	TAGCAATTTT	GGATGACCTT	CTGCCTCTTA	CCATATTTGA	CTTCATCCAG	TTGTTATTAA	3600
	TTGTGATTGG	AGCTATAGCA	GTTGTGCGAG	TTTTACAACC	CTACATCTTT	GTTGCAACAG	3660
	TGCCAGTGAT	AGTGGCTTTT	ATTATGTTGA	GAGCATATTT	CCTCCAAACC	TCACAGCAAC	3720
10	TCAAACAAC	GGAATCTGAA	GGCAGGAGTC	CAATTTTCAC	TCATCTTGTT	ACAAGCTTAA	3780
	AAGGACTATG	GACACTTCGT	GCCTTCGGAC	GGCAGCCTTA	CTTTGAAACT	CTGTTCCACA	3840
15	AAGCTCTGAA	TTTACATACT	GCCAACTGGT	TCTTGTTACCT	GTCAACACTG	CGCTGGTTCC	3900
	AAATGAGAAT	AGAAATGATT	TTTGTCTCAT	TCTTCATTGC	TGTTACCTTC	ATTTCCATTT	3960
	TAACAACAGG	AGAAGGAGAA	GGAAGAGTTG	GTATTATCCT	GACTTTAGCC	ATGAATATCA	4020
20	TGAGTACATT	GCAGTGGGCT	GTAAACTCCA	GCATAGATGT	GGATAGCTTG	ATGCGATCTG	4080
	TGAGCCGAGT	CTTTAAGTTC	ATTGACATGC	CAACAGAAAG	TAAACCTACC	AAGTCAACCA	4140
	AACCATACAA	GAATGGCCAA	CTCTCGAAAG	TTATGATTAT	TGAGAATTCA	CACGTGAAGA	4200
25	AAGATGACAT	CTGGCCCTCA	GGGGGCCAAA	TGACTGTCAA	AGATCTCACA	GCAAAATACA	4260
	CAGAAGGTGG	AAATGCCATA	TTAGAGAACA	TTTCCTTCTC	AATAAGTCCT	GGCCAGAGGG	4320
30	TGGGCCTCTT	GGGAAGAACT	GGATCAGGGA	AGAGTACTTT	GTTATCAGCT	TTTTTGAGAC	4380
	TACTGAACAC	TGAAGGAGAA	ATCCAGATCG	ATGGTGTGTC	TTGGGATTCA	ATAACTTTGC	4440
35	AACAGTGGAG	GAAAGCCTTT	GGAGTGATAC	CACAGAAAGT	ATTTATTTTT	TCTGGAACAT	4500
	TTAGAAAAAA	CTTGGATCCC	TATGAACAGT	GGAGTGATCA	AGAAATATGG	AAAGTTGCAG	4560
	ATGAGGTTGG	GCTCAGATCT	GTGATAGAAC	AGTTTCCTGG	GAAGCTTGAC	TTTGTCTTGG	4620
40	TGGATGGGGG	CTGTGTCCTA	AGCCATGGCC	ACAAGCAGTT	GATGTGCTTG	GCTAGATCTG	4680
	TTCTCAGTAA	GGCGAAGATC	TTGCTGCTTG	ATGAACCCAG	TGCTCATTTG	GATCCAGTAA	4740
45	CATACCAAAT	AATTAGAAGA	ACTCTAAAAC	AAGCATTGTC	TGATTGCACA	GTAATTCTCT	4800
	GTGAACACAG	GATAGAAGCA	ATGCTGGAAT	GCCAACAATT	TTTGGTCATA	GAAGAGAACA	4860
	AAGTGCGGCA	GTACGATTCC	ATCCAGAAAC	TGCTGAACGA	GAGGAGCCTC	TTCCGGCAAG	4920
50	CCATCAGCCC	CTCCGACAGG	GTGAAGCTCT	TTCCCCACCG	GAAGTCAAGC	AAGTGCAAGT	4980
	CTAAGCCCCA	GATTGCTGCT	CTGAAAGAGG	AGACAGAAGA	AGAGGTGCAA	GATACAAGGC	5040
55	TTTAGAGAGC	AGCATAAATG	TTGACATGGG	ACATTTGCTC	ATGGAATTGG	AGGTAGCGGA	5100
	TTGAGGTACT	GAAATGTGTG	GGCGTGGCTT	AAGGGTGGGA	AAGAATATAT	AAGGTGGGGG	5160
	TCTCATGTAG	TTTTGTATCT	GTTTTGCAGC	AGCCGCCGCC	ATGAGCGCCA	ACTCGTTTGA	5220

TGGAAGCATT GTGAGCTCAT ATTTGACAAC GCGCATGCCC CCATGGGCGG GGGTGCCTCA 5280
 GAATGTGATG GGCTCCAGCA TTGATGGTCG CCCCCTCCTG CCCGCAAAC CTACTACCTT 5340
 5 GACCTACGAG ACCGTGTCTG GAACGCCGTT GGAGACTGCA GCCTCCGCCG CCGCTTCAGC 5400
 CGCTGCAGCC ACCGCCCGCG GGATTGTGAC TGACTTTGCT TTCCTGAGCC CGCTTGCAAG 5460
 CAGTGCAGCT TCCCCTTCAT CCGCCCGCGA TGACAAGTTG ACGGCTCTTT TGGCACAATT 5520
 10 GGATTCTTTG ACCCGGGAAC TTAATGTCGT TTCTCAGCAG CTGTTGGATC TGCGCCAGCA 5580
 GGTTTCTGCC CTGAAGGCTT CCTCCCCTCC CAATGCGGTT TAAACATAA ATAAA 5635

15 (2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 36 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

ACTCTTGAGT GCCAGCGAGT AGAGTTTTCT CCTCCG 36

(2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 29 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

GCAAAGGAGC GATCCACACG AAATGTGCC 29

(2) INFORMATION FOR SEQ ID NO:6:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 24 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

CTCCTCCGAG CCGCTCCGAG CTAG

24

(2) INFORMATION FOR SEQ ID NO:7:

5

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 31 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

10

(ii) MOLECULE TYPE: cDNA

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

CCAAAAATGG CTGGGTGTAG GAGCAGTGTC C

31

20 (2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 34 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

25

(ii) MOLECULE TYPE: cDNA

30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

CGGATCCTTT ATTATAGGGG AAGTCCACGC CTAC

34

35

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 32 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

40

(ii) MOLECULE TYPE: cDNA

45

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

50 CGGGATCCAT CGATGAAATA TGACTACGTC CG

32

Claims

1. An adenovirus-based gene therapy vector comprising the genome of an adenovirus 2 serotype in which the Ela and Elb regions of the genome, which are involved in early stages
5 of viral replication, have been deleted and replaced by genetic material of interest.
2. The adenovirus-based gene therapy vector of claim 1, wherein the genetic material of interest is DNA encoding cystic fibrosis transmembrane conductance regulator
- 10 3. The adenovirus-based gene therapy vector of claim 1 further comprising PGK promoter operably linked to the genetic material of interest.
4. The adenovirus-based gene therapy vector of claim 2 having substantially the same nucleotide sequence as shown in Table II (SEQ ID NO:3).
15
5. An adenovirus-based gene therapy vector comprising adenovirus inverted terminal repeat nucleotide sequences and the minimal nucleotide sequences necessary for efficient replication and packaging and genetic material of interest.
- 20 6. The adenovirus-based gene therapy vector of claim 5 having the adenovirus 2 sequences shown in Figure 17.
7. The adenovirus-based gene therapy vector of claim 5 further comprising PGK promoter operably linked to the genetic material of interest.
25
8. The adenovirus-based gene therapy vector of claim 5 in which the genetic material of interest is selected from the group consisting of DNA encoding: cystic fibrosis transmembrane conductance regulator, Factor VIII, and Factor IX.
- 30 9. An adenovirus-based gene therapy vector comprising an adenovirus genome which has been deleted for all E4 open reading frames, except open reading frame 6, and additionally comprising genetic material of interest.
10. The adenovirus-based gene therapy vector of claim 9 further comprising PGK
35 promoter operably linked to the genetic material of interest.
11. The adenovirus-based gene therapy vector of claim 9 in which the Ela and Elb regions of the genome, which are involved in early stages of viral replication, have been deleted.

12. The adenovirus-based gene therapy vector of claim 9 in which the E3 region has been deleted.
13. An adenovirus-based gene therapy vector comprising an adenovirus genome which
5 has been deleted for all E4 open reading frames, except open reading frame 3, and additionally comprising genetic material of interest.
14. The adenovirus-based gene therapy vector of claim 13 in which the Ela and Elb
10 regions of the genome, which are involved in early stages of viral replication, have been deleted.
15. The adenovirus-based gene therapy vector of claim 13 further comprising PGK promoter operably linked to the genetic material of interest.
- 15 16. The adenovirus-based gene therapy vector of claim 13 in which the E3 region has been deleted.
17. A method for treating or preventing cystic fibrosis in a patient comprising
administering to the pulmonary airways of the patient, a gene therapy vector comprising
20 DNA encoding cystic fibrosis transmembrane conductance regulator.
18. The method of claim 17 wherein the gene therapy vector is an adenovirus-based gene therapy vector comprising the genome of an adenovirus 2 serotype in which the Ela and Elb
25 regions of the genome, which are involved in early stages of viral replication, have been deleted and replaced by DNA encoding cystic fibrosis transmembrane conductance regulator.
19. The method of claim 17 wherein the gene therapy vector further comprises PGK promoter operably linked to the DNA encoding cystic fibrosis transmembrane conductance
30 regulator.
20. The method of claim 17 wherein the gene therapy vector is an adenovirus-based gene therapy vector comprising adenovirus inverted terminal repeats and the minimal sequences necessary for efficient replication and packaging and DNA encoding cystic fibrosis
35 transmembrane conductance regulator.
21. The method of claim 20 wherein the gene therapy vector further comprises PGK promoter operably linked to the DNA encoding cystic fibrosis transmembrane conductance regulator.

22. The method of claim 17 wherein the gene therapy vector is an adenovirus-based gene therapy vector comprising an adenovirus genome which has been deleted for all E4 open reading frames, except open reading frame 6, and additionally comprising DNA encoding
5 cystic fibrosis transmembrane conductance regulator.

23. The method of claim 22 wherein the gene therapy vector further comprises PGK promoter operably linked to the DNA encoding cystic fibrosis transmembrane conductance regulator.

10

24. The method of claim 17 wherein the gene therapy vector is an adenovirus-based gene therapy vector comprising an adenovirus genome which has been deleted for all E4 open reading frames, except open reading frame 6, and has been deleted for the Ela and Elb regions of the genome, which are involved in early stages of viral replication, and additionally
15 comprising DNA encoding cystic fibrosis transmembrane conductance regulator.

25. The method of claim 24 wherein the gene therapy vector further comprises PGK promoter operably linked to the DNA encoding cystic fibrosis transmembrane conductance regulator.

1/50

PARTIAL cDNA CLONES OF THE CFTR GENE




Figure 1

2/50

STRATEGY FOR CONSTRUCTING pKK- CFTR1




Figure 2

3/50

CONSTRUCTION OF THE pKK- CFTR2 PLASMID




Figure 3

4/50

STRATEGY FOR CONSTRUCTING THE pSC- CFTR2 PLASMID




Figure 4

5/50




Figure 5

6/50

```

S          bp 1716
P          |
h          |-----x-----xSynthetic Intronx-----x-----x
l          |
|-----1195RG-----
CCAACTAGAAAGAGGTAAGGGGCTCACCAGTTCAAAATCTGAAGTGGAGACAGGAC
GTACGGTTGATCTTCTCCATTCCCCGAGTGGTCAAGTTTTAGACTTCACCTCTGTCCTG
<-----1198RG-----
                                     bp 1717
=====|
                                     |
----->|-----
CTGAGGTGACAAATGACATCTACTCTGACATTCTCTCCTCAGGACATCTCCAAGTTTGCAG
GACTCCACTGTTACTGTAGATGAGACTGTAAGAGAGGAGTCCTGTAGAGGTTCAAACGTC
-----|<-----1197RG-----
                                     H
                                     i
                                     n
                                     c
                                     I
                                     I
-----1196RG----->
AGAAAGACAAATATAGTTCTTGGAGAAAGGTGGATCACAAGTGGAGGTC
TCTTTCTGTTATATCAAGAACCTCTTCCACCTTAGTGTGACTCACCTCCAG
-----|

```

Figure 6

7/50

CONSTRUCTION OF THE pKK- CFTR3 cDNA




Figure 7A

8/50

CONSTRUCTION OF THE pKK- CFTR3 CLONE (cont'd.)





Figure 7B

9/50

MAP OF pKK- CFTR3








-  CFTR coding region
-  CFTR noncoding region
-  85 bp intron
-  T11-derived non-CFTR DNA
-  pKK- 223- 3

Figure 8




Figure 9

11/50





Figure 10B

Figure 10A

12/50



13/50

Figure 12A

Figure 12B




Figure 12C

Figure 12D




Figure 13

FIGURE 1
MAP OF VECTOR




Figure 14




Figure 15

17/50




Figure 16

18/50




Figure 17

19/50




Figure 18A




Figure 18B




Figure 19




Figure 20A




Figure 20B




Figure 21





Figure 22

ANTIBODY TITERS






Figure 24




Figure 25

27/50



Figure 26




Figure 27




Figure 28A

Figure 28B




Figure 29A




Figure 29B

2 min




Figure 30A




Figure 30B




Figure 30C




Figure 30D




Figure 30E




Figure 30F






Figure 31



- ① Adenovirus Type 2 packaging signal and E1 enhancer Region
- ② Adenovirus Type 2 major Late Promoter and Tri-partite leader
- ③ SV40 T-antigen Nuclear Localization Signal
- ④ SV40 Poly Adenylation Signal

PAVII




- ① Adenovirus Type 2 packaging signal and E1 enhancer Region
- ② Adenovirus Type 2 major Late Promoter and Tri-partite leader
- ③ SV40 T-antigen Nuclear Localization Signal
- ④ SV40 Poly Adenylation Signal

PAV I/II LEC



- ⑤ EMC VIRUS Internal Ribosomal entry site - for polycistronic Translation

PAVI Cloning Cassette



Expression Cassette




Figure 32

36/50




Figure 33

Adenovirus Vector AD2-ORF6/PGK-CFTR




Figure 34




Figure 35

Figure 36 C




Figure 36D



Figure 36A



Figure 36B




Figure 37A




Figure 37C



Figure 37B



Figure 37D




Figure 38C



Figure 38D





Figure 38A



Figure 38B



CLINICAL SIGNS MONKEY C					AGE 7 YEARS
DATE	EXAMINATION	HEART RATE (beats/min)	RESP RATE (breath/min)	TEMPERATURE (Celsius)	WEIGHT (Kg)
5/11/93	NORMAL	112	16	37.8	6.4
5/11/93	INFECTION				
5/14/93	NORMAL	98	14	38.1	
5/18/93	NORMAL	104	16	38.3	
6/4/93	NORMAL	108	16	38.2	
6/18/93	NORMAL	112	16	38.4	
6/24/93	NORMAL	116	18	38.8	
6/24/93	INFECTION				
16/28/93	NORMAL	104	18	37.9	
7/5/93	granulation	116	16	37.4	
7/12/93	NORMAL	114	20	38.3	
9/17/93	NORMAL	108	16	38.3	7

Figure 39A

CLINICAL SIGNS MONKEY D					AGE 7 YEARS
DATE	EXAMINATION	HEART RATE (beats/min)	RESP RATE (breath/min)	TEMPERATURE (Celsius)	WEIGHT (Kg)
5/11/93	NORMAL	108	18	38.3	6.25
5/11/93	INFECTION				
5/14/93	NORMAL	100	20	38.4	
5/18/93	NORMAL	98	20	38.4	
6/4/93	NORMAL	106	18	37.9	
6/18/93	NORMAL	100	19	38.4	
6/24/93	NORMAL	106	16	37.8	
6/24/93	INFECTION				
16/28/93	NORMAL	104	16	37.4	
7/5/93	NORMAL	102	14	38.8	
7/12/93	granulation	114	16	38	
9/17/93	NORMAL	104	16	38.3	6.4

Figure 39B

CLINICAL SIGNS MONKEY E					AGE 11 YEARS
DATE	EXAMINATION	HEART RATE (beats/min)	RESP RATE (breath/min)	TEMPERATURE (Celsius)	WEIGHT (Kg)
5/11/93	NORMAL	120	18	28.3	10
5/11/93	INFECTION				
5/14/93	NORMAL	112	20	37.9	
5/18/93	NORMAL	108	22	38.4	
6/4/93	NORMAL	112	20	38.3	
6/18/93	NORMAL	106	20	38.3	
6/24/93	NORMAL	108	18	38.9	
6/24/93	INFECTION				
16/28/93	NORMAL	112	20	38	
7/5/93	NORMAL	106	22	38.3	
7/12/93	NORMAL	114	16	38	
9/17/93	NORMAL	114	16	38.3	8.75

Monkey C

Clinical Lab Results From Monkey C									
DATE	11-May	14-May	18-May	4-Jun	18-Jun	24-Jun	24-Jun	12-Jul	17-Sep
WBC/mm ³	6.7	9	8.9	7.1	7.9	7.3		10.6	8.1
NEUT/mm ³	1850	3990	3060	1480	3550	3450		2210	3950
LYMP/mm ³	4460	4220	4770	4780	3640	2670		7270	3770
MONO/mm ³	120	520	600	360	420	550		480	340
EOS/mm ³	30	110	190	120	80	400		250	70
HEMOG. gr/dl	12.2	12	12.6	12.8	14	13.5		13.7	13.9
HEMATOCR. %	38	38	42	41	45	39		46	43
PLAT k/mm ³	311	319	343	338	308	281		324	432
ESR	<1	1	1	1	0	<1		<1	<1
NA mEq/l	149	148	147		151	147		149	153
K mEq/l	3.6	3.6	2.6		3.8	3.1		3.4	3.6
Cl mEq/l	111	106	107		112	108		109	113
CO ₂ mEq/l	19	20	20		22	21		19	19
BUN mg/dl	11	18	11		14	13		16	23
CREAT mg/dl	1.1	1	1.2		1.1	1		1.1	1.2
GLUCOSE mg/dl	68	56	81		67	87		74	58
ALB gr/dl	4.7	4.3	4.7		4.9	4.2		4.5	4.5
T. PROT. gr/dl	7.3	6.7	7.1		7.4	6.9		7.1	7.4
CALCIUM mg/dl	10	9.3	9.9		10.2	9		10.1	9.5
PO ₄ mg/dl	3.3	5.9	5.7		2.9	5		3.7	3.4
ALK. PH IU/l	117	376	375		117	76		116	164
TOT BIL mg/dl	0.3	0.2	0.2		0.2	0.1		0.2	0.3
AST IU/l	38	37	45		28	25		45	34
LDH IU/l	601	599	740		277	408		458	220
URIC Ac mg/dl	0.1	0.1	<0.1		0.1	0.1		<0.1	0.1

Figure 40A

44/50

Monkey D

Clinical Lab Results From Monkey D									
DATE	11-May	14-May	18-May	4-Jun	18-Jun	24-Jun	24-Jun	12-Jul	17-Sep
WBC/mm ³	7	4.2	9.9	6.7	9.1	6.9		9.4	8.3
NBUT/mm ³	2860	1980	3060	1090	6230	1740			3180
LYMP/mm ³	3660	4180	6100	4770	1820	4750			3230
MONO/mm ³	160	410	340	500	500	190			670
EOS/mm ³	50	150	210	110	240	130			210
HEMOG. gr/dl	10.9	13.7	14.7	13.6	13.9	13.6			14.5
HEMATOCR.%	35	42	49	44	43	43		44	47
PLAT k/mm ³	268	277	413	369	265	300		284	348
ESR	1	2	<1	1	0	<1		<1	<1
							S E C O N D I N F E C T I O N		
NA mEq/l	147	150	150		149	147		148	148
K mEq/l	3.5	3.5	3.6		3.5	3.4		3.5	3
Cl mEq/l	109	106	110		111	108		109	109
CO ₂ mEq/l	19	20	20		23	20		19	16
BUN mg/dl	19	18	20		10	16		18	12
CREAT mg/dl	1.1	1	1.1		1.1	1		1	1
GLUCOSE mg/dl	65	81	72		92	78		66	88
ALB gr/dl	4.3	4.7	5.2		4.2	4.6		4.5	4.7
T. PROT. gr/dl	6.6	7.4	7.8		6.8	6.8		7.1	7.6
CALCIU. mg/dl	9.3	10.1	10.4		9.6	9		10.3	9.5
PO ₄ mg/dl	6.2	3.5	3.6		2.8	5		5.6	4.7
ALK. PH IU/l	426	104	116		82	337		328	101
TOT BIL. mg/dl	0.1	0.3	0.2		0.2	0.1		0.1	0.2
AST IU/l	29	32	103		55	27		25	21
LDH IU/l	520	496	912		768	615		252	227
URIC Ac mg/dl	0.1	<0.1	<0.1		0.1	0.1		<0.1	0.1

Figure 40B

45/50

Monkey E

Clinical Lab Results From Monkey E

DATE	11-May	13-May	14-May	18-May	4-Jun	18-Jun	24-Jun	24-Jun	12-Jul	17-Sep
WBC/mm ³	8.7				5.3	8.8	8.6		6.9	8.1
NEUT/mm ³	4850				3210	4480	2040			2592
LYMP/mm ³	3060				1510	3360	5610			5265
MONO/mm ³	120				280	350	460			182
EOS/mm ³	30				150	80	170			81
HEMOG. gr/dl	12.9				13.7	12.6	12.4		13.8	13.9
HEMATOCR. %	40				42	41	38		44	43
PLAT k/mm ³	291				287	291	300		269	432
ESR	1				1	0	<1		<1	<1
F I R S T I N F E C T I O N										
NA mEq/l	148				147	148	149		148	150
K mEq/l	3				2.6	3.7	3.6		3.1	3.8
Cl mEq/l	110				107	110	111		109	110
CO ₂ mEq/l	16				20	22	23		21	20
BUN mg/dl	8				11	15	13		14	17
CREAT mg/dl	1.1				1.2	1.1	1		1	1.2
GLUCOSE mg/dl	115				102	86	65		87	69
ALB gr/dl	4				4.4	4.5	4.8		4	4.5
T. PROT, gr/dl	6.7				7.1	7	7.3		6.8	7
CALCIUM mg/dl	9.3				9.4	9.8	9.7		9.7	9.4
PO ₄ mg/dl	3.5				4.2	5.1	3.3		4.6	4.1
ALK. PH IU/l	68				90	393	116		75	355
TOT BIL mg/dl	0.2				0.3	0.1	0.2		0.2	2
AST IU/l	32				47	27	28		28	24
LDH IU/l	416				571	277	481		247	200
URIC Ac mg/dl	0.1				<0.1	0.1	0.1		<0.1	<0.1
S E C O N D I N F E C T I O N										

Figure 40C

46/50

CYTOLOGY MONKEY C

DATE	5/11/93	5/11/93	5/18/93	6/4/93	6/18/93	6/24/93	6/24/93	6/28/93	9/17/93
LEFT NOSTRIL									
Sq. Epith.	88	F	78	63	72	74	S	B	89
Resp. Epith.	30	I	18	34	24	25	E	I	30
Neutrophils	1	R	2	3	2	0	C	O	0
Lymphocytes	1	S	2	0	1	1	O	P	0
Eosinophils	0	T	0	0	1	0	N	S	1
							D	Y	

CYTOLOGY MONKEY D

DATE	5/11/93	5/11/93	5/18/93	6/4/93	6/18/93	6/24/93	6/24/93	7/5/93	9/17/93
LEFT NOSTRIL									
Sq. Epith.	60	F	60	72	72	84	S	B	73
Resp. Epith.	39	I	39	26	25	14	E	I	25
Neutrophils	1	R	1	0	1	2	C	O	2
Lymphocytes	0	S	2	2	1	0	O	P	0
Eosinophils	0	T	0	0	1	0	N	S	0
							D	Y	

CYTOLOGY MONKEY E

DATE	5/11/93	5/11/93	5/18/93	6/4/93	6/18/93	6/24/93	6/24/93	7/12/93	9/17/93
LEFT NOSTRIL									
Sq. Epith.	60	F	60	72	72	84	S	B	73
Resp. Epith.	39	I	39	26	25	14	E	I	25
Neutrophils	1	R	1	0	1	2	C	O	2
Lymphocytes	0	S	2	2	1	0	O	P	0
Eosinophils	0	T	0	0	1	0	N	S	0
							D	Y	

Figure 41

47/50




Figure 42

48/50




Figure 43



Figure 44

50/50

NEUTRALIZING ANTIBODIES •

